

**STUDY OF FACTORS RESPONSIBLE FOR ACADEMIC BACKWARDNESS IN  
SCHOOL GOING CHILDREN****\*Dr. Sudhir Mehta**MBBS, DCH, DNB (PEDIATRICS) Department of Pediatrics, Sri Aurobindo Medical College and PG Institute,  
Indore, Madhya Pradesh, India.**\*Author for Correspondence: Dr. Sudhir Mehta**MBBS, DCH, DNB (PEDIATRICS) Department of Pediatrics, Sri Aurobindo Medical College and PG Institute, Indore, Madhya Pradesh,  
India.

Article Received on 23/12/2015

Article Revised on 12/01/2016

Article Accepted on 02/02/2016

**ABSTRACT**

**Background:** Academic backwardness is a problem with far reaching consequences. This study was undertaken with the aim of correlating IQ obtained through WI scale for children with poor academic performance and also to ascertain various factors associated with academic backwardness. **Methods:** All children in the age group 10 to 16 years brought with complaints of poor school performance were included in the study. They were evaluated with a thorough relevant history, physical examination. IQ assessment was done using Wechsler Intelligence Score for Children (WISC). Results: 36% (n=18) children had subnormal IQs, the majority having normal IQ. Behavioral disorders were the single largest group of associated factors followed by chronic medical illnesses. Among the various perinatal and postnatal factors, history of delayed development was significantly associated with subnormal IQ. Behavioural disorders were present in children performing poorly at school. **Conclusions:** Mental subnormality accounts for only a minority of children with academic backwardness. Behavioral disorders can either be the cause or effect of poor scholastic performance.

**KEYWORDS:** Wechsler intelligence score, Mental sub normality, Academic backwardness.**INTRODUCTION**

Academic performance and behavior of children represent final common pathway, the convergence of many forces including interaction between cognitive strengths and deficits, environmental and cultural factors, temperament, educational experience and intrinsic resiliency. The cause of academic backwardness may be anywhere; right from the home to the grading system or even the examiner. Individual intelligence tests are the best available predictors of academic success.<sup>[1]</sup> However, the highest correlations that are achieved between IQ and achievement grade equivalents range from 0.5 to 0.6.<sup>[2]</sup> This study was undertaken to elucidate the relationship between them and to find the factors associated with poor school performance.

**METHODS**

Children between 10-16 years of age presenting with poor school performance. A directed paediatric evaluation was done including; detailed history with respect to the complaints, detailed perinatal, past medical history including history of recurrent otitis media, meningitis, encephalitis, epilepsy, major head trauma, CNS irradiation, chronic medical illness, developmental History, family history of learning/neuro developmental disorders, speech and language disorders, genetic

syndromes or heritable diseases, external stress/socio emotional preoccupation: history of parental divorce, conflict or depression, family turmoil or violence, death/ illness of a family member, Child-school mismatch, Socio-economic class using Kuppusswamy scale. A thorough physical examination for evidence of chronic medical conditions and genetic syndrome phenotype. IQ was recorded using the Wechsler Intelligence scale for children by a trained clinical psychologist. Visual acuity, audiometry, iron studies were done in all to exclude cognitive abnormalities due to iron deficiency anemia. Statistical analysis was carried out with the help of the SPSS statistical package. The data was collected in a pre-planned proforma and analyzed. Correlation of academic performance with intelligence quotients was done using the Pearson's correlation coefficient. X = Independent variable i.e. IQ.

Y= Dependent variable i.e. marks obtained in previous school examination. The significance of the correlation coefficient thus obtained was tested by dividing the observed r by the standard error. The analysis was carried out using SPSS statistical package.

## RESULTS

The study group comprised of 50 children, 26 male and 24 female of age group between 10-16 years. 32(64%) had normal IQ and 18(36%) were having subnormal IQ. Of the 18 children with subnormal IQ; 17 had an IQ between 50-69 (mild MR) and one had IQ between 35-49 (moderate MR). Of the 32 children with normal intelligence quotients, half of them had average IQ and the half had borderline IQ.

30 children were born at term and 20 were preterm at birth. Of the 30 full term children, 22 had normal IQ and 8 had mental sub normality, while among the 20 preterm children, 13 had normal IQ and 7 had mental sub normality.

There were 3 children with history of perinatal insults and 2 with history of postnatal insults. There were 12 children with associated medical disorders of which 8 had epilepsy and were on antiepileptic drugs, 2 had cerebral palsy (spastic diplegia).

A history of developmental delay was obtained in 5 out of the total children with normal intelligence; in 10 out of the 16 children with borderline intelligence and 11 out of the 16 children with mild mental retardation.

## DISCUSSION

Of the total 50 children we studied, 44(77.19%) were male and 13(23.36%) female. 70% of the children were 10-13 years old and 30% 14-16 years old. This clustering in the early age group may be due to the relatively high prevalence of scholastic problems at the beginning of middle school as recorded by other studies also.<sup>[3]</sup> Literature shows a divided opinion on the causes of poor school performance and the predictors of academic success. No significant correlation of marks obtained in school examinations with IQ was found in our study, ( $r=0.185$ ) while earlier studies recorded it in the range of 0.5 to 0.6.<sup>[1]</sup>

Associated factors were found in only 65% of children with normal IQs; leaving almost one third i.e. 36% with no identifiable cause. Subtle micronutrient deficiencies or other environmental problems which could not be made out in a few visits, may account for them. Many studies have proved the role of micronutrient deficiency exclusive of other factors.<sup>[3,4]</sup> Behavioral disorders were found in 33%; medical problems in 20%; dyslexia in 2%; and family problems in 2%.

Among the children with subnormal intelligence, 26% had no other associated factor; behavioral disorders in 20%, medical problems were found in 18%; and family problems in 6%.

The single largest group (45%) was behavioral disorders; which are most likely to be due to a complex interaction of the personality and biological make-up of the child with both family relationships and the environment at

school. This association has also been described by other researchers.<sup>[5-7]</sup> It is also difficult to say which comes first, poor academic progress may cause emotional upsets through frustration. Another study found that behavioral maladjustment in many domains is strongly associated with learning difficulties even after intelligence and socioeconomic factors are controlled.<sup>[7]</sup> It has also been seen in recent years that improving survival of low birth weight infants is accompanied by a higher prevalence of behavior disorders.<sup>[8,9]</sup> In our study the majority of children (62%) with behavioral disorders were born with low birth weight.

The majority of children with behavioral disorders in the present study were having normal intelligence but this association was not statistically significant when compared to the same in mentally retarded. The next largest group was that of chronic illness which affect learning when students experience sensory, physical or other health related impairments or when the illness interferes with school. 24% had associated medical problems, out of which, the majority having epilepsy (on antiepileptic drugs). 60%. Almost 12% had chronic headaches<sup>[10]</sup>, 10% had spastic diplegia 10% had JRA and 8% had hypothyroidism. Epilepsy, its association with mental sub normality and its effect on school performance had always been an area of controversy.<sup>[11-13]</sup> Only 48% of the children with normal intelligence quotients had a history of developmental delay, while 84% of the children with mental subnormality had history of developmental delay, this difference was found to be statistically significant ( $p<0.01$ ). A strong and persistent connection between cognitive abilities and academic achievement has been documented by research.<sup>[14,15]</sup>

In the present study, the majority of children i.e. 85% belonged to the Kuppaswamy class 2-the upper middle class. The fact that we were dealing with a group of the society who were concerned with the education of their children and it's a hospital based study. The effect of poor socioeconomic conditions on scholastic backwardness is studied in different studies though in our study we cannot draw any inference.

## CONCLUSIONS

Factors other than mental retardation are associated with scholastic backwardness, in majority of children. The single largest group was of behavioral disorders, with majority having normal IQs; but this association was not statistically significant. Associated medical conditions are important in a large number of children performing poorly at school in the present study.

## Funding

No funding sources.

## Conflict of interest

None declared.

**REFERENCES**

1. Marcel Kinsbourne, William D. Graf. Disorders of mental development in child neurology. In: John H. Menkes, Harvey B. Sarnat, eds. A Book. 6<sup>th</sup> ed. Philadelphia: Lippincott Williams and Wilkins, 2002; 1156-1157.
2. Yule W, Rutter M. Reading and intelligence. In: Knights R, Bakker J, eds. Neuropsychology of Language Disorders. Theoretical Approaches. 1<sup>st</sup> ed. Baltimore: University Park Press, 1976.
3. Sudha Chaudhri, Madhumati, Anjali Chitale, Anand Pandit, Mahendra Hoge. Pune low birth weight study-cognitive abilities and educational performance at 12 y. *Indian Pediatr.*, 2004; 41: 121-8.
4. Lui J, Rane A, Venebles PH, Dalais C, Mednik SA. Malnutrition at age 3 years and lower cognitive ability at age 11 years: independence from psychosocial adversity. *Arch Pediatr Adolesc Med.*, 2003; 157: 593-600.
5. Byrd RS, Weitzman M, Aunger P. Increased behavioral problems associated with delayed school entry and delayed school progress. *Pediatrics.*, 1997; 100: 654-61.
6. Mostafa. A. Abdolfotouh. Behaviour disorders among urban school boys in Southwest Saudi Arabia. *East Mediterr Health J.*, 1997; 3(2): 274-83.
7. Lin Y-Q, Ebrahim GJ. Frequency of behaviour disorder and related factors in school children in Lusaka. *J Trop Pediatr.*, 1991; 37: 303-9.
8. Margot Prior, Diana Smart, Ann Sanson, Frank Berklaid. Relationships between learning difficulties and psychological problems in preadolescent children from a longitudinal sample. *J Am Acad Child Adolesc Psychiatry.*, 1999; 38(4): 429-36.
9. Pharoah PO, Stevenson CJ, Cooke RW, Stevenson RC. Prevalence of behaviour disorders in low birth weight infants. *Arch Dis Childhood.*, 1994; 70: 271-4.
10. Mehta S. Study of various social and demographic variables associated with primary headache disorders in 500 school going children of central India. *J Pediatr Neurosci*, 2015; 10: 13-7.
11. Ellen Bery JH, Hotz DG, Nelson KB. Do seizures in children cause intellectual deterioration? *NEJM.*, 1986; 314: 1085-8.
12. Sunil Karande, Madhuri Kulkarni. Poor school performance. *Indian J Pediatr.*, 2005; 72: 961-7.
13. Ibekwe RC, Ojinnaka NC, Iloeje SO. Academic performance of school children with epilepsy. *West Afr J Med.*, 2008; 27(2): 74-7.
14. Demonet JF, Taylor MJ, Chaix. Developmental dyslexia. *Lancet.*, 2004; 363: 1451-60.
15. Karande S, Kulkarni A. Specific learning disability: the invisible handicap. *Indian Pediatr.*, 2005; 45: 315-9.