

**EFFECT OF CORIANDRUM OIL ON THE RNA LEVELS IN THE OVARY OF
CORCYRA CEPHALONICA*****M. Madhavi**

Assistant Professor, Department of Zoology, Nizam College, Osmania University Hyderabad-500001.

***Corresponding Author: Dr. M. Madhavi**

Assistant Professor, Department of Zoology, Nizam College, Osmania University Hyderabad-500001.

Article Received on 31/01/2017

Article Revised on 21/02/2017

Article Accepted on 14/03/2017

ABSTRACT

Corcyra cephalonica is a threat to agricultural crop produces infesting cereals, and many other food products, hence an attempt was made to control the stored products pest by using medicinal plant extract Coriandrum oil. The RNA levels in the ovary increased gradually in the larvae, pupae and the adults of Corcyra cephalonica, whereas in the Coriandrum oil treated resultant larvae there was a prominent decrease in the protein content when compared with the controls.

KEY WORDS: Coriandrum oil, Corcyra cephalonica, RNA, ovary, larvae, pupae and adult.**INTRODUCTION**

Proteins are the first biological factors making their manifestation during development. During metamorphosis of an insect, process like destruction of certain larval tissue and rejuvenation and remolding of various tissues into adult. One is bound to take place involving synthesis and consumption of the macro molecules as well^[1]. The Fat body tissue plays a key role in storage proteins. Storage proteins increased during successive stages of development^[2-3].

RNA synthesis is a key biosynthetic pathway which operates actively during early larval development in holometabolous insects and is thought to be an important preparatory mechanism for active metabolic functions to be carried out later by different organs during late larval development^[4]. The resultant increase in nuclear volume and RNA are proportional to the increase in cell size. The studies on various tissues / organs show that there is a relationship between cellular RNA synthesis and capacity of the cells for differentiation^[5]. Mitosis is necessary for a cell to change from one developmental stage to another, presumably to a more mature stage, Coccinelled beetles, Mosquito^[6] show that during each larval moult cycle there is a definite temporal pattern of RNA synthesis in various tissues^[7-9].

Coriandrum oil an essential oil used as a tonic and vermifuge and also in smoking for relief from catarrh and headaches. They are also used as insect repellents^[10-11] Coriandrum oil induces morphological changes and biochemical changes^[12]. The RNA levels in the Ovary of Corcyra cephalonica, were studied in the Coriandrum oil treated instars.

MATERIALS AND METHODS

A rich standard culture of this insect was maintained in the laboratory on normal dietary medium composed of coarsely ground jowar (Sorghum vulgar) inside a glass container at 26±1°C temperature and 65±5% Relative humidity.

Coriander Essential Oil Extraction

Essential oil was extracted from the seed of coriander, coriander sativum L. of the family Apiaceae. The essential oil was extracted by steam distillation for 4-6 hours using a Clevenger type apparatus where 250g of seeds in 250 ml of water is subjected to hydrodistillation. The oil was separated dried over anhydrous sodium sulfate and stored in dark glass bottles at 4 °C in the refrigerator until used. The isolated oil is a colorless or pale yellow liquid with a characteristic odor and taste of coriander. The larvae and adults were treated with the coriander essential oils prepared using ethyl alcohol (95%) as solvent. After suitable gap of time the insects were transformed into the diet. The treated larvae were observed daily to note the changes and the resulting abnormal intermediates were collected from the diet media. Fat body is dissected and rinsed free of haemolymph with Ringers solution. 10% homogenate was prepared for the estimation of proteins and the protein was estimated by the method of Lowry et al 1951.^[13]

RESULTS

Statistical Analysis of the Data: The experimental data was analyzed statistically, mean and standard Deviation was calculated. The RNA level in the Ovary was estimated in the control of larval stages, pupa and Adult.

Larval stages

The RNA content in the ovaries of the control larvae was much less than in the fat body. The first day of the V instar (18 day old) larvae recorded a value of 0.460 ± 0.003 mg/gm weight of the tissue. On the 2nd day it showed 0.490 ± 0.082 mg/gm weight of the tissue. On the 3rd day the RNA content further increased to 0.500 ± 0.033 mg/gm weight of the tissue (Figure 1).

Pupal stages

The RNA content in the ovaries of the freshly pupated pupa was 0.511 ± 0.034 mg/gm weight of the tissue. On the 2nd day the recorded value was 0.592 ± 0.039 mg/gm weight of the tissue of RNA was recorded. It further increased to 0.654 ± 0.043 mg/gm weight of the tissue on the 3rd day. The RNA content further increased from 0.692 ± 0.046 mg/g m weight of the tissue on the fourth day to 0.721 ± 0.048 mg/gm weight of the tissue on the 5th day (Figure 1).

Adult stage

The first day of the adult stage recorded a value of 0.850 ± 0.056 mg/gm weight of the tissue. The second day recorded a value of 1.420 ± 0.094 mg/gm weight of the tissue. The third day recorded a value of 0.982 ± 0.065 mg/gm weight of the tissue and 0.741 ± 0.049 mg/gm weight of the tissue on the fourth day (Figure 1).

Statistical Analysis of the Data: The experimental data was analyzed statistically, mean and standard Deviation

was calculated. The RNA levels in the Ovary was estimated in the treated of larval, pupa and Adult.

Larval stage

The RNA content in the ovaries of the treated resultant V instar larvae showed a marked decrease when compared to the controls. The first day of the V instar larvae noted a value of 0.162 ± 0.010 mg/gm weight of the tissue. It changed from 0.171 ± 0.0114 mg/gm weight of the tissue on the 2nd day to 0.184 ± 0.012 mg/gm weight of the tissue on the 3rd day of the V instar larvae (Figure 1).

Pupal period

The RNA content increased to 0.190 ± 0.012 mg/gm weight of the tissue on the 1st day of the pupal period. The RNA content increased further to 0.192 ± 0.012 mg/gm weight of the tissue on the 2nd day of the pupal period. On the 3rd, 4th, 5th days the RNA content recorded was 0.198 ± 0.012 mg/gm weight of the tissue, 0.203 ± 0.013 mg/gm weight of the tissue, 0.216 ± 0.014 mg/gm weight of the tissue respectively (Figure 1).

Adult:

The first day of the treated adult recorded a RNA level of 0.218 ± 0.014 mg/ gm weight of the tissue in the ovaries. The second and third day recorded values were 0.210 ± 0.014 mg/gm weight of the tissue and 0.028 ± 0.012 mg/gm weight of the tissue respectively (Figure 1).




Figure.1: Quantitative changes in the RNA content of the ovaries of the control and Coriandrum oil treated V instar resultant insects during the development of *Corcyra cephalonica*.

DISCUSSIONS

Corcyra cephalonica were treated with Coriandrum oil treated resultants showed a decline in the RNA content of the Ovary compared to the control larvae. The increase in the RNA content of the ovaries coincided with increase in protein content during larval, pupal and adult development in the control insects. The increased amounts of RNA in the ovaries of the controls are probably associated with mitosis of ovarian tissues during maturation of ovaries showed that RNA synthesis is correlated with increased ecdysteroid titres.^[14]

Coriandrum oil acts antagonistic to that of 20-hydroxyecdysone at the target site epidermis, inhibiting ecdysis.^[15-16] This may be due to the fact that Coriandrum oil inhibits mitosis thus inducing degeneration of cells, preventing growth, resulting in reduced levels of RNA in the tissues of the treated resultant *Corcyra cephalonica*.^[17-18]

CONCLUSION

The current study revealed that, in the Coriandrum oil treated insects, the cells of the ovary were smaller than the control larvae. The remnants of the ovary cells dispersed through the haemolymph. Hence building material for further tissue synthesis was unavailable and the ovary which depends on the fat body proteins for growth, degenerated, showing abnormally reduced amounts of RNA. The morphological observation and biochemical analysis of RNA, confirm the fact that Coriandrum oil deranges the development of *Corcyra cephalonica* by interfering with the hormonal milieu.

REFERENCES

- Venugopal, K.J and Dinesh Kumar, (1997), Electrophoretic studies on the development profiles of protein in Haemolymph, Fat body and ovary of red cotton bug, *Dysdercus Koenigii* Entomon, 22: 185-191.
- Kanost, M.R, Dawooga, J.K, Ryan, R.O., Husden, M.D, and Zeilger R., (1990), Insect haemolymph proteins in Adv. Insect Physiology, 22: 299-397.
- Rajathi A., Pandiaajan J., and Krishnan M., (2010) Effect of RH-2485 on the development, metamorphosis and synthesis of major proteins in female silkworm, *Bombyx mori* Biologia 65/5, 903-913.
- Dean, R.L, Bollenbacher, W.E, Locke, M, Gilgert L.I., and Smith S.L., (1985) Haemolymph ecdysteroid levels and cellular events in the intermoult/ moult sequence of *Calpodes ethilus* Journal of Insect physiology, 26: 267-280.
- Krishna Kumaran A., Berry S.J., Oberlander H, and Schnciderman H.A., (1967), Nucleic acid synthesis during insect development. II control of DNA synthesis in the *cecropia* silkworm and saturnid moths J. Insect. Physio. 13: 1-57.
- Dittman F., Koge P.H., and Hagedorn H.H., (1989) Ploidy levels and DNA synthesis in fat body cells of the adult mosquito, *Aedes aegypti*; the role of juvenile hormone, Arch.Insect, Biochem. Physiol 12: 133-143.
- Anitha, H.R., Sabita Raja S., Manjula C., and Raman C.V., (1999) Effect of Precocene –II on the nucleic acid content in the ovaries of *Chilo partellus*, Swinehoe, Entomon, 24: 307-313.
- Anuradha P., and Amarjith Kaur., (2010) Influence of Solasodine on the protein content of *Bombyx mori* L, J. Insect Physiol, 23: 15-18.
- Manjula C., and Sabita Raja S., (2001), Effect of precocene-II on the protein changes in the Haemolymph, Fatbody and ovaries of *Chilo partellus* during ontogenesis, Convergence, 2: 18-.
- Dharmasri M.G., Jayakody J.R.A.C., and Galhena G., (2003) Anti-inflammatory and analgesic activities of mature fresh leaves of *Vitex negundo*, J. Ethnopharmacol. 87: 199-202.
- Umamaheshwari M., Asok Kumar K., and Somasundaram A., (2007) Xanthine oxidase inhibitory activity of some Indian medical plants J. Ethnopharmacol, 109: 547-551.
- Ignacimuthu S., (1998) Nature's ecofriendly arsenal of pesticides, Curr. Sci, 74: 1037.
- Lowry O.H., Rosebrough, J.J., Farr A.L., and Randall R.J., (1951) Protein measurement with the folin phenol reagent, Journal Biology of Chemistry, 193: 263-275.
- Vanderberg J. P., (1963) Synthesis and transfer of DNA, RNA and protein during Vitellogenesis in *Rhodinus Prolixus* (Hemiptera), Biol. Bull, 125: 556-575.
- Tefler W. H., (1965), The mechanism and control of yolk formation, Ann-Rev. Ent, 10: 161-184.
- Schluter U., (1987) Effects of Azadirachtin on developing tissues of various insect larvae. Proc. 3, Int. Neem Conf, Eschborn, Germany, 331-348.
- Lafont R., Manchamps, B, Blais C., and Penetier J.L., (1977), Ecdysones and imaginal disc development during the last larval instar larvae of *Pieris brassicae*, J. Insect Physiol, 23: 277-283.
- Lobbecke E.A., (1969) Autoradiographische Bestimmung der DNA-Synthese-Daner Von Zellen der Flugelmainalanlage von *Ephestia kulneilla* Z. Wilhelm Roxus Arch. Entw. Mech. Org, 162: 1-18.