

HERBAL & DIETARY SUPPLEMENT CAUSING HEPATOTOXICITYJyoti Sharma*¹, Aakarshan Kumar², Pooja Verma², Damit Kumar² and Shubham Bhardwaj*³¹Student, Chandigarh University, Chandigarh.²Assistant Professor, IEC Univeristy, Baddi, H.P.³IEC University, Baddi, H.P.

*Corresponding Author: Jyoti Sharma

Student, Chandigarh University, Chandigarh.

Article Received on 19/01/2021

Article Revised on 09/02/2021

Article Accepted on 02/03/2021

ABSTRACT

The liver which is also known as the hepatic is the biggest organ and centre for the drug and foreign particle metabolism. Metabolism is mostly influenced by the medication or any xenobiotic activity whether it is harmful or not. Herbal drug induced hepatotoxicity is common and its frequency is difficult to note because of not reporting or underreporting, difficulties in the detection of amount or identification of the materials and incomplete observation of drug exposure. The death rate is high regarding drug induced in liver cause harm. Intense liver failure is due to supported medications is drawn more than 50% found in exceedingly studies. Although absolutely one-of-a-kind studies location unit available uncovering the mechanistic aspects of drugs induced hepatotoxicity, we will be inclined to vicinity unit inside the quandary concerning the virtual tale. The growing prevalence and effectiveness of writing and natural product inside the remedy of assorted issues crystal rectifier the investigators to seem into their capacity in countering drug-triggered liver toxicity. Much herbal merchandise is mentioned thus far to mitigate drug-prompted toxicity. The nutritional nature and less negative reactions of the natural product offer them an additional aspect over one-of-a-kind candidates of supplementary remedy.

KEYWORDS: xenobiotic, hepatotoxicity, quandary, prevalence, mitigate.**INTRODUCTION**

The liver plays a great function in the metabolism and elimination of drugs. Detoxification of medication and xenobiotics within the liver by using drug-metabolizing enzymes (DMEs) is a crucial development inside the acquisition of physiological conditions.^[1] Alteration in equilibrium status outcomes in a shift in the dynamic equilibrium of metabolism toward the (reactive oxygen species) ROS era thereby aerobic strain resulting in organ malfunction. Liver insufficiency and harm resulted from exposure to environmental toxicants, unique combinations, or dosages of prescribed drugs and microbial metabolites are major reasons of sickness and loss of life worldwide. Intake of drugs, their metabolism, and elimination build a circumstance of dynamic equilibrium to attend to the physiological circumstance. Phase I clinical trial and clinical check enzymes play a important role within the metabolism and cleansing of varied medication and xenobiotics. For the purchase of dynamic physiological conditions, extremely tuned metabolic control of medication or xenobiotics through xenobiotic-metabolizing enzymes is required. Any imbalance within the hobby of those enzymes ultimately outcomes within the shifting of equilibrium towards an intensive era that could eventually bind to macromolecules like deoxyribonucleic acid to purpose mutation, macromolecule to motive membrane harm, or

proteins to exchange their activities.^[2] The generation of reactive intermediates could be a common event in liver damage resulting from a number toxic medicinal drug and solvents. The toxic reactions due to remedy are also summarized as acute reactions (which include hepatocellular necrosis), disillusioned (with or at the same time as not infection), and miscellaneous reactions; however, a few medication will cause persistent harm and should even motive tumor growth. In the course of this article, we have stated drug-precipitated hepatotoxicity, factors a good way to augment its toxicity, mechanism, and bar. The listing of the 10 most in most cases used flavoring arrangements for various situations, enclosed Echinacea, garlic, Ginkgo Biloba, ginseng, grape seed extract, green tea, St.John's wort, bilberry, and succulent.^[3] especially, the usage of HDS is in particular not unusual in humans with persistent disease, as turned into stated via 30–62% of sufferers World Health Organization ate silymarin (milk thistle).^[4] This populace is in principle plenty vulnerable for hepatotoxicity similarly to lots of likely to develop severe internal organ destructive results.

Several Factors That Affect Hepatotoxicity

There are several factors that affect a man or woman's susceptibility to a doubtless toxic drug. Such as Advancing age of the person, gender like man or women,

manner factors, obesity due to any issues like hormonal changes or eating habits, nutritional standing in the body, genetic background, and dose of the drug required, and regime of medication may have an effect on the body to react the danger of herbs that consume and show poisonous reactions. Persons struck by exceptional sicknesses, along with, human immunological disease virus, hepatitis C, rheumatism, cirrhosis and autoimmune disease, are quite a few at risk of adverse drug reactions. Drug composition and interaction between drug-drug molecules can be a reason for the augmented risk of drug-caused hepatotoxicity. As an example, bound medication containing a nitro-aromatic moiety or medication interacting with nuclear receptors like phenobarbitone may motive organ-selective toxicity or may raise the toxicity of various medicine.^[5]

Mechanism of Hepatotoxicity

The liver synthesizes, concentrates, and secretes bile acids and excretes out the toxicants, such as bilirubin. Herbs induced injury to hepatocytes can lead to cholestasis. In turn, cholestasis causes intrahepatic accumulation of harmful toxin and excreted products which also lead for hepatic injury or hepatotoxicity. The liver has regenerative power, but the regeneration of liver cells lost by necrosis or apoptosis of cell death may mask the hepatotoxicity induced by herbs. The mechanism of herb-caused hepatotoxicity is variable and sophisticated in so many manners. Some herbs directly deadly and start to showcase poisonous reactions, that unit dose-related, within hours, while others could in all likelihood turn out to be liver harm only in inclined individuals and symptoms appear as soon as few days or perhaps weeks later. These reactions not often allergic or further as it should be harmful for any persons. Herb-drug interaction, though the herb shouldn't work as medicine does no longer seem toxic themselves, and also play an essential role in the propagation of toxicity.^[6] The biological technique of herb-induced toxicity mediate through reactive metabolites has been a centre of examining enthusiasm for the reason those developing to attack the uncovered the association among these metabolites an extensive reason for poisonous reactions may also be a drug-brought on intrahepatic sickness that frequently takes place all through the drug discovery and development method. The necessary roles of ROS among the cellular harm investigated and has been recommended that the valency binding of ROS, as well as reactive intermediates to macromolecules, could probably in all probability make a contribution to the severe dangerous drug reactions. Several studies recommend that the technology of unfastened radicals from poisonous medicinal drugs.^[7] Membrane lipid peroxidation is without delay associated with the depletion of tissue GSH (and bodily item antioxidant) main to the altered practical integrity of those structures and if the deterioration is intense; it must rather be lethal.^[8] Membrane lipid peroxidation would in all likely to result in alteration in membrane liquidness and porosity, augmented prices of natural compound

degradation, and in the end death. The concept is upheld with the aid of the style that aerobic harm to RBCs causes lack of membrane capability through enhancing lipid peroxidation (LPO) and enhancing the red blood cell rely on The attention of bodily item GSH, consequently, is that the important thing determinant of membrane integrity and hence the quantity of toxicant-brought about organ cellular harm.^[9]

Herb-Drug Interactions

In addition to the capability for direct effectiveness of the herbs, they also cause hepatotoxicity; some of these herbs ought to have interactions with certain prescription medications with the aid of various mechanisms resulting in damaging, collectively with the synergy of threat for hepatotoxicity, excretory organ toxicity, abnormal trauma, graft rejection, and vas crumble.^[10] Numerous herbs are known as substrates, inhibitors, and/or inducers of assorted hemoprotein P450 enzymes, such as john's wort, garlic, pepper, liquorice, flavonoids, triterpenoids, and anthraquinones.^[11] For instance, St.John's wort may be a potent inducer of CYP3A4, mediated by activation of the pregnane X receptor, which may then raise the intrinsic hepatotoxicity of various substances, like subshrub and other which are not subshrubs but likely to subshrub, through the way of partner degree enlarged conversion to toxic metabolites.^[12] It even increases the plasma clearance of a range of medication, like cyclosporine and enzyme inhibitors, which may additionally complicate the management of post transplant immunological sickness, further as HIV and viral hepatitis clinical resource.^[13] Additionally, coadministration of St.John's wort appreciably enlarged the overall exposure and toxicity of the immunosuppressive drugs in an incredibly rat model. Many herbs, together with garlic, papaya, tamarind, feverfew, and ginkgo Biloba, are related to partner degree enlarged danger of trauma in patients UN company surrounding area unit on decoagulant and/or pain pill. Distinctive herb-drug interactions so that you can end in hepatotoxicity or appreciably have an effect on the study.

Herbalife Products

Herbalife merchandise unit distributed as seasoner and dietary supplements at intervals the diversity of drinks, tablets, capsules, powders and energy bars for weight management, fitness, cosmetics, method to support and improvement in well-being. It's one altogether the foremost vital weight management and method supplement corporations at intervals the earth, with activity in nearly sixty countries, and sales of over three billion U.S. greenbacks.^[14,15] Pattern of liver injury was hepatocellular at intervals the majority of cases, however mixed and cholestatic patterns. Severity ranged from mild-to-severe liver hurt and fenced in cases that developed disease of the liver and ALF requiring LT.^[16] Relation was assessed by varied wide used scores, and it absolutely was thought of 'probable' in most cases, though 'certain' (with positive recalling) cases were

precise mechanism of liver injury has not been established, however Elinav *et al.* hypothesised that immune-mediated injury are often a gettable explanation, supported their observation of lymphocyte infiltrates and occasional transient auto antibodies presence in some cases.^[17] Recently, Stickel *et al.* according on a mix of cases of severe hepatotoxicity related to consumption of Herbalife merchandise contaminated with Bacilli which is obtained from the animal products. Relation in line with council for international organizations of medical sciences (CIOMS) was scored as 'probable' in each cases, and anatomy showed cholestatic and lobular/portal and biliary problem with disease of the liver in one patient. Further unassailable that culture supernatants of the Bacilli isolated from the merchandise iatrogenic escape of lactate dehydrogenase (LDH) from HepaG2 cells, that they taken as a results of the thought for the liver injury, thus raising concern of adulteration of Herbalife merchandise with hepatotoxin-producing organism strains. Therefore, it's quite apparently that hepatotoxicity will occur among some those that receive these merchandise, however the precise mechanism or the accountable agent at intervals the merchandise is unsure, half as a results of, to date, the entire listing of the ingredients of those merchandise isn't illustrative and collectively the manufacturer apparently is unwilling to provide the specified information about the heabal products.^[18] Besides, Herbalife representatives got to be compelled to the present time challenged the tributary relationship between consumption of their merchandise and drug-induced liver injury (DILI), still as confirming smart management for ingredients in the products and contaminants in their production lines.

KAVA

Kava (*Piper methysticum*) could be a plant native to the Pacific Islands, together with Hawaii, Vanuatu, Melanesia and a few elements of Micronesia, wherever Associate in nursing of rootstalk extract from kava to prepare a social and recreational functions with standard foods. In Western countries, alcoholic beverages found in herbal supplements are promoted as Associate in nursing agent to alleviate stress, anxiety, and tension, also as for passing state and biological time symptoms.^[19] Its effectiveness for the hysteria treatment is supported by a meta-analysis cochrane. Varied severe hepatotoxicity reports are delineate within the USA and Europe, and a few of that are established by structured, quantitative and liver-specific relation measurement ways.^[20] In Associate in Nursing analysis of thirty six cases of alcoholic beverage hepato-toxicity, the injury pattern was each hepatocellular and cholestasis; girls are than boys in the majority of the patients, the additive dose and latency were extremely variable, and acute liver failure (ALF) developed in nine patients and eight of them undergo Liver Treatment.^[21] The US-FDA began to advise of herbal merchandise on the sevre liver injury risk related to the employment dietary supplements within the year 2002 and also the merchandise are prohibited from some of the countries in Europe, though

they're still on the market within the USA, Canada, Australia, New Sjaelland and Pacific Islands, also as from the web.^[22]

Pennyroyal

Pennyroyal (squawmint oil) is associate herb containing pulegone and monoterpenes in a smaller quantity, which is in a very sort of oil. It has been used as associated with the abortifacient, in spite of this kava is consuming, without a doubt which is deadly toxic and toxin effects are showing.^[23] Pennyroyal's toxicity is supposed to be mainly from menthofuran, a substance of pulegone, which is oxidized by hemoprotein P450. Additionally, pulegone shows the deterioration of glutathione in liver tissue and plasma is measured.^[24] So, drug inhibiting hemoprotein P450 (i.e.antiacid and alcohol inhibiting drug) and/or glutathione replacement with N-acetylcysteine could be in theory to improve or forestall pennyroyal hepatotoxicity.^[25] Anderson *et al.* revealed in the literature review of hepatotoxicity twenty-two cases related to pennyroyal oil. Patients usually have severely developed epithelial duct upset and central systema nervosum effects at Severe/fatal internal organ death and multiorgan failure looked as if it would occur once quite fifteen millilitres was eaten.^[26]

Pyrrrolidine Alkaloids

Pyrrrolidine alkaloids (PA) found in additional more than 350 plant species worldwide and measure through the square method. These alkaloids are imposing the liver toxicity for over seventy years, significantly with Senecio, Heliotropium, Crotalaria, and with other similar species. The Greek fret of liver injury is VOD (veno-occlusive disease), recently termed curved obstruction syndrome (SOS), that was 1st reportable in Jamaican kids World Health Organization drank bush tea for his or her malady.^[27] after, many cases of VOD related to associate activity of PA-containing plants (most usually as flavourer tea) are reportable chiefly from the Southern US, Africa, and Asia, yet a periodically from Western Countries.^[28] VOD in pyrrrolidine alkaloids associated usually presents through pathology, dropsy and abnormal condition. The clinical onset of action of herbs and severity square measure variable helps in the diagnosis hepatotoxicity. Within the acute kind liver injury, which shows the symptoms of developing the pain in abdominal, usually with jaundice and clearly elevated altitude levels and with rapidly damaging in the tissues level and will eventually ends in death?^[29] Morbific method inflicting obstruction to the curved blood flow starts with harm to sinusoid epithelial tissue that ends up in partial curved lumens obstruction.^[30] Liver microscopic anatomy is study to consider the eventually, haemorrhagical centrilobular death by non thrombotic obstacle of tiny ending of internal organ venules, resulting in curved dilatation.^[31] Acute VOD is fatal in adults as compare to the kids who are not consume till date, almost 15–20% of patients of it in adults gets worsen cases. Complete recovery has been ascertained in regarding half of the

patients, characterized by perivenular and bridging pathology, and a few could die as they withdrawn the course due to not having the compliance to the patients disease many years later. A smaller proportion of patients could have acute or chronic diseases problems, which can perniciously reach liver disease and malignant hypertension. In animal models, biotransformation by hemoprotein P450 3A4 causes the hepatotoxicity of PA appears to be unstable toxicant metabolites (pyrrole derivatives) that will act as alkylating agents in further process and this relies on the sort of PA total dose eaten, together with status to the alkaloids.^[32] Especially, herb–drug or herb–herb interactions involving hemoprotein P450 square measure doubtless to have an effect on VOD status and severity of the cases have toxicity may be amplified by concomitant use of Purple Heart via the hemoprotein P450 induction. Acute VOD could is caused by a brief time exposure to high doses, whereas chronic liver injury is caused by prolonged exposure to even tiny doses of PA.^[33] Defibrotide, a polydisperse oligonucleotide with native antithrombotic, anti-ischaemic and anti-inflammatory activity, seems to be effective for severe VOD following biological process vegetative cell transplant;^[34] but, As before long as acute liver failure (ALF) or chronic liver failure seems at hand, LT is also the sole effective medical aid.^[35]

Other Herbal and Dietary Supplement

Numerous alternative flavourer product, together with oil (Cinnamomum camphora, Vicks VapoRub),^[36] rattle-top, Serenoa repens, Noni juice (Morinda citrifolia), Green tea (*Camellia sinensis*), cascara sagrada, mistletoe (*Viscus album*),^[37] skullcap (*Scutellaria*), flower (*Valeriana officinalis*), shrub (*Cassia angustifolia* and *C.acutifolia*), Germander (*Teucrium chamaedrys*), usnic acid,^[38] tree oil of variety of neem (*Antelaea Azadirachta*, *Azadirachta indica*), thirty eight and burn plant,^[39] yet as dietary supplements, together with vitamin A are delineated to cause hepatotoxicity.^[40]

CONCLUSION

As flavouring merchandise of herbs still be used wide round the World, flavouring hepatotoxicity can still be determined. Such events don't seem to be essentially distinctive to flavouring medications as they'll be seen with prescription like antibiotics, anticonvulsants, etc. it's so essential that the popularity and reportage of flavouring hepatotoxicity be command to an equivalent standards as prescription medications. Liver injury is generally hepatocellular, however mixed and cholestatic patterns may also occur, and therefore the severity ranges from gentle injury to Acute Liver Failure, still likely to evolve the liver chronicity. The identification of herbs which causes the vital organ failure or damage due to toxicity depends on a correct data of the offered literature on hepatotoxicity with the spectrum of herb dosages in today lifestyle also matter as how much required to intake for effectiveness and not for the toxicity through checking the metabolism of different age and additionally on a heightened awareness. As anything

taken in high dose or on regular basis whether it's natural can also cause harm to the body by accumulating at a specific site, if not metabolised. Advances within the understanding of the frequency, the pathological process, the clinical manifestations and outcomes square measure required to be able to improve herb as a medication safety.

REFERENCES

- Ostapowicz, G., Fontana, R. J., Schiodt, F. V., Larson, A., Davern, T. J., Han, S. H., et al. Results of a prospective study of acute liver failure at 17 tertiary care centers in the United States. *Ann. Intern. Med.*, 2002; 137: 947–954.
- Upadhyay, G., Gupta, S. P., Prakash, O., and Singh, M. P. Pyrogallol-mediated toxicity and natural antioxidants: triumphs and pitfalls of preclinical findings and their translational limitations. *Chem. Biol. Interact.*, 2010a; 183: 333–340.
- Bent S, Ko R. Commonly used herbal medicines in the United States: a review. *Am J Med.*, 2004; 116: 478–85.
- Verma S, Thuluvath PJ. Complementary and alternative medicine in hepatology: review of the evidence of efficacy. *Clin Gastroenterol Hepatol*, 2007; 5: 408–16.
- Boelsterli, U. A., Ho, H. K., Zhou, S., and Leow, K. Y. Bioactivation and hepatotoxicity of nitroaromatic drugs. *Curr. Drug Metab*, 2006; 7: 715–727.
- Yamazaki, Y., Kakizaki, S., Horiguchi, N., Takagi, H., Mori, M., and Negishi, M. Role of nuclear receptor CAR in carbon tetrachloride-induced hepatotoxicity. *World J. Gastroenterol*, 2005; 11: 5966–5972.
- Racknagel, R. O., Glende, E. J. R., Dolak, J. A., and Waller, R. I. Mechanisms of carbon tetrachloride toxicity. *Pharmacol. Ther.*, 1989; 43: 130–154.
- Park, B. K., Kitteringham, N. R., Maggs, J. L., Pirmohamed, M., and Williams, D. P. The role of metabolic activation in drug-induced hepatotoxicity. *Ann. Rev. Pharmacol. Toxicol*, 2005; 45: 177–202.
- Ross, D. Glutathione, free radicals and chemotherapeutic agents. *Pharmacol. Ther.*, 1988; 37: 231–249.
- Fasinu PS, Bouic PJ, Rosenkranz B. An overview of the evidence and mechanisms of herb-drug interactions. *Front Pharmacol*, 2012; 3: 69.
- Zhou S, Gao Y, Jiang W, Huang M, Xu A, Paxton JW. Interactions of herbs with cytochrome P450. *Drug Metab Rev.*, 2003; 35: 35–98.
- Schiano TD. Hepatotoxicity and complementary and alternative medicines. *Clin Liver Dis.*, 2003; 7: 453–73.
- Posadzki P, Watson L, Ernst E. Herb-drug interactions: an overview of systematic reviews. *Br J Clin Pharmacol*, 2012.
- Stickel F, Kessebohm K, Weimann R, Seitz HK. Review of liver injury associated with dietary supplements. *Liver Int*, 2011; 31: 595–605.

15. Chao S, Anders M, Turbay M, Olaiz E, McCormack L, Mastai R. Toxic hepatitis by consumption Herbalife products a case report. *Acta Gastroenterol Latinoam*, 2008; 38: 274–7.
16. Stickel F, Droz S, Patsenker E, Bogli-Stubler K, Aebi B, Leib SL. Severe hepatotoxicity following ingestion of Herbalife nutritional supplements contaminated with *Bacillus subtilis*. *J Hepatol*, 2009; 50: 111–7.
17. Elinav E, Pinsker G, Safadi R, et al. Association between consumption of Herbalife nutritional supplements and acute hepatotoxicity. *J Hepatol*, 2007; 47: 514–20.
18. Seeff LB. Are herbals as safe as their advocates believe? *J Hepatol*, 2009; 50: 13–6.
19. Teschke R. Kava hepatotoxicity: pathogenetic aspects and prospective considerations. *Liver Int*, 2010; 30: 1270–9.
20. Seeff LB. Herbal hepatotoxicity. *Clin Liver Dis.*, 2007; 11: 577–96.
21. Teschke R. Hepatotoxicity by drugs and dietary supplements: safety perspectives on clinical and regulatory issues. *Ann Hepatol*, 2009; 8: 184–95.
22. Stickel F, Baumuller HM, Seitz K, et al. Hepatitis induced by Kava (*Piper methysticum rhizoma*). *J Hepatol*, 2003; 39: 62–7.
23. Teschke R, Schulze J. Risk of kava hepatotoxicity and the FDA consumer advisory. *JAMA*, 2010; 304: 2174–5.
24. Schiano TD. Hepatotoxicity and complementary and alternative medicines. *Clin Liver Dis.*, 2003; 7: 453–73.
25. Sztajnkrzyer MD, Otten EJ, Bond GR, Lindsell CJ, Goetz RJ. Mitigation of pennyroyal oil hepatotoxicity in the mouse. *Acad Emerg Med.*, 2003; 10: 1024–8.
26. Thomassen D, Slattery JT, Nelson SD. Menthofuran-dependent and independent aspects of pulegone hepatotoxicity: roles of glutathione. *J Pharmacol Exp Ther.*, 1990; 253: 567–72.
27. Bras G, Jelliffe DB, Stuart KL. Veno-occlusive disease of liver with nonportal type of cirrhosis, occurring in Jamaica. *AMA Arch Pathol*, 1954; 57: 285–300.
28. Tandon RK, Tandon BN, Tandon HD, et al. Study of an epidemic of venoocclusive disease in India. *Gut*, 1976; 17: 849–55.
29. Mohabbat O, Younos MS, Merzad AA, Srivastava RN, Sediq GG, Aram GN. An outbreak of hepatic veno-occlusive disease in north-western Afghanistan. *Lancet*, 1976; 2: 269–71.
30. Stillman AS, Huxtable R, Consroe P, Kohnen P, Smith S. Hepatic veno-occlusive disease due to pyrrolizidine (Senecio) poisoning in Arizona. *Gastroenterology*, 1977; 73: 349–52.
31. Bach N, Thung SN, Schaffner F. Comfrey herb tea-induced hepatic veno-occlusive disease. *Am J Med.*, 1989; 87: 97–9.
32. Ridker PM, Ohkuma S, McDermott WV, Trey C, Huxtable RJ. Hepatic venoocclusive disease associated with the consumption of pyrrolizidine-containing dietary supplements. *Gastroenterology*, 1985; 88: 1050–4.
33. DeLeve LD, Wang X, Kuhlenkamp JF, Kaplowitz N. Toxicity of azathioprine and monocrotaline in murine sinusoidal endothelial cells and hepatocytes: the role of glutathione and relevance to hepatic venoocclusive disease. *Hepatology*, 1996; 23: 589–99.
34. Huxtable RJ. New aspects of the toxicology and pharmacology of pyrrolizidine alkaloids. *Gen Pharmacol*, 1979; 10: 159–67.
35. Mattocks AR. Toxicity of pyrrolizidine alkaloids. *Nature*, 1968; 217: 723–8.
36. Jimenez JF, Brown AL, Arnold WC, Byrne WJ. Chronic camphor ingestion mimicking Reye's syndrome. *Gastroenterology* 1983; 84: 394–8.
37. Singh YN, Devkota AK, Sneed DC, Singh KK, Halaweish F. Hepatotoxicity potential of saw palmetto (*Serenoa repens*) in rats. *Phytomedicine* 2007; 14: 204–8.
38. Stadlbauer V, Fickert P, Lackner C, et al. Hepatotoxicity of NONI juice: report of two cases. *World J Gastroenterol* 2005; 11: 4758–60.
39. Rabe C, Musch A, Schirmacher P, Kruis W, Hoffmann R. Acute hepatitis induced by an *Aloe vera* preparation: a case report. *World J Gastroenterol*, 2005; 11: 303–4.
40. Geubel AP, De Galocsy C, Alves N, Rahier J, Dive C. Liver damage caused by therapeutic vitamin A administration: estimate of dose-related toxicity in 41 cases. *Gastroenterology*, 1991; 100: 1701–9.