



BACTERIOLOGICAL PROFILE IN COMMUNITY ACQUIRED PNEUMONIA IN A TERTIARY CARE HOSPITAL

Vinut S. Math*¹, Sanjeev Chetty² and A. M. Koppad³

Department of Paediatrics, Navodaya Medical College Hospital and Research Centre, Raichur.

***Corresponding Author: Vinut S. Math**

Department of Paediatrics, Navodaya Medical College Hospital and Research Centre, Raichur.

Article Received on 21/01/2023

Article Revised on 10/02/2023

Article Accepted on 02/03/2023

ABSTRACT

Introduction: Community acquired pneumonia (CAP) refers to clinical signs and symptoms of pneumonia acquired outside a hospital setting.^[1] **Methods and Material:** It was a prospective study conducted at department of Paediatrics, Navodaya Medical College Hospital and Research Centre, Raichur for a period of 1 year on children with community acquired pneumonia admitted in ward and PICU. Total 60 cases were enrolled in the study according to recent WHO guidelines of community acquired pneumonia. Oropharyngeal swab and blood culture samples were collected from every child and sent for culture and sensitivity. **Results:** Out of 60 cases, 43 (71.6%) were diagnosed as pneumonia and 17 (28.4%) were as severe pneumonia. On blood culture only 11 (18.3%) had positive growth and on oropharyngeal swab culture 14 (23.33%) had growth. Microorganisms isolated in blood culture were predominantly *Staphylococcus aureus*, while in throat swab culture *Streptococcus* predominated. **Conclusion:** The overall rate of isolation of organism in CAP is low. The yield of culture positivity was 23% in throat swabs, with *Streptococcus pneumoniae* being most common isolate, whereas 18% in blood culture with *Staphylococcus aureus* being predominantly isolated organism.

KEYWORDS: CAP, Blood culture, Oropharyngeal swab culture.

INTRODUCTION

Acute respiratory tract infections are the largest cause of morbidity and mortality among under-five children worldwide.^[2-6] Pneumonia are the number one cause of under-5 child mortality, responsible for nearly 400,000 deaths in India annually.^[7] In paediatric age group, the developing immune system and age related exposures result in infection caused by varied yet consistent set of bacterial and viral pathogens, which are different in different countries and changing with season within the same country. Recent systematic reviews of childhood pneumonia etiology suggest that in developing countries, a few bacteria (*S. pneumoniae* and *H. influenzae*) and viruses (respiratory syncytial virus, influenza virus) are associated with majority of childhood CAP.^[7]

In India, there are only a few systematic review or studies regarding the etiology of pneumonia in children.^[2,5] In general, the etiology can be identified in only 30%–50% of cases using conventional methods.^[6] This lack of information leads to inappropriate use of antibiotics with consequent antibiotic resistance and increased cost.^[6] Ascertaining the etiology in childhood CAP is important because it plays an important role in shaping the individual treatment decisions, antibiotic policy in the community, and also rationalize immunization policy at a national level. Our study is

aimed at recognising the common etiological agents and sensitivity pattern of organisms in children suffering from CAP and admitted in our centre.

METHODS

Study setting- This study was a hospital based prospective study and was conducted in the department of Paediatrics, Navodaya Medical College Hospital and Research Centre Raichur, after the ethical committee clearance.

Study Design- The study design was prospective and observational study.

Study Period- This study was conducted from August 2019 to September 2020.

Sample Size- The sample size of study was 60.

Inclusion Criteria- Children aged 2 months to 12 years who presented with history of fever, cough and difficulty in breathing fulfilling WHO case definition of CAP were enrolled. According to WHO^[8] pneumonia and severe pneumonia are defined as follows.

Pneumonia: Patient present with cough, cold, fever, rapid breathing and chest indrawing and tachypnoea is defined as respiratory rate >60/minute for infant <2

months, >50/minute for infants 2-12 months, >40/minute for children >12-60 months, >30/minute for children >60-144 months.^[8]

Severe pneumonia: Patient present with sign and symptoms of pneumonia and associated with danger signs, like inability to drink, persistent vomiting, convulsion, lethargic or unconsciousness, stridor in calm child with severe malnutrition.

Exclusion Criteria- Children with duration of illness >7 days, those who had received antibiotics for >24 hours at presentation or those with previous hospitalization for >48 hours in last 2 weeks, children with wheeze who had received a single dose of bronchodilator (salbutamol @0.15mg/kg by nebulization) and symptoms disappeared were excluded.

Sampling Method- Detailed history and clinical examination done, sample for throat swab and blood for culture and sensitivity were collected with standard precautions from every child and sent for bacteriological lab within 30 minutes of collection.

Statistical Analysis- The data was collected and compiled systematically in a tabular form and then analysed statistically.

RESULTS

Total of 60 patients were enrolled in our study based on inclusion criteria. Most of the children i.e. 43 (71.6%) were diagnosed as pneumonia and rest 17 (28.4%) were classified as severe pneumonia. Based on the age group, out of 60, 38 (63.33%) children belonged to 2-12 months, 13 (21.6%) children belonged to age group of 13-24 months, and remaining 9 (15%) belonged to >2 years. Based on gender, 38 (63.33%) were male and 22 (33.66%) were female, ratio being 1.7:1. Most of them belonged to low SES 32 (53.33%), followed by lower middle class 14 (23.33%). It was observed that majority 33 (55%) were having no infiltrates on chest xray. Current study revealed positive blood culture was seen only in 11 (18.3%), while rest were negative. Children with severe pneumonia had 8 (47.05%) culture positives when compared to children with pneumonia who had only 3 (6.9%) positive culture growth. On applying chi square test we found significant result (P value = 0.0003) which revealed growth on blood culture depends on severity of illness. Bacteriologically, blood culture showed predominant growth of *Staphylococcus aureus* 6 (10%), followed by *Streptococcus pneumoniae* 2 (3.33%). On throat swab culture, 14 cases yielded growth, in which 5 had pneumonia and 9 had severe pneumonia, amongst which *Streptococcus pneumoniae* predominated 6 (42.8%) followed by *Staphylococcus aureus* 4 (28%), followed by *Klebsiella* 2 (14.2%) and *E. coli* 2 (14.2%) which were equal.

Table-1: Result of Blood Culture.

Blood C & S	Pneumonia	Severe pneumonia	Total	%
Growth positive	3	8	11	18.3%
Growth negative	40	9	49	81.7%
Total	43	17	60	-

Table-2: Result of Oropharyngeal Swab Culture.

Oropharyngeal swab culture	Pneumonia	Severe Pneumonia	Total	%
Growth Positive	5	9	14	23.33%
Growth negative	38	8	46	76.66%
Total	43	17	60	-

Table- 3: Bacterial Culture in Clinical Specimen.

Organisms	Blood culture {n=60}	Oropharyngeal swab culture {n=60}
<i>Staphylococcus aureus</i>	6(10%)	4(28%)
<i>Streptococcus pneumoniae</i>	2(3.33%)	6(42.8%)
<i>E. coli</i>	-	2(14.2%)
<i>Klebsiella species</i>	1(1.6%)	2(14.2%)
<i>Pseudomonas aeruginosa</i>	1(1.6%)	-
<i>Hemophilus influenzae</i>	1(1.6%)	-

DISCUSSION

There is a constant threat of emerging antibiotic resistance due to liberal travel across geographical borders, in-advert, irrational antibiotic use. Owing to limitation of data regarding antibiotic sensitivity and resistance pattern, this study was undertaken to create a

baseline antibiotic resistance database for formulating an effective infection control policy in our centre.

60 children who came with fever, cough, cold, rapid breathing and refusal to feed were included after excluding the exclusion criteria. In present study, 43 (71.6%) subjects were diagnosed as pneumonia and 17

(28.4%) diagnosed as severe pneumonia. Similar results were found in other studies done by different author.^[5,9,10,11,12]

Males were affected more commonly than females (63.33% Vs 36.66%) with ratio 1.7:1 in our study, which is similar to the study done by Aditi et al^[13] where Males predominated females i.e. 64.6% vs 35.4%, similar observations were made in other studies.^[10,11,12] Whereas a study from Bulgaria done by Guergana Petroval et al^[14] had almost equal male and female population enrolled in the study (48% vs 52%), which implies female neglect and gender biased attitude of our society.

In our study, majority of patients i.e. 76% were belonged to age group 2-12 months followed by 13-24 months (10%). In a study done in 2015^[13] majority of

patients 78.2% belonged to age group 2-12 months followed by 21.7% belonged to 12-59 months age group. Similar results were obtained in other studies.^[10,11,12,15]

This may be due to lack of exclusive breastfeeding for first 6 months, lack of immunization, vitamin A deficiency which aggravates the already deficit immune condition in infants.

Majority of subjects i.e. 28 (56%) belonged to lower socioeconomic status, which is again similar to others study.^[12,16,17,18] This could be attributed to overcrowding, poor parental education, indoor air pollution in deprived community.

The comparison of blood culture and oropharyngeal swab culture of various studies are given in table no. 4

Table-4: Comparison of organisms isolated in various study and their sensitivity pattern.

Sr. No.	Author	Sample/Test	Result
1	Patwari et al (1996) ^[19]	Throat swab, NPA, lung aspirate and blood	1. In infants <3 months - Escherichia coli [50%], Klebsiella [25%] and S. pneumoniae [18%]. 2. Between 7 and 24 months - H. Influenza [31%] 3. In all age groups - S. pneumoniae and S. aureus : common
2	Pandey et al (2000) ^[20]	Blood	1. S. pneumoniae isolated in 12.8% of all ALRTI 2. S. aureus isolated in 7.1% of all ALRTI
2	The SENTRY antimicrobial surveillance study (2000) ^[21]	Blood culture	Organism recovered were Staph. aureus (28%) Pseudomonas aeruginosa (10%) Strep. pneumoniae (9.1%) Klebsiella (7.5%) H. influenzae (7.3%)
4	Kanungo and rajalakshmi (2001) ^[22]	Blood	1. 7.3% of isolated S. pneumoniae were resistant to penicillin.
5	Awasthi et al. (2004; ISCAP) ^[23]	NPA and nasopharyngeal swab	1. 66.3% of isolated S. pneumoniae were resistant to cotrimoxazole, 4.1% to chloramphenicol, 15.9% to oxacillin and 2.9% to erythromycin
6	Capoor et al. (2006) ^[24]	Blood culture	1. blood culture positivity: 27.4% (by combine method of Bac T and conventional methods). 2. S. pneumoniae (35.3%), staphylococcus aureus (23.5%), klebsiella pneumoniae (20.5%) and haemophilus influenzae (8.8%)
7	Neuman MI et al 2007-11 ^[25]	Blood culture	Strep. pneumoniae accounted for 78% of all detected pathogen 82% susceptible for penicillin
8	Tiewsoh et al. (2009) ^[26]	Blood culture	1. blood culture positivity : 15% 2. the most common organism S. pneumoniae (40%)
9	Joseph et al 2011-13 ^[9]	Blood culture, PCR, BAL	NPA & blood culture Yielded bacteria in only 322 (13.7%) and 49 (2.1%) children respectively in NPA, Strep. pneumoniae predominated, followed by H. influenzae (9.6%) and Staph. aureus (6.8%). In blood S. aureus (30.6) dominated followed by S. pneumoniae (20.4%) and Klebsiella, pneumoniae (12.2%) M. pneumoniae and C. pneumoniae serology were positive in 4.3% and 1.1% respectively.
10	Anusmita das et al 2013-14 ^[10]	Blood culture	Blood culture was positive in 3% cases while oropharyngeal positive in 16.8% S. pneumoniae was major isolate
11	Yudhavir et al (2016) ^[11]	Blood culture	On blood culture S. aureus (10%) predominated followed by S. pneumoniae (3.1%) On NPA Strep. pneumoniae was predominated 18.5%

12	Chaudhary et al(2016-2017) ^[12]	Blood culture, oropharyngeal culture	On blood culture- staph. aureus was the most common organism isolated(8%) oropharyngeal culture- strep. pneumoniae was most common bacteria(10%)
----	--	--------------------------------------	---

In our study (2019-2020), oropharyngeal culture isolated predominately *Streptococcus pneumoniae*, whereas in blood culture the most common organism isolated was *Staphylococcus aureus* which is similar to the studies done above.

Out of 60 patients, 49 (81.67%) recovered and discharged uneventfully and 3(5%) children died, remaining 8(13.33%) patients in our study either absconded or were discharged against medical advice. Out of the 3 mortality, all 3 were from severe pneumonia group which can be concluded that mortality is more common in severe pneumonia age group.

Our finding is similar to the study conducted by Anusmita das et al (2016) who also reported similar mortality rate (i.e. 8.8%).^[10]

CONCLUSION

Thus following conclusions were made from our study- Incidence of CAP is highest during infancy and decreases with advancing age. Also the increased prevalence seen in Lower socioeconomic strata highlights the importance in addressing the preventable risk factors such as malnutrition, illiteracy, overcrowding, poor attitude towards vaccination, indoor air pollution. Vaccine coverage against *Streptococcal pneumoniae* and *H.influenza* needs to improve to reduce the mortality and morbidity. Identifying and changing barriers to accessing health care is important. The results achieved in our study may help in formulation of better antibiotic policy against community acquired pneumoniae in children less than 12 years in our region. There is a need of advanced diagnostic methods which improve the diagnostic yield by detection of fastidious and atypical organisms.

Limitation of the study-*Hemophilus Influenzae* could not be isolated in any of the cases as throat swabs were done by conventional method. Isolation of these organisms require PCR and DNA studies.

What this study adds to existing Knowledge: Blood culture and throat swab culture can help in diagnosis and management of resistant cases. But most of the results comes negative so clinical diagnosis and treatment is still useful in community acquired pneumonia.

REFERENCES

- Harris M, Clark J, Coote N, et al. British thoracic guidelines for the management of community acquired pneumonia in children: update, 2011; 66(2): 1-23. [PubMed:2190369].
- Mathew JL, Patwari AK, Gupta P, Shah D, Gera T, Gogia S, et al. Acute respiratory infection and pneumonia in India: A systematic review of literature for advocacy and action: UNICEF-PHFI series on newborn and child health, India. *Indian Pediatr*, 2011; 48: 191-218.
- Huong Ple T, Hien PT, Lan NT, Binh TQ, Tuan DM, Anh DD. First report on prevalence and risk factors of severe atypical pneumonia in Vietnamese children aged 1-15 years. *BMC Public Health*, 2014; 14: 1304.
- Wu Z, Li Y, Gu J, Zheng H, Tong Y, Wu Q. Detection of viruses and atypical bacteria associated with acute respiratory infection of children in Hubei, China. *Respirology*, 2014; 19: 218-24.
- Mathew JL, Singhi S, Ray P, Hagel E, Sagharian-Hedengren S, Bansal A, et al. Etiology of community acquired pneumonia among children in India: Prospective, cohort study. *J Glob Health*, 2015; 5: 050418.
- Liu J, Ai H, Xiong Y, Li F, Wen Z, Liu W, et al. Prevalence and correlation of infectious agents in hospitalized children with acute respiratory tract infections in Central China. *PLoS One*, 2015; 10: e0119170.
- Dubish S, Singh V. *Pneumonia*. In: Piyush G., PSN M, Siddhart R, Rakesh L. PG Text book of pediatrics. 2nd ed. New Delhi: Jaypee Brothers, 2018; 2: 2033-40.
- WHO. (2014) Revised WHO classification and treatment of childhood pneumonia at health facilities. Evidence Summaries, WHO 2014. Available at: http://apps.who.int/iris/bitstream/10665/137319/1/9789241507813_eng.pdf (accessed 18 August 2017).
- Agweyu A, Kibore M, Digolo L. Prevalence and correlates of treatment failure among Kenyan children hospitalized with severe community-acquired pneumonia: a prospective study of the clinical effectiveness of WHO pneumonia case management guidelines, 2014; 19(11): 1310-1320. doi:10.1111/tmi.12368.
- Das A, Patgiri SJ, Saikia L, Dowerah P, Nath R et al. Bacterial pathogens associated with community-acquired pneumonia in children aged below five years. *Indian pediatrics*, 53; 2016: 225-227. [PubMed].
- Shekhawat YS, Sharma P, Singh A, Payal V et al. Bacteriological and clinical profile of community acquired pneumonia in hospitalised children with associated co-morbidity in tertiary care centre of Western Rajasthan, India. *Int J Contempediatr*, 2016; 3(4):1380-1384. [ijcp20163682](http://dx.doi.org/10.18203/2349-3291). doi: <http://dx.doi.org/10.18203/2349-3291>.
- Chaudhary G.S, Kumar S, Kankane A, Gupta S. Microbiological profile in community acquired

- pneumonia in children. *Int J Pediatr Res.*, 2018; 5(5): 263-267. doi:10.17511/ijpr.2018.5.04.
13. Aditi Baruah, Arpita Gogoi, Helina Rahman.
 14. Microbiological Profile And Their Resistance Pattern Among Children with Community Acquired Pneumonia in A Tertiary Care Hospital in North-East India} *IOSR Journal of Dental and Medical Sciences (IOSR-JDMS)* e-ISSN: 2279-0853, p-ISSN: 2279-0861, October. 2016; 15(10): 01-05.
 15. Petrova, Guergana & Miteva, Dimitrinka & Parina, Snejana & Georgieva, Biliiana & Bijeva, Svetomira & Marteva-Proevska, Yulia & Lazova, Snezhina & Tzotcheva, Iren & Velinov, Tzvetan & Perenovska, Penka. Etiological profile of community acquired pneumonia in hospitalized children in Bulgaria in the era of the pneumococcal vaccine, 2019; 1-5. 10.15761/LBJ.1000138.
 16. Kumar K J, Ashok Chowdary K V, Usha H C, Kulkarni M, Manjunath V G. Etiology of community acquired pneumonia among children in India with special reference to atypical pathogens. *Lung India*, 2018; 35: 116-20.
 17. Nag VL, Ayyagari A, Venkatesh V, Ghar M, Yadav V, Prasad KN et al. Drug resistant Haemophilus influenza from respiratory tract infection in a tertiary care hospital in North India. *Indian J chest dis allied sci.*, 2001; 43: 13-17.
 18. Kilabuko JH, Nakai S. Effects of cooking fuels on acute respiratory infections in children in Tanzania. *Int J Environ Res Public Health*, 2007 Dec; 4(4): 283-8.
 19. Neuman MI, Hall M, Lipsett SC, Hersh AL, Williams DJ, Gerber JS, Brogan TV, Blaschke AJ, Grijalva CG, Parikh K, Ambroggio L, Shah SS; Pediatric Research in Inpatient Settings Network. Utility of Blood Culture Among Children Hospitalized with Community-Acquired Pneumonia. *Pediatrics*, 2017 Sep; 140(3): pii: e20171013. doi: 10.1542/peds.2017-1013. Epub 2017 Aug 23. [PubMed]
 20. Patwari, A., Bisht, S., Srinivasan, A. Aetiology of pneumonia in hospitalized children. *J Trop Pediatr*, 1996; 42: 15-20. Google Scholar | Crossref | Medline | ISI
 21. Pandey, A., Chaudhry, R., Nisar, N. Acute respiratory tract infections in Indian children with special reference to Mycoplasma pneumoniae. *J Trop Pediatr*, 2000; 46: 371-374. Google Scholar | Crossref | Medline | ISI
 22. Hoban DJ, Biedenbach DJ, Mutnick AH, Jones RN. Pathogen of occurrence and susceptibility patterns associated with pneumonia in hospitalized patients in North America: results of the SENTRY Antimicrobial Surveillance Study (2000). *Diagn Microbiol Infect Dis*, 2003 Apr; 45(4): 279-85. [PubMed]
 23. Kanungo, R., Rajalakshmi, B. (2001) Serotype distribution & antimicrobial resistance in Streptococcus pneumoniae causing invasive & other infections in south India. *Indian J Med Res.*, 114: 127-132. Google Scholar | Medline | ISI
 24. Awasthi, S., Nichter, M., Verma, T. Revisiting community case management of childhood pneumonia: perceptions of caregivers and grass root health providers in Uttar Pradesh and Bihar, Northern India. *PLoS ONE*, 2015; 10: e0123135. Google Scholar | Crossref | ISI
 25. Capoor, M., Nair, D., Aggarwal, P. Rapid diagnosis of community-acquired pneumonia using the BacT/Alert 3D system. *Braz J Infect Dis*, 2006; 10: 352-356. Google Scholar | Crossref | Medline
 26. Neuman MI, Hall M, Lipsett SC, Hersh AL, Williams DJ, Gerber JS, Brogan TV, Blaschke AJ, Grijalva CG, Parikh K, Ambroggio L, Shah SS; Pediatric Research in Inpatient Settings Network. Utility of Blood Culture Among Children Hospitalized with Community-Acquired Pneumonia. *Pediatrics*, 2017 Sep; 140(3): pii: e20171013. doi: 10.1542/peds.2017-1013. Epub 2017 Aug 23. [PubMed]
 27. Tiewsoh., Karalanglin Lodha . et al (2009); Factors determining the outcome of children hospitalized with severe pneumonia", *BMC Pediatrics*, 2009; 23(9): 15.