



**A COMPARATIVE STUDY TO ASSESS KNOWLEDGE, ATTITUDE AND PRACTICE
OF OTC DRUGS AMONG SECOND YEAR MEDICAL STUDENTS OF 2018 BATCH AND
2021 BATCH AT A TERTIARY CARE TEACHING HOSPITAL**

Dr. Pooja Agrawal¹, Dr. Virendra Kushwaha^{*1}, Dr. Vijay Kumar Singh², Dr. Vipul Shukla¹

¹Department of Pharmacology, GSVM Medical College, Kanpur, UP.

²Department of Pharmacology, Rani Durgawati Medical College, Banda, UP.

***Corresponding Author: Dr. Virendra Kushwaha**

Professor & Head, Department of Pharmacology, GSVM Medical College, Kanpur, UP.

Article Received on 31/05/2023

Article Revised on 20/06/2023

Article Accepted on 10/07/2023

ABSTRACT

Introduction: Over-the-counter (OTC) drugs are more commonly used than prescription drugs for self-medication worldwide¹. The Advantages of OTC drugs are decreased doctor visits and lower costs compared with prescription drugs. Overuse of OTC drugs can lead to adverse reactions, drug interactions, overdosing, and other medication-related issues. **Method:** A cross sectional survey is conducted among 2nd year medical students, in the form of pre-test containing questions about OTC drugs. After sensitisation on topic same questions were provided in the form of post-test and results were analysed. **Result:** Out of 250 students, 217(87%) students responded in pre-test while 220(88%) responded during post-test. During test, 39% were female and 61% were male. Most common ailment is fever for which self medication has been done. We found significant improvement in knowledge and change in attitude after sensitisation towards use of OTC drugs and majority of students (80%) agreed that self medication should not be done. **Conclusion:** We concluded that majority of participants practiced self-medication. Awareness and dangers of misuse of OTC medications among all participants was less. Therefore, it is suggested that proper education/awareness programmes should be implemented regarding self-administration of OTC drugs.

KEYWORDS: OTC drugs, self medication, prescription.

INTRODUCTION

Over-the-counter (OTC) drugs are more commonly used than prescription drugs for self-medication worldwide.^[1] OTC drugs are drugs that can be legally purchased without requiring a prescription from a registered medical practitioner. OTC drugs are primarily used for conditions, when there is no need for medical attention or supervision, and OTC drugs must be reasonably safe and well-tolerated.^[2-3]

Many people use OTC drugs as their first line of treatment, without formal prescription, because they are affordable and easily accessible from drug stores.^[4]

The Advantages of OTC drugs are decreased doctor visits and lower costs compared with prescription drugs. Overuse of OTC drugs can lead to adverse reactions, drug interactions, overdosing, and other medication-related issues. The public should use OTC drugs carefully to avoid adverse events associated with overuse of OTC drugs and increase their knowledge and understanding of potential dangers of overuse of OTC drugs.^[5-7]

OTC drugs are classified into many categories according to WHO Anatomical Therapeutic Chemical (ATC) classification. They include analgesics, laxatives, antacids, cough and cold preparations, antihistamines, dermatological, throat preparations, nasal preparations, and antidiarrheals.^[8] The Food, Drugs, and Cosmetic Act was implemented in 1938 which authorized the FDA to issue clear guidelines for drugs to be sold by prescription only and to be sold as OTC.^[9] In India, "over the counter drugs" has no legal definition, so all the drugs not included in the list of "prescription drugs" are considered as OTC drugs.

Many people use OTC drugs, without reading the directions to use them.^[10] Significant problems and malpractices were identified with OTC drugs like sharing of OTC drugs, expired medicine usage, doubling the drug dose when they are ineffective, OTC storage, and not reading labels and expiry dates.^[11] Therefore, government should introduce awareness programmes through television/newspaper/social media regarding proper use of OTC drugs and difference between OTC drugs and drugs which should be used with prescription only.

The aim of the study was, therefore, to assess the practice of self-medication with OTC drugs, the prevalence of the risky practice, and its associated factors.

MATERIAL AND METHOD

It is a cross sectional survey organised in Department of Pharmacology, GSVM Medical College, Kanpur, UP. As a part of undergraduate teaching first we organised a pre-test, and provided 15 questions on google form online (time - 20 minutes) about knowledge, attitude and practice regarding OTC drugs among second year medical students (2021 batch). In, the questionnaire we included five questions based on Likert scale, four questions were open ended, four questions were closed ended while rest two questions were based on attitude/practice towards OTC drugs. We recorded the answers, and then students were sensitised about OTC drugs. After sensitisation same questionnaire was provided online on google form in the form of post-test

(time - 20 minutes) and answers were analysed. Consent regarding questionnaire was taken from students.

RESULT

Out of 250 students, 217(87%) students responded in pre-test while 220(88%) responded during post-test. (Table 1)

Table 1.

	Total students	Responded	Not responded
Pre test	250	217(87%)	33(13%)
Post test	250	220(88%)	30(12%)

Gender and Age distribution of students during pre and post-test has been shown in Figure 1 and figure 2 respectively. During Pre and Post, both test 31% of participants were female while 69% were male. During both test about 45% students belong to age group of 21-22 years, while about 27% belong to 19-20 years and about 25% belong to 23-24 years age group.




Fig 1.




Fig 2.

Table 2 shows responses (based on Likert scale) of students over Knowledge/Awareness

Table 2.

Are you aware about OTC drugs	Not aware	Slightly aware	Somewhat aware	Moderately aware	Extremely aware
Pre test (n=217)	7	3	90	75	42
%	(3)	(1)	(41)	(35)	(20)
Post test (n=220)	4	2	5	90	119
%	(2)	(1)	(2)	(41)	(54)

Do you know about drugs included in OTC list	Not aware	Slightly aware	Somewhat aware	Moderately aware	Extremely aware
Pre test (n=217) %	23 (10)	46 (21)	84 (39)	30 (14)	34 (16)
Post test (n=220) %	13 (6)	22 (10)	38 (17)	104 (47)	43 (20)
Have you read the instruction on the package insert before using the OTC drug	Never	Almost never	Occasio-nally	Almost every time	Always
Pre test (n=217) %	36 (17)	51 (24)	84 (39)	29 (13)	17 (7)
Post test (n=220) %	28 (12)	18 (9)	45 (20)	84 (39)	45 (20)
How much did you understand the instruction	Never	Almost never	Occasio-nally	Almost every time	Always
Pre test (n=217) %	58 (27)	44 (20)	49 (23)	45 (20)	21 (10)
Post test (n=220) %	20 (9)	36 (16)	13 (6)	93 (42)	58 (27)
How frequently you took self-medication/ OTC drug	Never use	Almost never	Occasio-nally/ Sometimes	Almost every time	Frequently
Pre test (n=217) %	49 (23)	31 (14)	61 (28)	34 (16)	42 (19)
Post test (n=220) %	5 (22)	39 (18)	64 (29)	34 (15)	33 (19)

Table 2 shows that during pre- test 76% students either somewhat/moderately aware about OTC drugs while during post-test 95% students were moderately/Extremely aware. Table also shows that during pre-test maximum number of students were neither aware about drugs included in OTC list nor they

were reading the instructions written over package. However frequency of using OTC drugs was almost similar during both pre and post test.

Figure 3 shows the disease/ medical condition for which OTC drugs have been used




Fig 3.

Figure 3 shows that maximum use of OTC drugs was for fever (22%) followed by skin problems (16%), Acidity

(14%) and Headache (11%). Response was almost same during pre and post test.




Figure 4: Shows the source of information about OTC drugs.

Fig 4 shows that during pre-test source of information about OTC was Peer(27%), Lecture(26%), self learning(21%), text books(17%) and 7% from media

advertisements. During post test 71% students reported that source of information was Lecture.




Figure 5: Shows the sources of obtaining OTC drugs for self-medication.

Fig 5 shows that medical store(55%) was the main source of obtaining the OTC drugs during pre and post test, while other sources were online shopping(18%), left

over medicines(15%) and about 11% obtained from friends/relatives.




Figure 6 shows the reason for using OTC drugs.

Fig 6 shows that (during both pre and post test) maximum students used OTC medicines because visit to a doctor was costly (about 40%), while 35% think that for

minor ailments, why to visit a doctor. About 18% used OTC based on previous experience of using OTC drugs.

Table 3 shows the responses of students towards open questions with closed answers.

Do you easily get antibiotics/ anti-anxiety drugs without prescription	Yes, Always	Yes, Sometimes	Never
Pre test (n=217)	113	73	31
Post test (n=220)	115	71	34
Did you get relief upon self-medication	Yes, Always	Yes, Sometimes	Never
Pre test (n=217)	98	77	42
Post test (n=220)	102	76	42
Have you experienced any adverse effects	Yes, Always	Yes, Sometimes	Never
Pre test (n=217)	43	124	50
Post test (n=220)	51	120	49
What did you do for the adverse reaction	Stop taking medication	Consulted pharmacy staff	Consulted a doctor
Pre test (n=217)	156	20	41
Post test (n=220)	152	16	52

Table 3 shows that about 50% students get antibiotic/antianxiety easily as OTC drugs during both pre and post test, while 34% sometimes and 14% never get these medicines. About 45% of them get relieve always, 35% sometimes and 20% never get relieved during both the tests. About 20% of students get always adverse effects, 57% get sometimes and 23% never have

adverse effects. On getting adverse effect about 70% stopped medication, 10% consulted a pharmacist/ healthcare staff and 20% consulted a doctor.

Figure 7 and 8 shows responses of students towards practice- based questions over OTC Drugs.




Fig 7.

Figure 7 shows that during pretest 62% students think that OTC drug use is a good practice, 15% think it acceptable while 23% think it to be not acceptable.

During posttest 65% think it not acceptable, 20% acceptable and 15% thinks it is a good practice.




Fig 8.

Figure 8 shows that during pretest 56% students think that common ailments should be treated by self, 28% were not sure and 16% think that it should never be done. Posttest shows that only 20% students agreed that it should be done, 16% were not sure and 64% think that it should never be done.

DISCUSSION

Use of OTC drugs is very common among Indians, without having knowledge about their indication or adverse effects. Several studies have shown that resistance of pathogens, adverse drug reactions and drug dependence increase to a very high level due to this inappropriate use of drugs without expert opinion.^[12-14]

This study was conducted among second year medical students and maximum participants belong to age group of 21-22 years(45%), while about 27% belong to 19-20 years and about 25% belong to 23-24 years age group. Out of them 31% were female and 69% were male.

This study shows that during pre- test 76% students were either somewhat/moderately aware about OTC drugs while during post-test 95% students were moderately/Extremely aware. About 63% of students were using OTC drugs. Another questionnaire based study had been conducted by our department with students of 2018 batch shows that majority (98.90%) of the students were aware of OTC drugs and self-medication was seen among (95.09%) of the medical students.^[15] In another study conducted among medical students in Karnataka revealed that 53% students practicing self medication^[16], while study conducted by Mourya et al stated that the overall prevalence of self-medication was 77.33%.^[17]

This study shows that maximum use of OTC drugs was for fever (22%) followed by skin problems (16%), Acidity (14%) and Headache (11%). Same departmental study during 2020 stated that the most common ailment for which maximum number of students have taken OTC

was Fever (36.07%).^[15] Mourya et al shows that Highest % of indication for self-medication was found to be fever (31.551%).^[17] Another study in South India revealed that cough and cold are the most common reason for OTC drugs.^[18]

This study shows that (during both pre and post test) maximum students used OTC medicines because visit to a doctor was costly(about 40%), while 35% think that for minor ailments, why to visit a doctor. Previous study in our department stated that most common reason for self-medication or use of OTC drug was “for minor ailment why see Dr advice” (55.19%) followed by the reason that “Going to doctor were cumbersome (12.02%) and “free availability in market” (22.96%).^[15] Maurya et al stated that the most common reason for self-medication was easy accessibility (21.55%) followed by self-confidence (17.24%), time saving (16.03%), no enough time for consultation (7.07%), safe and well tolerated (5.69%), low cost (3.28%).^[17]

In our study we found that students easily get (around 80%) antibiotics or even antianxiety drugs, and most of the times they get relieved(80%) by self medication with incidence of adverse effects was found always associated in 19% and sometimes in 57% of participants. Previous study in our department showed that only 16.54% students reported any adverse drug reaction.^[15] Another study reported that 18.62% of participants experienced side effects from OTC drugs.^[17]

This study shows that during pretest 62% students think that OTC drug use is a good practice, 15% think it acceptable while 23% think it to be not acceptable. After sensitisation their attitude changed towards OTC drugs and in posttest 65% think it is not acceptable, 20% acceptable and only 15% thinks it is a good practice. Previous study of our department showed that maximum students (68%) think that self-medication with OTC is a good practice.^[15] Maurya et al also showed that majority

of the participants believed that use of OTC drug is a good practice.^[17]

This study shows that during pretest 56% students think that common ailments should be treated by self, 28% were not sure and 16% think that it should never be done. Posttest shows that only 20% students agreed that it should be done. Previous study shows that 44% think that common ailments can be treated by using OTC drugs.^[15] We can clearly observe that after sensitisation there is change in attitude towards use of OTC drugs. Adequate counselling/education can improve self-medication practice and made them understand how it will improve their quality of life and in better management of the disease.

CONCLUSION

This cross-sectional study showed that use of OTC drugs is very common among medical students, facilitated by the easy availability of drugs. Majority of the them believed, use of OTC drug is a good practice, but their knowledge about OTC is inadequate, they are not aware about correct drug indication, what is the ideal duration for the drugs, what is the dose regimen of the drugs and when to stop the drugs. Excessive use of non OTC drugs as OTC can lead to extra burden of cost and development of resistance. The scenario among general population may be worse leading to more complications and more adverse events associated with excessive/inappropriate use of drugs. So, there is need to implement strict laws by the government about categorisation/dispensing of OTC/nonOTC drugs, conducting awareness programs and restricting drug advertisements for public.

REFERENCE

1. World Health Organization. The role of the pharmacist in self – care and self-medication. Report of the 4th WHO Consultative Group on the Role of the Pharmacist. The Hague, 1998. Cited on 20/04/23. Available from: <http://www.who.int/medicines/library/dap/who-dap-98-13/who-dap-98-13.pdf>.
2. Pawaskar MD, Balkrishnan R. Switching from prescription to over the counter medications: a consumer and managed care perspective. *Manag Care Interface*, 2007; 20: 40–47.
3. WHO Guideline for the Regulatory Assessment of Medicinal Products for Use in Self- Medication, 2000. Cited on 20/02/23. Available from: www.who.int/medicines/library/qsm/who-edm-qsm-2000-1/who-edm-qsm-00_1.ht
4. Ocan M, Bwanga F, Bbosa GS, Bagenda D, Waako P, Ogwal-Okeng Jet al Obua. Patterns and predictors of self-medication in northern Uganda. *PLoS One.*, 2014; 9(3): e92323.
5. Alshahrani SM, Alakhali KM, Al-Worafi YM, Najim Z, Alshahrani NZ. Awareness and use of over the counter analgesic medication: a survey in the Aseer region population, Saudi Arabia. *Int J Advan Appl Sci.*, 2020; 7(3): 130–134.
6. Kagashe GA, Francis L. Dispensing of drugs with and without a prescription from private pharmacies in Dar es Salaam. *Tanzania Med J.*, 2004; 19(1): 36–39.
7. Alghanim SA. Self-medication practice among patients in a public health care system. *East Mediterr Health J.*, 2011; 17(5): 409–416.
8. WHO Collaborating Centre for Drug Statistics Methodology, Guidelines for ATC classification and DDD assignment, 2023. Oslo, 2022. Cited on 30/04/2023.
9. Marathe PA, Kamat SK, Tripathi RK, Raut SB, Khatri NP. Over-the-counter medicines: Global perspective and Indian scenario. *J Postgrad Med.*, 2020; 66(1): 28-34.
10. Barrenberg E, Garbe E. Use of over-the-counter (OTC) drugs and perceptions of OTC drug safety among German adults. *Eur J Clin Pharmacol*, 2015; 71(11): 1389-96.
11. Bekele KM, Abay AM, Mengistu KA, Atsbeha BW, Demeke CA, Belay WS, et al. Knowledge, attitude, and practice on over-the-counter drugs among pharmacy and medical students: A facility-based cross-sectional study. *Integr Pharm Res Pract.*, 2020; 9: 135-46.
12. Badiger S, Kundapur R, Jain A, Kumar A, Pattanshetty S, Thakolkaran N et al. Self medications pattern among medical students in South India. *Australasian Medical Journal*, 2012; 5(4): 217-20.
13. James H, Handu SS, Khalid AJ, Khaja A, Otoom S, Sequeira RP. Evaluation of the Knowledge, Attitude and Practice of Self-Medication among first-Year Medical Students. *Med Princ Pract*, 2006; 15: 270–275.
14. Hem E, Stokke G, Reidar Tyssen R, Grønvold NT, Vaglum P, Ekeberg O. Self-prescribing among young Norwegian doctors: a nine-year follow-up study of a nationwide sample. *BMC Med.*, 2005; 3: 16.
15. Agrawal P, Kushwaha V, Tripathi M, Singh S and Barman SK. *EJPBS*, 2020; 7(8): 538-46.
16. G. K. Nalini. Self-Medication among Allopathic medical Doctors in Karnataka, India. *BJMP*, 2010; 3: 325
17. Mourya A, Mary C, James C, Jose J, Srinivasan R. A Survey on Over the Counter Drug usage in the Community. *Journal of Drug Delivery & Therapeutics*, 2019; 9(2): 406-16.
18. Gaikwad NR, Patil AB, Khan TA. Comparative evaluation of knowledge, attitude and practice of self-medication among first and second year medical students. *J Datta Meghe Inst Med Sci Univ.*, 2010; 5: 157–62.