



**ANTI-HYPERURICEMIA ACTIVITY OF SOME MEDICINAL PLANTS FROM
KARAWANG, WEST JAVA, INDONESIA: A REVIEW**

Zafira Fatiha Nurulhadi, Siti Mudrikah, Tanti Amelia, Della Putri Valentina, Iin Kurniawati, Reza Rizky Yuniar, Dida Fahdona Azzahra, Tiurida Pandiangan, Silvia Ismayanti, Erna Sri Rahmawati, Ferdyan Pranata Saputra, Helsen, Ilham Bintang Pratama, Deden Sukandar and Maulana Yusuf Alkandahri*

Faculty of Pharmacy, Universitas Buana Perjuangan Karawang, Karawang, West Java, Indonesia.



*Corresponding Author: Maulana Yusuf Alkandahri

Faculty of Pharmacy, Universitas Buana Perjuangan Karawang, Karawang, West Java, Indonesia.

Article Received on 04/01/2024

Article Revised on 25/01/2024

Article Accepted on 15/02/2024

ABSTRACT

The utilization of various botanical species for the treatment of anti-uricemia has garnered interest as part of the search for alternative therapies. This review provides details regarding the potential therapeutic effects of specific botanical species in mitigating hyperuricemia. Treating hyperuricemia is primarily complicated by the limited efficacy of conventional therapies and the condition's high prevalence. Age is a common determinant of hyperuricemia; inactivity and an unhealthy diet are potential risk factors. A number of studies have additionally emphasized the correlation between uric acid accumulation in the body and genetic factors, physical activity, and diet. Up-to-date information regarding medicinal plants and their potential as anti-hyperuricemia treatments is gathered through a systematic search of multiple sources of data. As a result, further investigation into the use of medicinal plants as an alternative treatment for hyperuricemia is a compelling area of study. A number of botanical species indigenous to Karawang, West Java, Indonesia, have exhibited promise as uric acid inhibitors.

KEYWORDS: Anti-hyperuricemia, Uric acid, Medicinal plants, Karawang, Indonesia.

INTRODUCTION

At present, the global incidence of hyperuricemia is estimated to be between 0.1 and 0.3%, while the prevalence ranges from 1% to 4%. Males are significantly more susceptible to hyperuricemia than females, by a ratio of 3:1 to 10:1. The occurrence and prevalence of hyperuricemia exhibit a consistent upward trend as individual age. Specifically, the prevalence rises by 11–13% in the over-80 age group, while the incidence rises to 0.4%.^[1] Despite the fact that this condition can be effectively managed with currently available medications, these options have numerous adverse effects.^[2,3] Low patient adherence to treatment also contributes to the recurrence of elevated uric acid levels.^[4,5] Hyperuricemia is the main cause of gout arthritis. Monosodium urate (MSU) crystal deposition in and around the joints is the primary cause of this inflammatory arthritis disease.^[6] Globally, the utilization of medicinal plants to treat a variety of ailments has increased due to their reputation for being considerably safer than synthetic drugs.^[7,8] Active compound explosion from natural sources, especially medicinal plants that have been used to treat hyperuricemia in the past in different countries, is still one way that researchers are trying to find new natural medicines that can treat hyperuricemia.^[9,10] The objective is to identify novel anti-hyperuricemia compounds

characterized by minimal toxicity and mild side effects so as to prevent harm to patients.^[11,12] This review article will discuss active compounds and anti-hyperuricemia properties of some medicinal plants in Karawang, West Java, Indonesia.

Curcuma longa

Curcuma longa, a member of the *Zingiberaceae* family, is a medicinal plant. This plant is cultivated extensively throughout Asia, particularly in China and India. *C. longa* is a sterile plant that lacks the ability to produce seeds. This plant attains a height of 3 to 5 feet and produces yellow flowers. The rhizome is an underground stem that is thick and fleshy.^[13] It has been widely reported that *C. longa* is a medicinal plant with anti-inflammatory and analgesic properties. This effect is due to the presence of curcuminoids, which are at least three natural polyphenols found in this plant: curcumin, demethoxycurcumin, and bisdemethoxycurcumin.^[14,15] Giving mice 20 to 40 mg/kg/day of curcumin-containing *C. longa* for 14 days lowered their uric acid levels when they are exposed to potassium oxonate.^[16] In vitro studies, curcumin found in *C. longa* can stop the enzymes xanthine oxidase (XO) and urate transporter 1^[17] with an IC₅₀ value of 3.33 ± 0.52 μM.^[18] In addition, *C. longa* is also reported to increase uricosuric activity,

which can result in increased renal excretion of uric acid.^[19]

Apium graveolens

Apium graveolens is a plant belonging to the *Umbelliferae* family. *A. graveolens* seeds contain various substances, such as essential oils, flavonoids, coumarin, and linoleic acid. *A. graveolens* seeds have long been used to treat arthritis, uric acid, and help reduce muscle spasms, calm nerves, and reduce inflammation.^[20] According to reports, potassium oxonate-induced mice has lower uric acid levels after receiving *A. graveolens* at a dose of 1,000 mg/kg for two weeks.^[21] Apart from that, in tests carried out in vitro, *A. graveolens* is reported to be able to inhibit the xanthine oxidase enzyme at a concentration of 1,500 mg/L.^[22]

Orthosiphon stamineus

Orthosiphon stamineus is a traditional medicinal plant that has various properties for treating various diseases, including kidney nephritis, cancer, and arthritis. Triterpenes, betulinic acid, oleanolic acid, ursolic acid, β -sitosterol, and more than 20 other phenolic compounds are said to be bioactive and found in this plant.^[23] Administration of *O. stamineus* at doses of 500, 1000, and 2000 mg/kg can significantly reduce the serum uric acid of mouse, which is induced by potassium oxonate.^[24] On the other hand, tests done in vitro showed that *O. stamineus* can stop the xanthine oxidase enzyme working, with IC_{50} values between 3.94 and 8.76 mg/mL.^[24] Apart from that, *O. stamineus* has also been reported to inhibit the expression of inflammatory mediators such as iNOS, COX-2, TNF- α , and PGE2.^[25] Therefore, this plant can be used as an alternative treatment for gouty arthritis caused by hyperuricemia.

Kaempferia galanga

Kaempferia galanga is an herbaceous plant renowned for the culinary and traditional medicinal uses of its rhizomes. This plant produces flowers and green leaves that protrude above the soil. *K. galanga* rhizome is utilized in numerous traditional medicines due to its fragrant aroma and purported anti-inflammatory, antimicrobial, and antioxidant properties.^[26] *K. galanga* is reported to have quite high total phenolic content, namely 50.35 mgGAE/100 g.^[27] In tests carried out in vitro, *K. galanga* is reported to inhibit the xanthine oxidase enzyme with an IC_{50} value of 139.92 μ g/mL.^[27]

Annona muricata

Annona muricata tropical is characterized by oval, dark green leaves and white, spiny green-skinned fruits. The fruit of this plant offers a combination of sweet and sour flavors, rendering it a frequent ingredient in juice, ice cream, and other culinary preparations.^[28] The *A. muricata* plant is reported to have properties for treating several diseases, such as cancer, uric acid, tumors, hypertension, diabetes mellitus, ulcers, diarrhea, and allergies.^[29] Administration of *A. muricata* at a dose of 75 mg/kg can reduce the serum uric acid of rats induced

by potassium oxonate.^[30] Apart from that, in tests carried out in vitro, *A. muricata* is reported to inhibit the xanthine oxidase enzyme with an IC_{50} value of > 200 μ g/mL.^[30]

Psidium guajava

Psidium guajava, more commonly referred to as guava, is a long-standing medicinal plant with a rich historical background in tropical regions, including Indonesia. *P. guajava* is a member of the *Myrtaceae* family and possesses a variety of therapeutic properties, including analgesic, anti-inflammatory, anti-diabetic, anti-hypertensive, antioxidant, antibacterial, and antitumor.^[31,32] *P. guajava* can lessen the potassium oxonate-induced uric acid of mouse serum when administered at doses of 5, 10, and 20 mL/kg/day.^[33] Besides, tests done in vitro showed that *P. guajava* can stop the xanthine oxidase enzyme working, with an IC_{50} value of 38.24 μ g/mL.^[31]

Zingiber officinale

Zingiber officinale is a spice that has been utilized historically for both culinary and medicinal purposes. *Z. officinale*, a botanical species classified within the *Zingiberaceae* family, has a rich history of utilization as an immunomodulatory, anticancer, nephroprotective, hepatoprotective, anti-microbial, and anti-diabetic agent.^[34] Chicken liver juice induces the production of uric acid in mouse serum, which *Z. officinale* can inhibit at a dose of 100 mg/kg.^[35] Furthermore, it has been reported that *Z. officinale* inhibits the xanthine oxidase enzyme in vitro at an IC_{50} value of 188.5 μ g/mL.^[36]

Andrographis paniculata

Andrographis paniculata is a medicinal plant that is empirically used to treat a variety of conditions, including respiratory ailments, diabetes, cancer, obesity, skin infections, herpes, dysentery, fever, sore throat, urinary tract infections, diarrhea, and inflammation.^[37] This plant has the main content of andrographolide lactone group compounds \pm 2.5% in dried simplicial.^[38] Administration of *A. paniculata* at doses of 50, 100, and 200 mg/kg can reduce the serum uric acid of rats induced by potassium oxonate.^[39] Besides, in tests carried out in vitro, *A. paniculata* is reported to inhibit the xanthine oxidase enzyme at a concentration of 400 μ g/mL.^[39]

CONCLUSION

Indonesia is home to a diverse array of medicinal plants that possess promising attributes that can be harnessed to formulate alternative therapies for hyperuricemia. It has been empirically and scientifically demonstrated that each of these plants can reduce uric acid levels. This is inextricably linked to the active compounds present in these plants, which employ various mechanisms to exert their anti-uricemic effects. Research on this botanical species holds promise for enhancing contemporary anti-uricemia treatments, which are presently characterized by a multitude of detrimental side effects.

REFERENCES

- Roddy E, Choi HK. Epidemiology of gout. *Rheum Dis Clin North Am.*, 2014; 40(2): 155-175.
- Dalbeth N, Merriman TR, Stamp LK. Gout. *Lancet.*, 2016; 388(10055): 2039-2052.
- Alkandahri MY, Sholih MG, Fadilah NN, Arfania M, Amal S, Frianto D, *et al.* Evaluation of antidiarrheal, antispasmodic, and antisecretory activities of extract and fractions of *Castanopsis costata* leaves in animal models. *Pharmacogn J.*, 2023; 15(1): 31-37.
- Neogi T, Hunter DJ, Chaisson CE, Allensworth-Davies D, Zhang Y. Frequency and predictors of inappropriate management of recurrent gout attacks in a longitudinal study. *J Rheumatol*, 2006; 33(1): 104-109.
- Arfania M, Frianto D, Astuti D, Anggraeny EN, Kurniawati T, Alivian R, *et al.* Measurement of adherence level of pulmonary tuberculosis drugs use in patients in the primary health centers in Karawang Regency, West Java, Indonesia, using MMAS instrument. *J Pharm Res Int.*, 2021; 33(54A): 115-120.
- Stewart DJ, Langlois V, Noone D. Hyperuricemia and hypertension: Links and risks. *Integr Blood Press Control.*, 2019; 12: 43-62.
- Alkandahri MY, Berbudi A, Subarnas A. Active compounds and antimalaria properties of some medicinal plants in Indonesia – A review. *Sys Rev Pharm.*, 2018; 9(1): 64-69.
- Alkandahri MY, Maulana YE, Subarnas A, Kwarteng A, Berbudi A. Antimalarial activity of extract and fractions of *Cayratia trifolia* (L.) Domin. *Int J Pharm Res.*, 2020; 12(1): 1435-1441.
- Alkandahri MY, Kusumiyati K, Renggana H, Arfania M, Frianto D, Wahyuningsih ES, *et al.* Antihyperlipidemic activity of extract and fractions of *Castanopsis costata* leaves on rats fed with high cholesterol diet. *RASĀYAN J Chem.*, 2022; 15(4): 2350-2358.
- Alkandahri, MY, Arfania M, Abriyani E, Ridwanuloh D, Farhamzah, Fikayuniar L, *et al.* Evaluation of antioxidant and antipyretic effects of ethanolic extract of Cep-cepan leaves (*Castanopsis costata* (Blume) A.DC). *J Adv Pharm Educ Res.*, 2022; 12(3): 107-112.
- Alkandahri MY, Berbudi A, Utami NV, Subarnas A. Antimalarial activity of extract and fractions of *Castanopsis costata* (Blume) A.DC. *Avicenna J Phytomed.*, 2019; 9(5): 474-481.
- Alkandahri MY, Yuniarsih N, Berbudi A, Subarnas A. Antimalaria activities of several active compounds from medicinal plants. *Pharmacogn J.*, 2022; 14(1): 245-252.
- Verma RK, Kumari P, Maurya RK, Kumar V, Verma RB, Singh RK. Medicinal properties of turmeric (*Curcuma longa* L.): A review. *Int J Chem Stud.*, 2018; 6(4): 1354-1357.
- Alkandahri, MY., Patala, R., Berbudi, A., and Subarnas, A. Antimalarial activity of curcumin and kaempferol using structure based drug design method. *J Adv Pharm Educ Res.*, 2021; 11(4): 86- 90.
- Alkandahri MY, Berbudi A, Subarnas A. Evaluation of experimental cerebral malaria of curcumin and kaempferol in *Plasmodium berghei* ANKA-infected mice. *Pharmacogn J.*, 2022; 14(6): 905-911.
- Chen Y, Li C, Duan S, Yuan X, Liang J, Hou S. Curcumin attenuates potassium oxonate-induced hyperuricemia and kidney inflammation in mice. *Biomed Pharmacother*, 2019; 118: 1-7.
- Hu Q, Zhou M, Zhu H, Lu G, Zheng D, Li H, *et al.* (E)-2-(4-bromophenyl)-1-(2, 4-dihydroxyphenyl) ethanone oxime is a potential therapeutic agent for treatment of hyperuricemia through its dual inhibitory effects on XOD and URAT1. *Biomed Pharmacother*, 2017; 86: 88-94.
- Ao GZ, Zhou MZ, Li YY, Li SN, Wang HN, Wan Q, *et al.* Discovery of novel curcumin derivatives targeting xanthine oxidase and urate transporter 1 as anti-hyperuricemic agents. *Bioorg Med Chem.*, 2017; 25(1): 166-174.
- Bupparenoo P, Pakchotanon R, Narongroeknawin P, Asavatanabodee P, Chaiamnuy S. Effect of curcumin on serum urate in asymptomatic hyperuricemia: A randomized placebo-controlled trial. *J Diet.*, 2021; 18(3): 248-260.
- Al-Asmari AK, Athar MT, Kadasah SG. An updated phytopharmacological review on medicinal plant of Arab Region: *Apium graveolens* Linn. *Pharmacogn Rev.*, 2017; 11(21): 13-18.
- Dolati K, Rakhshandeh H, Golestani M, Forouzanfar F, Sadeghnia R, Sadeghnia HR. Inhibitory effects of *Apium graveolens* on xanthine oxidase activity and serum uric acid levels in hyperuricemic mice. *Prev Nutr Food Sci.*, 2018; 23(2): 127-133.
- Iswantini D, Nadinah, Darusman LK, Trivadila. Inhibition kinetic of *Apium graveolens* L. ethanol extract and its fraction on the activity of xanthine oxidase and its active compound. *J Biol Sci.*, 2012; 12: 51-56.
- Ashraf K, Sultan S, Adam A. *Orthosiphon stamineus* Benth. is an outstanding food medicine: Review of phytochemical and pharmacological activities. *J Pharm Bioallied Sci.*, 2018; 10(3): 109-118.
- Xu WH, Wang HT, Sun Y, Xue ZC, Liang ML, Su WK. Antihyperuricemic and nephroprotective effects of extracts from *Orthosiphon stamineus* in hyperuricemic mice. *J Pharm Pharmacol*, 2020; 72(4): 551-560.
- Yam MF, Lim V, Salman IM, Ameer OZ, Ang LF, Rosidah N, *et al.* HPLC and anti-inflammatory studies of the flavonoid rich chloroform extract fraction of *Orthosiphon stamineus* leaves. *Molecules*, 2010; 15(6): 4452-4466.
- Kumar A. Phytochemistry, pharmacological activities and uses of traditional medicinal plant *Kaempferia galanga* L. - An overview. *J Ethnopharmacol*, 2020; 253: 1-19.

27. Julianti TB, Bakar MF, Wikantyasning ER. Phytochemical, antioxidant analysis and in vitro xanthine oxidase inhibitory activity of *Kaempferia parviflora* and *Kaempferia galanga*. *Trop J Nat Prod Res.*, 2022; 6(12): 1981-1985.
28. Balderrama-Carmona AP, Silva-Beltrán NP, Gálvez-Ruiz JC, Ruíz-Cruz S, Chaidez-Quiroz C, Morán-Palacio EF. Antiviral, antioxidant, and antihemolytic effect of *Annona muricata* L. leaves extracts. *Plants (Basel)*., 2020; 9(12): 1-11.
29. Mutakin M, Fauziati R, Fadhilah FN, Zuhrotun A, Amalia R, Hadisaputri YE. Pharmacological activities of soursop (*Annona muricata* Lin.). *Molecules.*, 2022; 27(4): 1-17.
30. Sunarni T, Leviana F, Fidrianny I, Iwo MI, Wirasutisna KR. Antihyperuricemic activity of four plants Annonaceae using hyperuricemic rats model and enzyme assay. *Asian J Pharm Clin Res.*, 2015; 8(6): 250-253.
31. Irondi EA, Agboola SO, Oboh G, Boligon AA, Athayde ML, Shode FO. Guava leaves polyphenolics-rich extract inhibits vital enzymes implicated in gout and hypertension in vitro. *J Intercult Ethnopharmacol*, 2016; 5(2): 122-130.
32. Ugbo EA, Emmanuel O, Uche ME, Dike ED, Okoro BC, Ibe C, *et al.* The ethnobotanical, phytochemistry and pharmacological activities of *Psidium guajava* L. *Arabian J Chem.*, 2022; 15(5): 1-26.
33. Aprillinda FWD, Paramasari D, Wasita B. The effect of red guava juice (*Psidium guajava* Linn.) in decreasing uric acid and creatinine levels of hyperuricemic white mice (*Mus musculus*). *Bali Med J.*, 2018; 7(2): 323-329.
34. Mao QQ, Xu XY, Cao SY, Gan RY, Corke H, Beta T, *et al.* Bioactive compounds and bioactivities of ginger (*Zingiber officinale* Roscoe). *Foods.*, 2019; 8(6): 1-21.
35. Afriani T, Yulion R, Andriani M, Syufyani F, Fadri D. The effect of Ginger (*Zingiber officinale* Roscoe) fractionation in decreasing uric acid level of hyperuricemic white mice. *International Conference on Health Informatics and Medical Application Technology*, 467-474.
36. Muthusamy S, Jeyabalan S. In-silico and in-vitro evaluation of xanthine oxidase inhibition of *Zingiber officinalae* for hypouricemic activity. *Research J Pharm Tech.*, 2019; 12(1): 314-320.
37. Dai Y, Chen SR, Chai L, Zhao J, Wang Y, Wang Y. Overview of pharmacological activities of *Andrographis paniculata* and its major compound andrographolide. *Crit Rev Food Sci Nutr.*, 2019; 59(1): S17-S29.
38. Resi EM. Effect of antimalaria herbal Sambiloto (*Andrographis paniculata* Nees.) on morphology changes of development and parasite *Plasmodium falciparum*. *Jurnal Info Kesehatan*, 2014; 12(1): 661-669.
39. Rahmi EP, Kumolosasi E, Jalil J, Buang F, Jamal JA. Extracts of *Andrographis paniculata* (Burm.f.) Nees leaves exert anti-gout effects by lowering uric acid levels and reducing monosodium urate crystal-induced inflammation. *Front Pharmacol*, 2022; 12: 1-12.