

Measures	Indiana Health System	Florida A&M	National Avg.	Missouri
Patients discharged on medication within 30 days after last health care visit, based on patients' blood sugar and cholesterol level percentages in the past year	 68%	 56%	 73%	Higher than Florida 5 % Lower 5 % Higher that national average. Data Source: IPEDS-150000
Patients assessed and given insurance information	 66%	 70%	 73%	Higher than Florida 27 % Higher 22 % Higher that national average. Data Source: IPEDS-150000

Measures	Before Health System	Florida Age	National Age	Men's Life
Patients who visited about who received treatment for alcohol abuse (significant p)	26%	27%	40%	Highest in Nation 21% Higher (vs. Florida Average) 20% Higher (vs. National Average) Data Range: 1990-2000
Patients who visited about who received treatment for alcohol abuse (not significant p)	27%	40%	27%	Highest in Nation 17% Higher (vs. Florida Average) 20% Higher (vs. National Average) Data Range: 1990-2000
Patients who presented problem for alcohol abuse or drug abuse (not significant p)	46%	52%	25%	Highest in Nation 28% Higher (vs. Florida Average) 25% Higher (vs. National Average) Data Range: 1990-2000
Patients who presented problem for alcohol abuse or drug abuse (not significant p)	42%	25%	14%	Highest in Nation 27% Higher (vs. Florida Average) 46% Higher (vs. National Average) Data Range: 1990-2000
Patients who are Hispanic and received treatment for alcohol abuse or drug abuse (not significant p)	46%	25%	22%	Highest in Nation 12% Higher (vs. Florida Average) 15% Higher (vs. National Average) Data Range: 1990-2000
Patients who are Hispanic and received treatment for alcohol abuse or drug abuse (not significant p)	46%	26%	40%	Highest in Nation 21% Higher (vs. Florida Average) 21% Higher (vs. National Average) Data Range: 1990-2000
Patients who are Hispanic and received treatment for alcohol abuse or drug abuse (not significant p)	36%	40%	15%	Highest in Nation 10% Higher (vs. Florida Average) 20% Higher (vs. National Average) Data Range: 1990-2000
Patients who are Hispanic and received treatment for alcohol abuse or drug abuse (not significant p)	42%	25%	14%	Highest in Nation 27% Higher (vs. Florida Average) 46% Higher (vs. National Average) Data Range: 1990-2000

Measure	Indian Health System	Florida Reg.	National Reg.	Main Issue
Shows that patients spend in physical medicine for every 1,000 hours of patient care				Lower is Better 0.95 Higher than Florida average 0.7 Higher than National average Date Range: 10/01/00-10/01/00
	1.00	0.99	0.83	
Shows that patients spend in activities for every 1,000 hours of patient care				Lower is Better 0.16 Higher than Florida average 0.01 Lower than National average Date Range: 10/01/00-10/01/00
	0.07	0.0	0.06	

[illegible]

Measure	Indiana Health System	National Benchmark	More Info
Patients readmitted to any hospital within 30 days of discharge from the inpatient psychiatric facility	 Warren Group an Indiana Health System	26.93%	Lower is Better Best Range 1/2000 - 1/10,000