




Minnesota State Horticultural Society Records.

Copyright Notice:

This material may be protected by copyright law (U.S. Code, Title 17). Researchers are liable for any infringement. For more information, visit www.mnhs.org/copyright.

Rash Seedling No 1 Lancaster Minn

- Size.— 4 to 5 wgt 5 oz
Form.— oblate tapering a little toward
the eye very slightly angular
Color.— yellow ground marked deep
and stripes and splashes, carmine
Cavity.— broad open rather deep red
Stem.— medium Calyx closed
Basin.— broad and shallow mirror smooth
Flesh.— light yellow tender juicy
Flavor.— burk subacid very good
Season.— Jan to later
Free, Etc.—




Described.— Jan 27th 1899

Barb Seedling No 2 Laneshore Minn

- Size.— 5 Weight 6oz
Form.— Smooth round oboate
 slightly oblique
Color.— yellow ground with thin blush
 in sun, scattering brown dots over
Cavity.— narrow rather deep
Stem.— long; calyx partly open
Basin.— medium wrinkled
Flesh.— yellow tender nearly fine
Flavor.— rich, sweet, good

Season.— January
Free, Etc.—



Described.— Jan 27th 1899

A seedling on root sprout from Willow Lumber Tree

Size.— 4 to 5

Form.— round or globular

Color.— when ripe light greenish yellow greenish
and light and deep red stripes and blotches

Cavity.— medium rather deep, regular russeted at bottom

Stem.— very long rather slender elastic berries to bunch and

Basin.— medium slightly ribbed on summit calyx closed

Flesh.— white fine grained firm juicy


Flavor.— acid best for cooking

Season.— Jan to April

Free, Etc.— moderately hard, heavy berries in alternate years
Grown by J. S. Harris

Described.— Jan 12th 1901

Northwestern Grinning





Minnesota

State Horticultural Society.

President, W. W. PENDERGAST, Hutchinson.
Secretary, A. W. LATHAM, 207 Kasota Block, Minneapolis.

OFFICE OF

J. S. HARRIS,

Member of Executive Board.

Send to the Secretary for folder describing
the Society and its work and copy
of its Magazine.

PUBLISHES A MONTHLY MAGAZINE,
ENTITLED

"The Minnesota Horticulturist"

Subscription \$1.00 per annum,
including membership.

La Crescent, Minn., 1899

Long John by L. H. Alling Homer
Size 3 weight 5 oz
Form round inclining to pyriform
color yellowish green nearly
covered with deep and lighter
red in light stripes below
Stem long cavity regular small for
size of the fruit calyx partly open
basin broad medium deep wrinkled
flesh white fine grained flavor brisk
Subacid good Season November
The tree is from the root of a Farmhouse that killed
to the ground in 1884-5



J. S. Harris,
La Crescent,
Minn.

LA CROSSE BOOK & STATIONERY CO,

LA CROSSE, WISCONSIN.

No. 1278


Date

DONATED TO THE

Minnesota State Horticultural Society,

Case 34

No. 26



Apple Descriptions
by
John S. Harris
December 1901

Mill Seedling

Size 5

Form round

Color green to red
Striped and flushed

Stem long cavity

medium, green marked at bottom

Calyx closed basin broad rather

shallow corrugated

flesh yellowish nearly pure


core closed flavor subacid

Season early winter

Specimen grown by E. H. Sargent Antenna

has said to be a
seedling from transcendent

Crab
discovered Oct 20, 1899



Apple Descriptions
by
John S. Harris
December 1901

Bush Seedling No. 1, Lancaster Mass

Size 4 to 5 Weight 5 oz

Form oblate tapering a little

toward the eye very
slightly angular

Color orange yellow
ground in shade,
washed orange red
and striped & splashed
with brighter red in sun

Stem medium

Cavity broad open rather

deep, orange russeted calyx closed or partly

open basin broad medium shallow


ribbed or wrinkled, flesh light yellowish

tender juicy Flavor brisk Subacid

very good; Core medium partly open

Origin Lancaster Mass, Season Jan & later

Described Jan 27th 1899



Arch Seeding No 2 Lancaster Minn

Size 5

Weight 6oz

Form ^{smooth} round ovate

Slightly oblique

Color yellow with

blush on the sun

side with scattering

brown dots over the

surface Stem long

set in a narrow rather

deep thin rimmed cavity

calyx partly open basin narrow or small


medium, wrinkled; flesh yellow tender

nearly fine flavor rich Sweet should

think it would prove excellent for baking


Season January

Described Jan 2nd 1899



Gberhards Seedling No 2

Size 6 weight $7\frac{1}{2}$ oz
 form round inclining to
 cylindric or elongated
 Skin smooth greenish
 yellow thinly washed
 with brownish bluish on
 sun side and having
 numerous gray dots
 scattered over the surface
 Stalk long medium



deeply inserted in a regular
 narrow-deep russeted
 cavity. Calyx open in a
 rather broad medium deep slightly ribbed basin,
 core medium, closed seeds small, flesh
 yellowish fine grained firm; Flavor mild
 subacid. Should judge it will keep well into
 the summer. A chance seedling fruited for
 first time in 1898

Described Feb 4th 1899

Days Seedling No 1

Size 3 1/4

Form roundish flattened
at the ends

color yellow green
striped and splashed
with bright crimson
in narrow stripes


Stem long & slender

cupid medium calyx

closed basin medium broad irregular

wrinkled flesh white fine juicy pleasant

acid flavor Season late winter




Described Feb 21st 1899

Highland Seedling. (Winter)

Size 3

form roundish conical
color light green striped
and splashed with light
red on the sun side
Stalk long rather short
in a broad open and
deep cavity; calyx closed



basin medium, shallow slightly wrinkled; flesh
color light yellowish green or white, fine grained
firm flavor Sweet Season late winter
Oregon Fallmore co. Ann. Described Feb 4th 1899

Days Seedling No 2

Size 4 weight 4.03

Form round oblate

Color yellowish green

ground nearly over

spread with dark

very red or vermillion

with numerous fine gray

and brown dots in the skin

Stem medium long

carry rather broad (medium deep, orange

russeted Calyx closed, basin shallow

med broad wrinkled cor medium closed

flesh greenish white fine grained

Flavor mild acid fairly good

Season late Spring Is an annual

bearer



Described Feb 27th 1899

Scotts Wamter

Size 4 to 5 weight 4 to 5 oz
 form roundish conical
 of tin oblique and slightly
 angular color orange
 yellow-ground marked
 orange red striped
 and splashed crimson
 Stem Short and Stout
 in a deep narrow socket





Early Calyx closed deeply set in a narrow
 abrupt irregular wrinkled basin; flesh
 yellow, fine, juicy; flavor brisk rather
 acid Season Late winter

As a top worked tree it is an early and
 profuse bearer and has done well here
 for a number of years

Described March 4th 1899

From Wm Sandrock Bushford Spinn


Size 5. Weight 5 1/2 oz
 from roundish
 tapering a little towards
 the eye color a light
 yellowish white nearly
 over spread with light
 and dark red in stripes
 and splashes sometimes
 nearly solid brilliant red
 on the sun side, stem medium, to long
 medium, fleshy, set in an irregular medium
 sometimes broad cavity calyx closed in
 a medium nearly smooth basin. Corolla open.
 flesh fine grained white tender, very
 flavo-sulacid good
 slightly aromatic
 Season Jan to March
 It evidently belongs
 to the Gamnet
 Family and may be
 called Redly Brilliant



Described Nov 13th 1899

Mrs E. S. Nash No 1.


Size 5 weight 3 oz
 form oblate tapering
 slightly towards the
 edge. Color a dark
 yellow ground much
 overspread with
 red and striped
 and splashed with
 deep or dark crimson
 red and quite numerous
 grayish dots showing in the colors; Stem short
 medium in a broad med. deep gray russet
 -ed cavity frequently extending well out on the
 base of fruit Calyx closed basin medium
 broad uneven wrinkled flesh firm yellowish
 white flavor vinous subacid good
 Season best Feb & March but keeps well
 into the summer. The tree is found in
 a row of wealths and was procured for
 that variety and is said to be a vigorous
 healthy grower and an annual and abundant
 bearer. The location is favorable at
 Lanesboro Minn



Described March 13th 1899

Joseph Roemer seedling No 1 Reads chinn

Size 6 weight 60g
form flat conical
somewhat oblique
color, a ground of
light pale yellowish
green mostly over
sprued with a
stains of dull red
and stripes and
splashes of a deeper but
not bright red and many
small gray or russet dots that show most
on the sun side stem short medium
in a broad mid deep cavity dark russeted
at the bottom calyx small closed in
a medium open slightly wrinkled basin
flesh yellowish nearly white tender
juicy flavor mild subacid only fair
core small closed Season late winter
said to be a seedling of a supposed Baldwin
seedling but is more likely to be of
Jennet's Seed planted in 1886



The specimen described has commenced
to decay

Described March 16th 1899

Joseph Roemer's Seedling No 2 - Ready for
 "Pride" No 2

Size 4 Weight 403

form smooth round
 conical often a little
 oblique skin very
 light greenish yellow
 stained on the sun side
 with a dull bluish
 shade between stripes
 and splashes darker red



Surface of colored portions quite thickly
 sprinkled over with gray or russet dots most
 numerous toward the eye. Stem medium deeply
 inserted in a rather broad russet cavity
 calyx closed in a medium slightly wrinkled
 basin flesh yellowish nearly white tender
 juicy of mild subacid flavor good
 Season Jan - to March cost small
 Picked This is said to be a seedling of a
 Baldern Seedling Tree 12 years old and
 has borne fruit 8 years

Described March 16th 1899

Joseph Boerner Seedling No 3


Size 4 to 5
 form flatish irregular
 conical, oblong
 color yellowish
 green ground suffu-
 sed splashed and blot-
 ted more or less with
 brownish red. Stem



medium deeply set in a
 broad medium deep cavity, russeted well out-
 calyx closed basin medium abrupt irregular
 in form. flesh yellowish not as melting
 as No 2. Fine? Flavor mild subacid
 Season late winter at this date the the
 fruit is in good condition to keep much later
 The skin is rather thick and tough very
 small and close

Joseph Boemer Seed No 4

Size 4 weight 4oz
roundish oblate
conical color
light greenish
or yellowish green
Striped and splashed
in the sun with a
dullish black red



Stem medium in a regular broad
medium brown russet cavity

Calyx small closed in a smallish medium
narrow nearly smooth basin

flesh yellowish tender juicy

Mild sub-acid flavor core small
and nearly closed

Season January to March

Described Mar 13th 1899


Bendans Minnesota Gum

Weight 6 oz

Form roundish to
oval, convex
Color yellow ground
Striped splashed
and marbled with
red quite brilliant
on the sun side
Stem long in a



deep medium cavity
Calyx closed large abrupt
medium broad wrinkled
flesh yellowish
plant sub-acid
quality from fair to
good
Season late winter




Described Aug 20 1899

Wolf River

Size 8 to 10
Weight 9 to 14 g

Form roundish
often oblate
Color a yellowish
green greenish
covered with
dark red with
numerous gray or
russet dots over the





Surface Stem medium
in a medium golden russet cavity
cavity open in a medium basin
flesh yellowish rather coarse flavor
acid very cooking to market
Season usually early winter

The above specimen was grown by S. Bader
of Hobart has shriveled a little but is
free from decay at this date

Described Nov 20th 1899

Wm. Beustel seedling *Malus baccata* Mm
 Size 4 to 5 Height 50
 from roundish inclining
 to conical
 color at full maturity
 a greenish yellow
 ground spotted
 splashed and sometimes
 striped with red on
 the sun side stalk and
 long and slender in a rather deep
 greenish russeted cavity calyx closed
 in a medium ribbed or much wrinkled
 basin flesh yellowish, nearly white
 fine grained flavor an agreeable acid
 season late winter
 was apparently picked a
 little too early



Recd March 25th 1899

Histing

Seed planted about 1885 or 86 supposed to
be from Willow Spring

Only one turned out to be a good winter
apple - Transplanted once at 3 yr old
commenced to bear about 6 years since,
bears annually - fruit keeps well
Tree never blighted appears hardy nice
growing.

Size 4 to 5
form round oblate
color yellow green
striped red and
brilliant nearly
solid on the sun
side




Hamburg.

Size 3; Weight 5 to 6 oz

Form round or at
color nearly a solid
crimson red over a
whitish ground.

Stem medium, in
a narrow medium
cavity. Calyx small,
closed in a shallow nearly
smooth basin. Core medium



nearly closed; flesh light yellow fine grained
juicy. flavor pleasant acid or slightly sub acid
Season October. The tree is a symmetrical grower
and was not apparently injured by the
last winter and is reported to be a free
annual bearer. Grown sprout from the
root of a Fameuse tree that was killed to
the ground in the winter of 1884 & 5
in the orchard of S. B. Bellamy Homer Minn.

Described Oct 1st 1899
Specimen kept well until Dec 1st

Commons Seedling Crab

Size medium large

Form, roundish oblate

Color yellow ground

over spread with

crimson red lightly

striped Stem long

medium slender set

in a medium deep

greenish light russeted cavity


calyx closed, basin broad shallow and

considerably corrugated; core small closed

flesh yellow fine flavor agreeable acid

Season 2 September to October

Described Oct-15th 1899



J. S. Harris,


La Crescent,

Houston Co.

Minn.

Harsh Bennett's of Petersen

Size 3 to 4
form long oval
Color pale yellow
with faint bluish
cheek.



Stem long cavity
narrow - not deep
calyx closed, basin
thallum wrinkled flesh fine white tender
juicy flavor sweet. Season September
Oregon Russett does not agree with the
Russian description.

described Oct 3rd 1899

D. H. No 6

Size 3

form round conical

Color yellow

ground nearly

covered with red

in stripes and

thickly sprinkled over

with irregular grayish

specks or dots Stem long

and medium slender Cany medium


russeted at bottom Calyx large open

basin narrow deep ribbed flesh

yellow flavor sweet or slightly sub


acid Season September

Described Oct 3rd 1899



LeDus Seedling

Size 5-6
form roundish
slightly conical
color green or
yellowish and red
striped and splashed
stem medium, slender
cavity broad medium deep



brown russeted, calyx closed, basin med deep
bowl shaped flesh nearly white, often stained
with pale red juicy tender -
flavor agreeable acid or slightly subacid
Season winter

Described Oct 3rd 1899

Cummings Seedling

Size 6

Form Conical
tapering towards the
eye Color greenish
with stripes and flush
of brown red in the sun

Stem short to medium

in a broad medium deep russeted cany

calyx open basin medium deep wrinkled

flesh nearly white. Flavor brisk sub

acid Season Oct 1st

Described Oct 3rd 1899



Solanum Seedling

Size 4-5 Weight 4 to 5 oz

Form rounded
oblate slightly
angular

color light green

turning to pale
yellow when ripe

often with light
bluish cheek.



Stem long slender
set in a medium often irregular
green russetted cavity Calyx
closed in a rather large, ribbed
and wrinkled basin corolla rather
open flesh yellowish firm grained juicy
firm; flavor mild and good. Season
winter. See about 20 yrs old. 20 inches
in circumference. Specimen from Mrs. C. W.

Thayer Solanum Nunn

Observed Oct 20th 1899

Has kept well into February


J. S. Harris,

La Crescent,

Houston Co,

Minn.

Joseph Boerner Seedling No 3



Size.— 4-5 5
 Form.— Irregular flatish conical
 not on the oblique
 Color.— Yellowish green ground, striped
 splashed and blotched brownish red
 Cavity.— Broad and deep, russeted
 Stem.— Medium — Calyx small closed
 Basin.— Mid almost irregular
 Flesh.— Yellowish tender, fairly juicy
 Flavor.— Mild, sub-acid
 Core small closed
 Season.— Jan to March
 Free, Etc.—

Described.— March 16th 1898



Walter Seedling By J. R. Cummins

Size 5

Fruit round elongated oblique Color
light green turning to light yellow
Stem medium. Cavity medium, Calyx small
closed basin medium corrugated flesh
Whit, fine tender flavor mild acid
Season October
Described Oct 20th 1899.



J. S. Harris,
La Crescent,
Houston Co. Minn.

Walfert Seedling By Andrew Walfert
Cleveland Minn

Size 6 1/2

Weight 7oz

Fruit oval cone
slightly angular
Color yellowish
ground washed
with orange red
and streaked and
splashed with cream.
Stalk short medium
Cavity narrow rather
deep funnel shaped
Calyx closed



Basin Small medium angular or ribbed
Core open seeds roundish plump dark chestnut
Flesh yellow half fine tender
Flavor pleasant acid or subacid
Season winter Described Dec 11th 1900

Described Dec 11th 1900
This tree was produced from seed of Ben Jones planted in
1866 and is the only survivor of about 200 trees
Began to bear about 1874 Fruit has been kept in
ordinary cellar until the 4th of July

Lords L

Size 5

Form round or oblong
conic

Color a light yellowish
ground nearly covered
with deep dark red
with numerous small
gray or russet dots
showing through color

Stem long set in a
rather narrow deep
orange russet cavity
calyx small closed in a narrow
deep wrinkled basin


Flesh fine white stained red juicy tender

Flavor mild acid good

Season for the year

Oregon Minnesota City

Described Jan 12th 1909



J. L. Payne Seedling

Size 4

Form smooth, roundish

Color pale yellow

ground nearly
covered with bright

red in stripes and splashes

with numerous small gray dots on surface

Stem long & slender, set in a regular
medium deep cavity.

Calyx small, closed. In a smallish
medium nearly smooth basin.

Flesh white, sometimes stained with red, firm

tender, juicy. Flavor, subacid good;

Season Jan to Apr.



Described Apr 9th 1900

Smith H L

Sizes 3. from roundish
oblate color yellow

Stems medium canty

broad medium deep

calyx large closed

basin large broad shape


core open, flesh red

yellow, firm fine grained,

flavor pleasant acid Season

Dec or later evidently a hybrid

Dec 3 - 1895



Lightly so t

Size Med

Form roundish
color yellow

when fully ripe thin

Striped with light

red Stem medium

Slender in a rather

narrow medium early

Calyx closed in a rather broad corugated

basin cor small closed flesh yellow

fine grained firm flavor pleasant

rather sharp acid Not cooking

Season November

Described Dec 15th 1899



D. f. Texas Seedling Bushford Min

Size 4 weight $3\frac{1}{2}$ to 4 oz


Form smooth roundish
color a yellow ground
nearly covered with
crimson and orange
red in stripes and
splashes



Stalk medium set in a
regular medium broad cavity yellow russeted
at the bottom

Calyx half open; basin medium but slightly
wrinkled Core medium nearly closed

Flesh yellowish fine grain Flavor
mild Subacid. Season About middle of Aug



Described Aug 5th 1900

An apple recd from W. B. de Sur Hastings Oct 25th 99


Size 4 - Weight 4 oz

Form, smooth, roundish inclining to conical; color yellow-green nearly covered with deep crimson red faintly striped with numerous minute brown dots in the skin

Stem long and slender, in a medium broad and deep cavity, dull inserted at the bottom calyx small closed in in a mid broad and abrupt wrinkled basin, cor, small and closed stained scarlet flesh fine tender juicy flavor subacid slightly aromatic like an extra fine Wealthy season for

Described Oct 27th 99

J. S. Harris,
La Crescent,
Houston Co. - Minn.




Seedling from A Allert Deep Haren

Size 5 to 6 Weight 6 oz

Form smooth roundish tapering towards
the eye, color light greenish yellow
ground nearest covered with red
more or less striped and the skin thick
sprinkled with minute gray or whitish dots
stem long and slender set in a medium
regular rather deep cany light russeted at
the bottom, calyx small, closed in
a medium nearly smooth basin -
flesh yellowish ^{white} ~~pink~~ stained with red - nearly
fine juicy flavor subacid aromatic almost
identical with the Wealthy

Season September or later

Described Jan 2nd 1899.




Missouri

Size 5 Weight 5 oz
 form smooth round
 color greenish yellow
 ground marbled orange
 red striped and flushed
 with crimson

Stem medium set on
 a deep narrow cavity
 sometimes russeted at the
 bottom Calyx closed
 Basin medium slightly constricted
 core small closed

flesh yellowish fine grained firm
 flavor sub acid good season late winter
 The specimen was grown at Spring Valley
 Minn. by L. Wallen and was identified by
 the Pomologist at Washington as Missouri
 tree is said to be hard and good bearer



Winter Crab

Size, medium
 Form, round flattened at ends
 Color dark red with
 orange yellow speck, shiny through
 Stem long and slender set
 in a medium broad regular
 cavity
 Calyx closed in a broad
 conical basin
 Flesh deep yellow firm fine grained
 Core small, closed Skin subacid
 Season Jan to March
 Described Feb 17th 1901

