



  
**VICTORY**  
HEALTH & WELLNESS

# **THE 4-STEP SYSTEM TO END LOW VITAMIN D LEVELS**

[VictoryHealthWellness.com](http://VictoryHealthWellness.com)

# INTRODUCTION

We are highly disciplined about our health; we take our Vitamin D supplement faithfully, knowing it is crucial for a strong immune system, healthy bones, and stable energy. Yet, far too many of us still get disappointing blood test results, or worse, feel run down. If you've ever wondered why your vitamin D supplement isn't working, the problem isn't the product itself; it's a silent, internal conflict. Our bodies contain two hidden traps that actively seize and deactivate the Vitamin D we consume. The biggest obstacle is a biological reality: the vitamin gets locked up in our fat tissue, and even the amount that escapes is chemically useless without a key mineral.

This quick-start guide breaks down the science behind the "dilution effect" and the "activation block," giving you a clear, science-backed path to bypass both traps.

By focusing on release and conversion, you will stop wasting your supplement and finally experience the powerful health benefits Vitamin D is supposed to deliver.



Step 1.

---

# OPTIMIZE INTAKE & CHOOSE YOUR FORM

The journey starts with the right foundation.

- Choose D3 over D2:** Always ensure your supplement is **cholecalciferol (D3)**, as it is far more effective at raising blood levels than ergocalciferol (D2).
  
- General Dosing:** A common starting point for adults is 2,000–5,000 IU of D3 daily, but remember, this must be adjusted based on your body composition. **Disclaimer:** Consult your physician for personalised dosing based on your blood test results.
  
- The Sun Booster:** Aim for **10–20 minutes** of midday sun (arms and legs exposed) several times per week, if possible, balancing this with skin safety. This creates the most biologically available form of Vitamin D.



Step 2.

# **FIX THE ACTIVATION KEY (THE MAGNESIUM DEEP DIVE)**

Magnesium ensures the D you take (or release) actually gets used.

- The Problem:** Magnesium is needed to run the enzymes that convert D into its active form. Without it, the D gets stuck in a chemical traffic jam.
- The Food Fix:** Prioritize these Magnesium-rich foods daily: leafy greens, pumpkin seeds, almonds, and dark chocolate (cacao).
- The Supplement Solution:** Not all magnesium is equal. Use this guide when buying:
  - **Magnesium Glycinate:** Great for sleep, relaxation, and highly bioavailable.
  - **Magnesium Citrate:** Good bioavailability, often used to aid mild constipation.
  - **Magnesium Malate:** Excellent for energy and muscle soreness.
  - **AVOID Magnesium Oxide:** It has poor absorption (only 4%) and is mainly used for short-term bowel movements.



Step 3.

# ADDRESS THE STORAGE LOCK (THE D-RELEASE PROTOCOL)

This focuses on reversing the dilution and inflammation caused by excess adipose tissue.

- Reduce Inflammation:** Systemic inflammation hinders Vitamin D. Start by cutting out **highly processed vegetable oils** and **excess added sugar**. These simple changes reduce the inflammatory signals that interfere with Dactivation.
- Trigger Release with Movement:** Engage in regular physical activity. Movement, particularly **resistance training**, helps increase metabolic rate and triggers the gentle release of stored nutrients, including Vitamin D, from fat reserves.
- The Protein Priority:** Increase protein intake. Protein is thermogenic and helps preserve lean mass during fat loss, which is key to sustainably managing the storage reservoir of Vitamin D.



Step 4.

---

# SMARTER MONITORING AND RE-TESTING

Stop wasting time and money on inaccurate re-tests.

- Wait for Stability:** Wait a minimum of **90 days (3 months)** after starting a new D/Mg/Body Comp protocol before your follow-up blood test. This ensures the change in your supplementation and body chemistry has had enough time to stabilize the D's long half-life.
- Consistency is King:** Always request the test be done at the **exact same lab** that performed your original test. Even if the numbers look different between labs, using the same one ensures your comparison is valid and accurate.
- Optimal vs. Normal:** Discuss the results with your doctor, aiming for **optimal** levels (often 40–60 ng/mL) rather than just clearing the "normal" threshold.

You now have the knowledge to move beyond simply increasing your supplement dosage.

The frustrating cycle of low Vitamin D levels and persistent fatigue ends when we address the fundamental biological environment. Remember, the 2Hidden Traps—the Adipose Storage Vault and the Magnesium Block—are the real villains holding your health hostage.

The most effective strategy is a combination approach: we must strategically work to release the Vitamin D that is physically locked away by prioritising body composition, and we must simultaneously ensure that every dose is activated by consistently supplying the essential cofactor, magnesium.

That shifts the focus from supplement chasing to proper biological optimisation. By targeting both traps, you guarantee that every dose you take is thoroughly utilized by your body, giving Vitamin D the green light to support your energy, immunity, and overall well-being.