

Ultrasound Guided Minimally invasive (Sonex) Post Operative Instructions

Food

Eat what you want

Dressing

Sometimes we apply the dressing very tight after surgery. Please feel free to unwrap it and re-wrap to make yourself more comfortable after an hour or two. You may remove the dressing and apply bandaids or other dressing of choice in 48 hrs.

Keep the dressing/incision dry for 48 hrs. After that, it may get wet but do not soak it or get it dirty until it heals (usually 2-3 weeks)

Activity

Keep your hand elevated for 24 hrs to reduce swelling and pain.

You may do anything that your pain allows. Most people can return to light duties in 2 days. Many can return to heavy activity in 4 days but it can take as long as 4 weeks.

Blood clots

Blood clots are rare after upper extremity surgery.

Swelling

The best way to improve swelling is elevation of your hand. Ice also works if you do not have a cast or splint.

Physical therapy

You will not need PT. I do want you do stretch and move your hand as much as you can tolerate after surgery.

Medications

For the best pain control: Rotate over the counter ibuprofen and Tylenol. For example take regular Tylenol followed by Ibuprofen 2 hrs later. Do not take more than the recommended daily dose for each. Narcotics are not needed.

Narcotics

Narcotics are not needed after surgery.

Numbness

Numbness lasts a few hours after surgery. If your numbness does not improve after a few days please make an appointment for evaluation.

Post Operative Visits

Make an appointment 4 weeks after surgery if you want. Most people do not want to follow up.

Concerns

Call the office at if you experience the following: Increased redness at incision site Pain that is uncontrolled and increasing over hours Uncontrolled bleeding Fever > 101 o F or shaking chills Painful swelling

Contact

Please refer to the website for the office phone number.