



DARIN W ALLRED M.D.
ORTHOPEDIC SURGERY AND SPORTS MEDICINE

ACL Reconstruction Post Operative Instructions for BEAR PROCEDURE ONLY

(If you had MCL, PCL, or LCL reconstruction with your acl recon, these instructions are not for you)

Food

Start slow. It is natural to want to eat a lot after surgery. We recommend you eat small amounts of your normal diet and increase your intake as your nausea improves.

Incision

Keep your incision dry for 48 hrs. After 48 hrs you may remove the dressing and shower. Leave the Steri-Strips or glue in place. You may let water run over the incision but do not soak your incision in a bath, swimming pool, lake or hot tub for three weeks. Keep a Band-Aid on the incision until the incision is no longer draining (about 7 days)

Weight bearing

You must limit your weight bearing to 50% only for 4 weeks

Activity/Brace

Keep the brace locked in full extension for 24 hrs. After that follow the table in the attached BEAR post op protocol.

Blood clots

My biggest concern after acl reconstruction is the formation of blood clots. To prevent blood clots, I recommend you be active and take One tablet (325 mg) of aspirin twice daily with food for 2 weeks. If you or your family have had a blood clot or other reason to have an increased risk for a blood clot, you may need more aggressive treatment. Please discuss this with Dr. Allred within 24 hrs after surgery. What does a blood clot feel like? Usually (not always) blood clots surface a few weeks after surgery. They usually (not always) cause calf pain and increased ankle swelling that is new. Rarely, they cause shortness of breath. If you think you may have a blood clot call the office immediately (during office hours) or go to the ER (if after hours)

Swelling

Ice machines or reusable ice pads can be purchased on Amazon if you search knee ice machine. They can be great for decreasing pain and swelling. I like them but frozen peas work as well. Be careful to not freeze your skin. I recommend 20 min on and 10 min off unless the machine has a way to prevent freezing (ie temperature settings)

Physical therapy

You should have an appointment scheduled for physical therapy within a few days after surgery. If you do not, call the office to arrange.

Medications

As discussed above, I recommend Aspirin for two weeks. Rotate over the counter ibuprofen and Tylenol. For example take regular Tylenol followed by Ibuprofen 2 hrs later. Do not take more than the recommended daily dose for each. I also recommend over the counter Pepcid or other antacid while on ibuprofen to prevent a stomach ulcer. Narcotics: see below.

Narcotics

You may need them for a few days but the quicker you stop using them the better you will feel. It is very rare to need a refill. We **DO NOT** refill medications after hours or on weekends so plan ahead. It takes about 72 hrs to get a refill. If you take narcotic medications regularly you will need to make arrangements with your prescribing provider.

Post Operative Visits

Call for an appointment 2 weeks after surgery.

Precautionary Measures

- 1) To prevent infection, you have been administered IV antibiotics prior to your surgery.
- 2) To prevent blood clots, you will need to be active. I want you to walk around your living room once every hour while you are awake. Also take Aspirin 325 mg twice daily for 2 weeks to prevent a blood clot.
- 3) If you are taking narcotic pain medication (Percocet or Vicodin), you may need a stool softener to prevent constipation. Over-the-counter medication such as Docusate or Milk of Magnesia is recommended.

Emergencies

Call the office at if you experience the following:

Incision opens

Increased redness at incision site

Pain uncontrolled by pain medicine

Uncontrollable bleeding

Fever > 101 ° F or shaking chills

Difficulty breathing or chest pain

Severe pain or redness in calf

Painful swelling

Contact

Please refer to the website for the office phone number.

BEAR[®] Implant

Patient Post-Operative Instructions





Table of Contents

Page 1	Introduction
Page 2 - 3	After Surgery Instructions For Patients
Page 4 - 7	Home Exercises
Page 8 - 9	Frequently Asked Questions

Instructions for Surgical Team:

Please provide this document and the BEAR[®] Implant Rehabilitation Protocol to the patient and/or parent/guardian following surgery.

Date of surgery: _____

Name of patient: _____

Name of operating surgeon: _____

Contact for operating surgeon: _____

Introduction


Congratulations, you just received the BEAR® Implant. Below are the important things you need to know and do as you progress through the stages of ACL healing and rehabilitation.

What You Need to Know


1. **The BEAR Procedure is different than ACL reconstruction (ACLR).** The BEAR Implant and Procedure are designed to help your body restore your own ACL, not replace it.
2. Following the BEAR Procedure, your body will be working hard to replace the BEAR Implant with your own ACL tissue. During this time your new ACL is delicate, so attention should be given to protecting your injured knee.
3. The BEAR Implant Rehabilitation Protocol is carefully designed to meet important rehabilitation goals. To obtain the best result, please follow this protocol and any instructions from your surgeon.



Torn ACL



Surgeon adds
patients own blood
to the BEAR Implant



Surgeon inserts the
BEAR Implant between
torn ends of ACL



As ACL heals, the BEAR Implant
is absorbed by the body,
usually within eight weeks

What You Need to Do

1. **Ensure your physical therapist (PT) receives the BEAR Implant Rehabilitation Protocol.**
2. **Schedule your first physical therapy visit within the first week after your surgery.** At this visit your physical therapist will review and provide instruction for your Home Exercises.
3. To meet your range of motion (ROM) goals, **begin the Home Exercises within the first week after your surgery.**
4. **Carefully read and follow the instructions** in the pages that follow.
5. If you have any **questions or concerns, please contact your surgeon's office.**

After Surgery Instructions For Patients

There are clinical scenarios when your surgeon may alter your specific rehabilitation protocol, for example, in the case of a secondary procedure to the BEAR® Implant, such as a meniscal repair or additional ligament reconstruction or repair. In these cases, please follow the instructions you receive from your surgeon/surgical team.

Cryotherapy and Elevation

- Keep your operative leg elevated and apply cold compression for the first 24 hours or until post-operative swelling is controlled. Apply cold compression device once every waking hour for 15 minutes.
- After that period, apply the cold compression device 3 times per day for 15 minutes.
- Do not sleep with automated device running while on your knee.
- Keep a layer of fabric or ace wrap between skin and icing device at all times.

Crutch Use Instructions

- Beginning the day after surgery, you will be partial weight bearing (PWB) with crutches while walking and standing.
- Your goal is to be able to put half (50%) of your total body weight on your operative leg without pain.
 - *For example, if you weigh 150 lbs, your goal is to put 75 lbs of weight on your operative leg while standing and walking.*
- Under the direction of your physical therapist and operating surgeon, you may begin to wean off crutches at 4 weeks; some people may take longer. Follow the instructions provided by your PT to do this safely.

Knee Brace Instructions

- Keep your brace locked fully straight (zero degrees) for the first 24 hours after surgery or until 1st post-op surgeon visit for adolescents.
- Once your brace is unlocked after surgery, allow your knee to bend to the specified degrees listed in the chart below during non-weight bearing activities (sitting activities) such as watching TV, riding in car, sitting at computer, eating meals, and while performing your rehab exercises at home and with your PT.
 - Gaining and maintaining your knee's range of motion (ROM) is important for leg strength, gait, and stability.
- During this time also continue to work on getting your leg fully straight (see leg exercises below).
- Caution: your PT should not force your knee into a flexed position to protect the healing ACL.

After Surgery Instructions For Patients Cont.

Time Frame	Brace Range Setting	Instructions
First 24 hours	Keep brace locked at zero degrees	Keep your brace locked at zero degrees until 24 hours after your surgery or until 1st post-op surgeon visit
1 day after surgery to 2 weeks	Set your brace range 0 to 45 degrees	Set your brace so that it allows your knee to bend from fully straight to 45 degrees while non-weight bearing and performing rehab exercises Lock your brace at zero while walking for the first 4 weeks and while sleeping for 6 weeks
Weeks 2-4	Set your brace range 0 to 90 degrees	Set your brace so that it allows your knee to bend from fully straight to 90 degrees while non-weight bearing and performing rehab exercises Lock your brace at zero while walking for the first 4 weeks and while sleeping for 6 weeks
Weeks 4-6	PT will advise	Your PT and surgeon will evaluate your ROM and provide further instructions At 6 weeks, brace is no longer required for sleeping
After week 6	PT will advise	Dependent upon your progress, your PT may advise moving to a functional knee brace after week 6

Glossary of Terms

- **Flexion/Extension:** Leg bent/leg straight
- **Partial Weight Bearing (PWB):** Only putting a portion of your body weight on your operative leg (example: half your body weight (50%) on your surgical leg)
- **Range of Motion (ROM):** The measure of flexion and extension for a specific joint
- **Passive Range of Motion (PROM):** The manual movement by PT of a joint (example: your PT manually bends your leg as opposed to allowing your own muscles to bend your leg for you)
- **Active Range of Motion (AROM):** Use your own muscles to bend and extend your leg (example: you bend your leg to achieve a 60-degree bend)

Home Exercises - Knee Straightening Exercise

Begin Within the First Week After Surgery

Wearing your brace set to the specified degrees, perform the following exercise.

You can do this in one of the following two ways.

1. Sit in a chair and rest the foot of your operative leg on an adjacent chair. Then gradually let your leg relax so that you feel a gentle stretch on the back of your knee.
2. Put a small roll, pillow or rolled towel under your ankle so the back of your leg is lifted off the bed or floor. Then gradually let your leg relax so that you feel a gentle stretch on the back of your knee.

You should not push on the top of your knee or force your knee manually, just allow the weight of your leg to do the work.

Do this two times per day for 4 to 5 minutes at a time.



Home Exercises - Wall Slides

Begin Within the First Week After Surgery

Lie on your back with the foot of your injured leg on the wall. Let your foot slide down causing your knee to bend. Depending on your time out from surgery, use the chart above to flex your knee to the specific degree range (45, 90, or >90 degrees). Keep your knee in this position for 5 to 10 seconds. Do not flex your knee past the specified degree range in the chart.

Do 1 to 2 sets of 10 repetitions with a 5 to 10 second hold, at least 2 times per day.




Home Exercises - Quad Set / Isometric Contraction Exercise

Begin Within the First Week After Surgery

While lying flat on a bed or floor with knee straight, place heel on a rolled towel. Heel must be high enough so that your thigh and calf are off the bed or floor. Tighten your quadriceps muscles on the front of your thigh by pressing your knee down into the bed or floor. Hold this position (muscle contraction) for 10 seconds, then release. To properly perform this exercise, hold the above position for 10 seconds for 3 sets, do this 10 times.

Do 3 sets of 10 repetitions, 2 to 3 times per day.



Your quadriceps are a group of muscles on the front of the thigh located above your knee. They are responsible for extending your leg and helping with movements such as walking and jumping.

If comfortable doing so, you may remove your brace for this exercise only, then put it back on immediately after.

Home Exercises - Patellar Mobilizations

Begin Within the First Week After Surgery

Manually move patella side to side and up/down using your hands.

Repeat for 1-2 minutes 3 times per day.



Frequently Asked Questions

After BEAR Implant surgery, patients often have questions about when they can resume everyday life activities, such as driving or going up and down stairs. While exact timing is based on your individual rehabilitation progress, there are general guidelines for when you can expect to get back to these activities.

1. When can I travel on an airplane?

Timing: No restrictions

There are no restrictions on air travel after surgery. In fact, many patients have flown across the U.S. or from another country to have the BEAR Implant procedure, and then they have returned home shortly after surgery. However, the bracing and load bearing restrictions may make flying uncomfortable. Here are some considerations for traveling on an airplane in the first six weeks after surgery:

- **Sitting:** Depending on your time out from surgery, you can unlock your brace between 45 and 90 degrees, so you'll be able to put your leg under the seat in front of you. Keep in mind that it may not be comfortable to be confined to one position for an extended period of time.
- **Load Bearing:** You should avoid activities that require you to put more than 50% of your body weight on your knee, so you need to make sure that you are able to maneuver on your crutches and not carry heavy items, like luggage.
- **Standing:** Your leg needs to be straight with the brace on while standing and walking, so you'll need extra time to get through the airport.

2. When can I go up and down stairs?

Timing: Between 7-12 weeks

At 4 - 6 weeks, you'll start stair training as part of your rehabilitation. Your physical therapist will clear you to use the stairs once you can safely go up and down without significant pain or instability. Until then, it is recommended that you "scoot" up and down the stairs in a seated position. If you live in a two-story house, you may want to consider relocating your essential activities to the first floor.

3. When can I drive a car?

Timing: At least 2 weeks

If you received the BEAR Implant in your left leg, you can drive a car with an automatic transmission once you are no longer taking narcotic pain relievers and you can safely get into and out of the car. Depending on the size or height of your car, you may find it difficult to get yourself in and situated with your knee brace locked at 45 degrees.

If you received the BEAR Implant in your right leg, you may be able to drive a car with an automatic transmission around 2 weeks. That's when your knee brace is unlocked to 90 degrees. Not everyone feels strong enough at 2 weeks to drive, so it's best to discuss your individual progress with your physical therapist.

Frequently Asked Questions

4. When can I play sports?

Timing: Between 9-12 months

You can return to sports once you meet all the return-to-sport criteria – typically around 9 months. Your physical therapist will conduct return-to-sport testing to ensure your knee is strong and able to perform the movements needed for your sport. Key criteria include that you feel confident when running, cutting and jumping at full speed, you can hop on your BEAR Implant knee at nearly the same level as your other knee and you are confident in the function of your knee.

5. When can I run?

Timing: After 18 weeks

Straight line running as part of physical therapy on a treadmill or protected environment can begin after 18 weeks if you've met your rehabilitation goals to that point. Running outside of physical therapy is typically on the same timeline as returning to sport.

6. When can I carry my baby?

Timing: At least 4 weeks

While you're using crutches, it is recommended that you hold your baby only while sitting. This is for the safety of the baby and so that you don't exceed the 50% load bearing restriction. You should be able to graduate off crutches starting at 4 weeks, and shortly after that you can carry heavier objects if you feel ready.

Who should you contact if you have questions?

It is common to have questions about your recovery and how your recovery is progressing. Each patient's BEAR Implant procedure and recovery are unique, so contact your surgical team with any questions. If they need further guidance, they will reach out to Miach Orthopaedics.