

DESERT KNOWLEDGE CRC

High Conservation Value in the Rangelands

Mark Stafford Smith
Andrew Ash

Report
19

December
2006

A report prepared for the
Australian Government Department of the
Environment and Heritage
by the Desert Knowledge
Cooperative Research Centre



High Conservation Value in the Rangelands

Mark Stafford Smith

Andrew Ash

Workshop, Adelaide, 22–23 November 2005

A report prepared for the Australian Government
Department of the Environment and Heritage by the
Desert Knowledge Cooperative Research Centre

Contributing author information:

Mark Stafford Smith is a research fellow with Desert Knowledge CRC and long-term arid zone researcher with CSIRO Sustainable Ecosystems.

Andrew Ash is Deputy Chief of CSIRO Sustainable Ecosystems with lengthy research experience in rangelands ecology.

Desert Knowledge CRC Report Number 19

Information contained in this publication may be copied or reproduced for study, research, information or educational purposes, subject to inclusion of an acknowledgement of the source.

ISBN: 1 74158 030 7 (Print copy)

ISBN: 1 74158 031 5 (Online copy)

ISSN: 1832 6684

Citation

Stafford Smith M, and Ash A 2006, *High Conservation Value in the Rangelands*, Report of a workshop for the Australian Government Department of the Environment and Heritage, Desert Knowledge Cooperative Research Centre, Alice Springs.

For additional information please contact:
Department of the Environment & Heritage
Mr John Lumb
GPO Box 787
CANBERRA ACT 2601
Phone: 02 6274 1036
Email: john.lumb@deh.gov.au

© Commonwealth of Australia 2006

This work is copyright. Apart from any use as permitted under the Copyright Act 1968, no part may be reproduced by any process without prior written permission from the Commonwealth, available from the Department of the Environment and Heritage. Requests and inquiries concerning reproduction and rights should be addressed to:

Assistant Secretary
Natural Resource Policy Branch
Department of the Environment and Heritage
GPO Box 787
Canberra ACT 2601

Disclaimer:

The views and opinions expressed in this publication are those of the authors and do not necessarily reflect those of the Australian Government or the Minister for the Environment and Heritage.

While reasonable efforts have been made to ensure that the contents of this publication are factually correct, the Commonwealth does not accept responsibility for the accuracy or completeness of the contents, and shall not be liable for any loss or damage that may be occasioned directly or indirectly through the use of, or reliance on, the contents of this publication.

Project report

for the Department of the Environment and Heritage

This document provides a short Final Report for Desert Knowledge CRC Project *Criteria for Defining Areas of High Conservation Value in the Rangelands*, a small project contracted by the Department of the Environment and Heritage (DEH) in late 2005 in the form of a workshop (see Appendix). The objective of the workshop was to draw on key ecological, management and policy expertise to define a state-of-the-art rationale for the mapping of high conservation value (HCV) areas in the rangelands, and, if necessary, to develop the approach required to test and implement this mapping.

Workshop rationale

For a number of years there have been concerns in various agencies that approaches to identifying areas of high conservation value used for other regions of Australia may not provide the best outcomes in the rangelands. In response to these concerns, the Australian Government DEH has sought to explore alternative approaches and this workshop was intended to bring together the thinking and expertise around the subject. Amongst other matters, a better basis for defining areas of high conservation value will assist priority-setting for public investment at a variety of scales, including both traditional activities such as reserve acquisition, as well as more contemporary issues such as regional plan accreditation, incentives and off-reserve conservation actions.

Key findings

The key points from the workshop were:

1. We conclude that a process for determining HCV areas cannot currently be operationalised fully in the same way in rangelands as in fragmented landscapes, because of some key distinguishing features of rangelands in practice (Box 1).
2. There are two classes of HCV areas in rangelands – (i) those that can be conserved by action in a specific locality or focal area, such as a national park or other off-reserve area, and (ii) those that are supported by maintaining certain diffuse ecosystem processes across large areas – such as genetic flow across landscapes, the integrity of water redistribution (potentially at local scale but with regional implications) and the spatio-temporal dynamics of disturbance regimes at large scales – which drive the maintenance of genetic diversity (α , β and γ) in semi-natural landscapes (see Box 2).
3. The same two classes probably apply in all land types, but in fragmented landscapes the importance of focal areas for conservation greatly outweighs the importance of diffuse processes (of course, diffuse processes remain important for production in fragmented landscapes). The significance of the two classes of HCV areas is relatively more even in rangelands.

4. We conclude that priority-setting for the first class – focal areas – can be handled in rangelands (as elsewhere) by the current tools of Systematic Conservation Planning (SCP) as applied under a conventional reserve design paradigm. However, the application of this approach in rangelands using tools such as Marxan or C-Plus may be constrained by issues to do with data types, data resolution and data availability, as identified at the workshop (Table 1).
5. By contrast, the second class – diffuse processes (Table 2) – requires an approach which is paradigmatically different from that of conventional reserve design techniques. We began outlining the features of this approach which, in practice in Australia, would be based around IBRAs¹ (or preferably sub-IBRAs²). It would take the form of (see Fig. 1):
 - (a) identifying any differential intrinsic values represented by each IBRA in a pristine condition
 - (b) identifying any differential values represented by each IBRA in their current condition
 - (c) identifying what management interventions might be appropriate to maintaining the diffuse processes in that IBRA (and their costs)
 - (d) identifying the degree of benefit to be achieved by making those management interventions.
6. Whilst we thought there are strong conceptual reasons to imagine that the intrinsic pristine differences between IBRAs exist (ie. step a)³, there is limited evidence for this at present⁴, so we suggest initially valuing all IBRAs equally in terms of their pristine intrinsic value. However, this assumption should be critically re-appraised in the future.
7. The current condition values of IBRAs (step b) would be assessed in terms of historical damage, both in terms of degree and, as importantly, longevity. Longevity of impact is important: although some processes that create local genetic diversity (Box 2) might be re-instated quite easily, if they have been absent for long enough, the diversity has been lost (as in many fragmented landscapes) and will not be re-instated until genetic drift is re-established, probably over centuries. In a general sense this type of description of many IBRAs has already been carried out by jurisdictional and other projects (e.g. see <http://www.deh.gov.au/land/management/rangelands/processes.html> and <http://www.nrm.gov.au/monitoring/indicators/vegetation-condition/index.html>) but not with a specific eye to these diffuse processes which support biodiversity.
8. The costs and benefits of conservation action (steps c and d) in terms of invasive species control, maintenance of water-remote areas, maintenance of landscape function in terms of water flows, re-establishment of wetland networks, etc, have also been summarised for many IBRAs, but again without a focus on diffuse processes – in general these have been discussed in terms of maintaining processes at the far less critical level required for production.

1. Interim Biogeographic Regions of Australia as originally proposed by Thackway and Cresswell (1995), and subsequently extensively revised as shown recently (version 5.1) in NLWRA (2002).

2. In practice any large regional unit could legitimately be used, but this report uses IBRAs as the pragmatic most likely approach, and as shorthand for any other regionalisation. The key issue is that the units must be large enough to encompass regional ($\sim 10^3$ – 10^6 km²) scale diffuse processes.

3. Because each of the examples of diffuse processes in Box 2 could be expected to be more significant in some IBRAs than others.

4. Whilst we acknowledge that conservation management is generally about maintaining or changing the current state, there are circumstances where the aspiration to restore or conserve a ‘pristine state’ may be a viable option, so it is important to include this consideration. The unconstrained flooding systems of some parts of the Lake Eyre Basin might be an example.

9. It is worth noting that the reality of thresholds and change in critical slow processes means that there may often be a good case for intervening in an IBRA which is no longer in particularly good condition before some additional threshold is crossed. Nonetheless, it is likely that, as the condition of a rangeland IBRA deteriorates and, in particular, becomes more fragmented, it will converge on the situation for fragmented agricultural landscapes where the conservation of focal areas becomes a higher priority.
10. Although the approach to prioritising for diffuse processes has some parallels to current reserve design methodologies, it is conceptualised at a far larger scale of geographic unit (not generally amenable to excision), and the issues of representativeness, geographic relationships and perhaps irreplaceability are not so meaningful. Complementarity remains important, as may a fourth basic element (perhaps 'synergy' or 'additionality'?) – the degree to which the protection of focal areas in a reserve network add up to reduce the costs of maintaining sufficient functioning processes across the remainder of an IBRA. Development of data or indicators to represent these diffuse processes requires a different conceptual approach to the way data is used for reserve design planning. Hence using the standard software such as Marxan and C-Plus is probably not the appropriate methodology.⁵ It is likely that some form of Multi-Criteria Analysis (MCA) would handle the diverse values and weightings adequately; the use of these types of approach to incorporate more ecological processes in conservation planning is beginning to be explored in the SCP community.⁶
11. We spent inadequate time debating the question of how one might judge the relative balance of resources to put into the two classes of HCV areas in rangelands.

Future workplan options

This analysis leads to a number of future steps for DEH to consider:

- (i) Commission a systematic review of the IBRAs with a view to establishing the datasets for steps (b–d) above in Key Point 5, at least in some qualitative form. This would review the current descriptions of the IBRAs with a specific focus on diffuse processes for maintaining genetic diversity values (specifically within- and between-species diversity at a 10^{2-5} km² type of scale) as opposed to production values (which might require some further debate on the precise list of these); the study would then send the summaries out to regional experts to complete the review.

Potential cost (assume a coordinating contractor @50k plus collaboration from each jurisdiction @40k each, using an ACRIS-type model): \$250k

- (ii) The current efforts to apply SCP to focal areas prioritisation nationally, including rangelands, should proceed, but take account of the data issues raised for rangelands, and potentially draw further rangelands skills into the project steering team.

Potential cost: drawing the expertise in should require negligible extra cost; however, there may be significant data management costs not included in the current project which cannot be estimated here.

5. Note: the current reserve design tools such as Marxan can be used *within* a region to determine where to support management with incentives, etc (eg. the maintenance of water-remote areas, cf. James et al. 1999, Biogroze 2000, Landsberg et al. 2002); however this should not be confused with establishing the priority of how much to invest in doing that action in this particular IBRA as opposed to other actions and in other IBRAs.

6. New developments such as MarZone from UQ take up this challenge.

- (iii) In the light of (i), an initial attempt at an actual prioritisation process would be commissioned.

Potential cost (assume a small project team developing and applying an MCA-type approach over 9 months, to data that is assumed already collated under (i)): \$110k

- (iv) Some further discussion may be needed as to how to balance the effort put into focal areas as opposed to diffuse processes from the conservation point of view in rangelands.

Potential cost: this could be added on to work in the existing project or under (iii), or could be the subject of perhaps 2 teleconferences with appropriate participants for negligible cost (\$2k if carried out formally?)

- (v) As a more theoretical question, a review of whether there is any literature which could justify differential values for IBRAs in pristine states (step a in Key Point 5 above) is needed. It seems to us likely that regions with different degrees of spatio-temporal variability might logically have different degrees of landscape-based genetic diversity; and there may be criteria by which to judge what regions play the most important roles in connecting this diversity at large scales. Elucidating ideas about this requires an initial review but could also lead to important basic research eventually.

Potential cost (for review only, assuming a post graduate or equivalent undertook a 4 week literature review assisted by an expert steering group through 2 teleconferences and email review, and one bout of consultative air travel): \$15k

[further resulting research if required could be promoted through an ARC Linkage Grant or equivalent]

Further details of the discussion during the workshop are provided informally in the unpublished Workshop Record, available from the authors on request, and a more detailed rationale for the findings is being drafted in a separate short journal paper.

Box 1: What are the unique characteristics of rangelands?

It is important to assess whether approaches typically used in systematic conservation planning for the intensive-use zone of Australia are equally applicable to the rangelands. We argue that rangelands are characterised by a unique set of attributes, due both to inherent ecological characteristics, and to pragmatic factors such as current land use regimes. In practice, the availability and adequacy of data useful for conservation planning is also an issue, as in other data-scarce areas.

1. *The rangelands comprise very extensive areas of natural or semi-natural vegetation.* In contrast to much of the mesic zone, rangelands remain relatively intact, which provides opportunities to undertake conservation planning across whole landscapes, and to make a genuine attempt to conserve genetic diversity (both in species with very broad distributions and those with high levels of within-species genetic variability across smaller ranges and within habitats) (Morton et al. 1995).
2. *High variability in space and time is a key feature of many ecological and landscape processes in rangelands.* In particular, the major drivers of rainfall and disturbance (notably fire and grazing) are highly variable both spatially and temporally. This variability is evident at a variety of spatial (e.g. hillslope, sub-catchment, catchment and regional scales for movement of water, nutrients and sediments) and temporal (within-year, between years and decadal or century-scale cycles) scales and may be highly unpredictable (Stafford Smith and Morton 1990). Consequently, patterning of biota is dynamic and even basic attributes such as local species richness are difficult to define or measure.

Two other features of rangelands are related to this high level of spatial and temporal variability:

3. *Landscape-scale processes are as significant (for conservation planning) as local richness or similar biodiversity attributes.* Concentration of conservation planning on localised areas can only ever protect some proportion of the biota. The maintenance of broadscale, diffuse ecosystem processes is necessary for conservation of many components. As rangelands comprise extensive areas of (semi-)natural vegetation this may still be achieved (Morton et al. 1995). Such processes either never operated at broad scales in the more productive mesic country or, following fragmentation, are no longer as important to conservation planning which concentrates on identifying and conserving remnants. For examples of diffuse processes, see Box 2.
4. *Some organisms operate over large to very large spatial scales.* Mobile organisms track variable resources or preferred habitat over space and time (eg. waterbirds moving between ephemeral wetlands (Roshier et al. 2001a, Roshier et al. 2001b); honeyeaters following intermittent flowering of nectarivorous plants). Even in some sedentary organisms, the persistence of regional populations may depend on the maintenance of habitat heterogeneity over very large areas (eg. species dependent on juxtaposition of particular seral stages after fire).
5. *Highest diversity of many organisms occurs in less fertile lands.* While restricted patches with relatively high productivity may be important for certain taxa, such as mammals and birds, many groups, such as lizards, termites, ants and gryllacridids, reach their greatest diversity in infertile habitats (Morton and James 1988, Stafford Smith and Morton 1990, James and Shine 2000). It is therefore unlikely that these taxa can be captured within a relatively small selection of biodiversity 'hotspots', nor is it likely to be possible to identify areas of high diversity through (easily mapped) surrogates such as primary productivity or vegetation types (even if it were possible to map them at high resolution, which is in fact unrealistic). This is further exacerbated by species complexes in

which high levels of unspecialised genetic diversity (eg. Kearney 2003, Ladiges et al. 2006) may be masked by lumping into superficially single species records.

6. *Environmental and biodiversity data are generally sparse or have coarse spatial resolution for most rangeland regions.* Moreover, environmental and biotic variation is often subtle and patterns are difficult to resolve or map at finer scales. Over much of the Australian rangelands, environmental mapping is available at scales of 1:250000 or poorer. Moreover, state- or continental-scale vegetation mapping typically shows rangelands to have relatively few very extensive map units. Such mapping does not adequately represent the often subtle variation in many rangeland environments and the well-known biotic patterning associated with them (for example, most rangelands are very flat, but slopes of a fraction of a degree are significant for landscape processes) (eg. Pickup 1985, Tongway and Ludwig 1994). Consequently, such mapping does not allow adequate assessment of representativeness, complementarity or irreplaceability in conventional conservation planning processes.

While high-quality data relating to species distribution is available for a few taxa and some localised areas, such point data is generally sparse and geographically highly biased. Some of this data also has to be interpreted cautiously – for example, records may show irruptive species to be widespread but their non-irruptive distribution may be very limited but most important to conservation planning.

Poor availability of data in the rangelands has led to the search for appropriate indicators of biodiversity at plot and landscape scales and to the use of remotely sensed vegetation and soil attributes as surrogates for habitat condition and biodiversity (Smyth and James 2004). The challenge with this approach is to identify indicators that will adequately represent biodiversity trends.

7. A related issue is that understanding the *spatial distribution of the condition of vegetation units may be particularly important* to conservation planning in rangelands. Again, differences in condition may be subtle and/or difficult to determine, and available spatial data is particularly poor. This issue is important in fragmented landscapes too, but is usually overwhelmed by the dramatic differences between land-uses, such as cropping and remnants (in whatever condition). By contrast in rangelands it is the major axis of variation within the extensive land uses (Morton et al. 1995).

Box 2: Diffuse processes in rangelands

Examples of diffuse processes include:

- maintenance of pollination processes – clearly, these are important everywhere
- supporting recruitment and migration, including management of the matrix between refugia, etc – these processes are important everywhere but they occur at large scales [eg. with waterbirds (Roshier et al. 2001a, Roshier et al. 2001b), rock wallabies (Pearson 1992)] in rangelands, and are more likely to be still occurring in the general landscape here than in fragmented landscapes
- ensuring there is continued opportunity for the intersection of multiple patchy processes (fire, grazing, rain, etc) at large scales – this will require maintaining diversity of grazing and fire regimes at landscape and regional scales to maintain species diversity, but more importantly within species genetic diversity. Low species diversity associated with species complexes and reticulate evolution in (eg.) *Senna artemisioides* and various *Acacia* sp may in fact be masking regions with high genetic diversity (Stafford Smith et al. 1996, Kearney 2003 and references therein);
- large scale landscape function processes such as sustained flows in major river systems (eg. Sheldon et al. 2002).

Subsequent discussions⁷ have suggested that these might be loosely categorised into the following groups, with examples and possible approaches:

Functional categories of processes	Example reasons (should be exhaustive; needs ecological justification!)	Possible surrogates (that need ecological justification!)	Possible selection strategy
1. Large scale quasi-universal/ spatially-continuous processes	1a. Undetected genetic diversity (eg places with substantial reticulate evolution, etc)	• Degree of variance in spatio-temporal disturbance regimes (fire, rainfall, etc) • Variance in GPP	Select reasonable geographic spread but also zone at regional scale for lower level investment in maintenance of landscape function
	1b. Maintenance of dispersal (including directional dispersal, eg. as needed in response to climate change)	• Connectivity of regions • Preferential orientation (eg. with respect to climate gradients) of connectivity if directional	
2. Processes focused on small areas but spread across large (~10 ²⁻³ km) spatial scales	2a. Focused on fixed localities, but with linkages among sites crucial for survival (eg. fish in permanent waterholes)	• Migration routes • Spatially-fixed high production spots in GPP • Permanent waterholes	Maintenance of connected network (maybe select geographically spread sets, but also zone regions with lower investment maintenance of the matrix)
	2b. Focused on varying localities of which some need to be linked accessibly at any time (eg. fire patch edges for organisms needing both recent and old patches – but larger-scale examples??)	• Regions with high incidence of high production spots in GPP at any time (but not fixed in space)	

7. Notes by Mark Stafford Smith after Spatial Prioritisation workshop, 9th Nov 2006

Table 1: Attributes of focal places and implications for SCP and data use.

This table summarises the types of attributes that would contribute to defining high conservation value for focal places, and the issues that might need special consideration in finding and using surrogates for these in rangelands (compared to other regions – ie. generic problems are generally not noted). [From the workshop.]

Attributes	What's important in Rangelands	Data layers/Indicators	Implications for planning and data in rangelands
COMPOSITION			
Rare species	Sparse data; definitional issues (irruptives, etc)	Species distributions	Should distinguish genuinely rare species from species that irrupt that may appear as common but are rare most of the time – consider treating refugia in non-irruptive phases as key focal points
Threatened species	Nil	Species distributions	Distinguish species that are localised and threatened from those that are widespread but threatened (localised ones have higher priority for focal conservation action; the latter require diffuse process management and less emphasis on specific point records)
Species richness	Richness in different taxa to non-rangelands often	Species distributions	Subdivide to major taxa, so that the comparative dominance of different taxa in rangelands doesn't distort outcomes (eg. reptiles, ants, etc)
Endemic species	Dispersed and patchy species under-represented	Species distributions – Number of species in spatial selection cells	Scale dependence of issue interacts with poorer spatial data resolution in rangelands (sensitivity analysis for this?) [Possibly less attention to appropriate levels of species classification in rangelands]
Species of restricted distribution	Some species are in restricted patches but very widely spread (eg. <i>Xanthorrhoea thorntonii</i>)		Occurs elsewhere but here over geographic scale that is larger than would be considered in non-rangelands environments
Habitats of restricted distribution	Larger spatial scales and habitats with little connection except during extreme events (eg wet periods)	Vegetation, landsystem, soil types, geological types wetlands	[eg. mound springs, mountain gorges/seepages, etc] Possible issue of occurring at smaller scale than the coarser vegetation mapping that is common in rangelands
Condition and intactness in relation to species complement	Nil	Productivity, remoteness from watering points and stocking rates	Important criterion for choosing between alternatives within one vegetation type in rangelands. Differences likely to be more in understorey state rather than woody structure. Could use cover (ACRIS) dataset; also water remoteness as key indicator. Ideally should recognise areas that have avoided grazing by chance but this generally needs local knowledge.
STRUCTURE			
Condition and intactness in relation to age structure and recruitment	Mapping not resolved enough (i.e. subtlety of gradients)	Productivity, remoteness from watering points and stocking rates	As above. Plus fire and rabbits important in driving this in rangelands woody vegetation. Some woody thickening may be detectable in AGO change data; still a question of whether this makes any difference to conservation value. Clearing is an issue in rangelands margins.
Complexity of assemblages and temporal variation in this	High temporal variability	Lack of knowledge of assemblages	See above

Attributes	What's important in Rangelands	Data layers/Indicators	Implications for planning and data in rangelands
FUNCTION			
Richness of habitat types, including local landscape diversity	More subtle variation and low resolution mapping	Vegetation, landsystem, soil types, geological types, geomorphic types	General absence of fine scale mapping – could use remote sensing to provide some indicator of spectral diversity (more sophisticated semi-variograms/local spatial autocorrelation measures?)
Specific ecological processes	e.g. wetlands for bird breeding; mound springs	Directory of important wetlands and RAMSAR wetlands and Dave Roshier layers	Breeding areas, etc. not generally apparent from available data layers. Include inventories of wetlands where possible.
Condition and intactness in relation to functioning processes		Productivity, remoteness from watering points and stocking rates Hydro-ecological processes, topographic diversity	See diffuse processes, but also indicators from previous condition entries above
Spatial unit area (km sq)	Rangelands of lower productivity per unit area	Use built-in aggregation rules in Marxan	Consider whether appropriate sizes need to be different in rangelands due to scale of ecological processes (and management efficiency)


- Priorities at holistic level: a fundamental consideration of SCP is complementarity (the selection of sets of priority areas that maximise biodiversity when assessed collectively, vs. hotspots which are areas of importance individually), for which there may be implications of environmental mapping being relatively coarser in rangelands; aggregation rules might need to favour larger areas where possible.
- Threat maps are not included here – weeds and ferals (both fairly coarse and often inconsistent across boundaries; some point data).
- There is a particular opportunity in rangelands to draw on better quality state-level databases rather than poorer national ones.

Table 2: Links between specific management interventions and the ultimate biodiversity conservation goals of maintaining diffuse processes at a regional scale. [From the workshop.]

Threats to diffuse processes	Specific regional management interventions	Immediate (observable, measurable) diffuse processes being sustained	Ultimate (hard-to-measure) biodiversity assets being sustained
<i>Intensification (fragmentation from fencing, roads; water points; clearing)</i> <i>Total Grazing Pressure</i> <i>Ecologically Significant Invasive Plants</i> <i>Altered Fire Regimes</i> <i>Water Diversion, Dams etc.</i>	<i>Fire management</i> <i>Grazing distributions and water point development intensity</i> <i>Regionally organised protection of water-remote areas</i> <i>Total grazing pressure</i> <i>Control of weed and feral animals</i>	<i>Fire regime diversity</i> <i>Grazing regime diversity</i> <i>Landscape connectivity at all scales</i> <i>Water flows at multiple scales</i> <i>Soil and microhabitat functioning with emergent regional implications</i> <i>Predation pressures</i>	<i>Local genetic diversity where dependent on cross-fertilisation or re-invasion (eg. rock wallabies), or on widespread local landscape function (eg. mulga groving, terminal lakes on large river systems), or on the diversity of landscape ephemeral patch outcomes which are not fixed in space or time (eg. intersection of burned patches with rainfall, chats)</i> <i>Local genetic diversity where this is dependent on continuous functioning landscapes</i>

Measures of current status of IBRAs include: Land Condition; History of intensification (Indicators: property size, water remoteness, distance to markets, time since settlement, ground cover, ACRIS collation [poor info on Indigenous lands]); Species status; Fire histories; Ecologically significant invasive species and ferals; Long-term climate history; Water diversion; Road/track network intensity?

Figure 1: Basic conceptualisation of the procedure for prioritising conservation actions across regions, aimed at diffuse processes and their influence on maintaining biodiversity assets. [From the workshop.]



Appendix: Workshop participants and agenda

Person	Affiliation	Email
Annemarie Watt	Department of Environment and Heritage, Canberra	Annemarie.Watt@deh.gov.au
John Morley	Department of Environment and Heritage, Canberra	John.Morley@deh.gov.au
Andrew Ash	CSIRO Sustainable Ecosystems, Brisbane	Andrew.Ash@csiro.au
Mark Stafford Smith	Desert Knowledge CRC, Alice Springs	Mark.StaffordSmith@csiro.au
Craig James	Desert Knowledge CRC and CSIRO Sustainable Ecosystems, Alice Springs	Craig.James@csiro.au
Josie Carwardine	Ecology Centre, The University of Qld, Brisbane	j.carwardine@uq.edu.au
Alaric Fisher	Department of Natural Resources, Environments and the Arts, Darwin	Alaric.Fisher@nt.gov.au
Rob Brandle	SA Department for Environment and Heritage, Adelaide	brandle.robert@saugov.sa.gov.au
Mike Fleming	NSW Department of Environment and Conservation, Dubbo	Mike.Fleming@environment.nsw.gov.au

(A Western Australian invitee was ultimately unable to attend)

Workshop Agenda: Criteria for Defining Areas of High Conservation Value in the Rangelands	
Tuesday 22 November 2005, Adelaide	
9.00 am	Introductions and Background – Mark Stafford Smith [MSS]
9.30 am	Discuss and clarify purpose of workshop and policy drivers for the question – ‘value’ for what, scale, intended uses of outputs, etc – Annemarie Watt, Andrew Ash
11.00	Identify alternative approaches to thinking about biodiversity value in the rangelands, and the consequent potential drivers – inputs from all [MSS]
1.30–5.00 pm	Define scientific and conceptual approaches to biodiversity value drivers, with ideal proximate causes – discussion groups
Wednesday 23 November 2005	
9.00–1230	Reflect on progress on the previous day [MSS/JM?] Review data and consequent options and limits to implementation – what data is there about or indicators of key proximate causes
1.30–3.00 pm	What would be necessary to test the proposals, and how could they be implemented in practice? – implementation plan(s)
3.15–4.00 pm	Wrap up and depart

References

- Biograzed. 2000. *Biograzed: Waterpoints and Wildlife*. CSIRO, Alice Springs.
- James, C. D., J. Landsberg, and S. R. Morton. 1999. Provision of watering points in the Australian arid zone: a review of effects on biota. *Journal of Arid Environments* **41**:97–121.
- James, C. D., and R. Shine. 2000. Why are there so many coexisting species of lizards in Australian deserts? *Oecologia* **125**:0127–0141.
- Kearney, M. R. 2003. Why is sex so unpopular in the Australian desert? *Trends in Ecology and Evolution* **18**:605–607.
- Ladiges, P. Y., S. R. Ariati, and D. J. Murphy. 2006. Biogeography of the *Acacia victoriae*, *pyrifolia* and *murrayana* species groups in arid Australia. *Journal of Arid Environments* **66**:462–476.
- Landsberg, J., C. D. James, J. Maconochie, A. O. Nicholls, J. Stol, and R. Tynan. 2002. Scale-related effects of grazing on native plant communities in an arid rangeland region of South Australia. *Journal of Applied Ecology* **39**:427–444.
- Morton, S. R., and C. D. James. 1988. The diversity and abundance of lizards in arid Australia: a new hypothesis. *American Naturalist* **132**:237–256.
- Morton, S. R., D. M. Stafford Smith, M. H. Friedel, G. F. Griffin, and G. Pickup. 1995. The stewardship of arid Australia: ecology and landscape management. *Journal of Environmental Management* **43**:195–217.
- NLWRA 2002. *National Land & Water Resources Audit. Australian Terrestrial Biodiversity Assessment 2002*. Canberra: Commonwealth of Australia.
- Pearson, D. J. 1992. Past and present distribution and abundance of the black-footed rock-wallaby in the Warburton region of Western Australia. *Wildlife Research* **19**: 605–622.
- Pickup, G. 1985. The erosion cell – a geomorphic approach to landscape classification in range assessment. *Australian Rangeland Journal* **7**:114–121.
- Roshier, D., A. Robertson, and R. Kingsford. 2001a. *The availability of wetland habitat for waterbirds in arid Australia*. Environment Australia and Land and Water Resources Research and Development Corporation, Canberra, Australia.
- Roshier, D. A., A. I. Robertson, R. T. Kingsford, and D. G. Green. 2001b. Continental-scale interactions with temporary resources may explain the paradox of large populations of desert waterbirds in Australia. *Landscape Ecology* **16**:547–556.
- Sheldon, F., A. J. Boulton, and J. T. Puckridge. 2002. Conservation value of variable connectivity: aquatic invertebrate assemblages of channel and floodplain habitats of a central Australian arid-zone river, Cooper Creek. *Biological Conservation* **103**:13–31.
- Smyth, A. K., and C. D. James. 2004. Characteristics of Australia's rangelands and key design issues for monitoring biodiversity. *Austral Ecology* **29**:3–15.
- Stafford Smith, D. M., and S. R. Morton. 1990. A framework for the ecology of arid Australia. *Journal of Arid Environments* **18**:255–278.
- Stafford Smith, M., J. West, and K. Thiele. 1996. Biogeographic patterns in inland Australia: the role of isolation and uncertainty. Pages 368–378 (Chapter 326) in S. R. Morton and D. J. Mulvaney, editors. *Exploring Central Australia: Society, the Environment and the 1894 Horn Expedition*. Surrey Beatty & Sons, Sydney.
- Thackway, R. and Cresswell, I. 1995. *An Interim Biogeographic Regionalisation for Australia: a framework for setting priorities in the National Reserve System Cooperative Program*. Canberra: Reserve Systems Unit, Australian Nature Conservation Agency.
- Tongway, D. J., and J. A. Ludwig. 1994. Small-scale resource heterogeneity in semi-arid landscapes. *Pacific Conservation Biology* **1**: 201–208.