



HQD Plus

NEUTRAL pH DISINFECTANT DETERGENT

The latest in dual formula of dual chained disinfectant that not only kills all disease bearing pathogens, cleans and deodorizes applied area. Cold pressed lemon fragrance is leaves a fresh, clean scent after cleaning.

OSHA has well defined the areas in house-keeping that will require a disinfectant/cleaner. If you're looking for the best Quaternary Kill with great cleaning ability HQD Plus is your #1 choice.

DIRECTIONS FOR USE

MIX 2 OZ. PER 1 GAL. OF WATER

HQD Plus used on hard non-porous surfaces will tolerate a moderate amount of organic soil when used at 2 oz. per gallon of water.

HQD Plus provides 600 ppm of active quaternary at 2 oz. per gallon of water for a rapid four minute contact time. For all other disinfection claims, EPA states a ten minute contact time.

KILLING TYPE 1 (HIV-1) VIRUS

Remove blood and other body fluids from surface by applying "DeVour" body fluid absorbent. Remove into sack liner using personal protection, latex gloves, gowns, masks or eye coverings.

Apply HQD Plus with mop, sponge, or cloth as well as soaking where necessary. Mix fresh solution for each use, then discard solution.

Bactericidal Stability of Use-Dilution: Tests confirm that this product when diluted in 400 ppm hard water, remains effective against Pseudomonas Aeruginosa, Staphylococcus Aureus, and Salmonella Choleraesuis for up to one year when stored in a sealed container such as a spray bottle.

BENEFITS

- Virucidal
- Fungicidal
- Pseudomonacidal
- Staphylocidal
- Salmonellacidal
- Cleans & Deoderizes
- Leaves no flim
- Neutral pH
- Fresh Lemon Fragrance
- Won't dull or haze floors
- Saves on labor costs

KILLS FOLLOWING

- Acinetobacter calcoaceticus
- Bordetella bronchiseptica
- Chlamydia psittaci
- Enterobacter aerogenes
- Enterobacter cloacae
- Enterobacter cloacae NDM-1
- Enterococcus faecalis (Vancomycin Resistant)
- Escherichia coli1
- Escherichia coli NDM-1
- Fusobacterium necrophorum
- Klebsiella pneumonia NDM-1
- Klebsiella pneumoniae1
- Legionella pneumophila
- Listeria monocytogenes
- Pasteurella multocida
- Proteus mirabilis
- Proteus vulgaris
- Pseudomonas aeruginosa1
- Salmonella typhimurium
- Salmonella enterica
- Salmonella enteritidis
- Salmonella typhi
- Serratia marcescens
- Shigella flexneri
- Shigella sonnei
- Staphylococcus aureus1
- Staphylococcus aureus
- Staphylococcus aureus (MRSA)
- Streptococcus faecalis1
- Streptococcus pyogenes
- Adenovirus Type 4
- Adenovirus Type 7
- Avian influenza (H5N1)
- Avian polyomavirus
- Canine distemper virus
- Equine herpes Virus Type 1
- Feline Coronavirus
- Feline leukemia virus
- Feline Panleukopenia virus
- Feline picornavirus
- Hepatitis B
- Hepatitis C (HCV)
- Herpes Simplex Type 1
- Herpes Simplex Type 2
- HIV-1 (AIDS Virus)
- Human Corona Virus
- Infectious bovine rhinotracheitis
- Infectious bronchitis (Avian IBV)
- Influenza A/ Hong Kong
- Newcastle Disease
- Porcine Parvovirus
- Pseudorabies virus
- Rabies virus
- Respiratory Syncytical Virus
- Rotavirus
- Rubella virus
- SARS Associated Coronavirus
- Transmissible gastroenteritis virus
- Vaccinia
- Aspergillus niger
- Candida Albicans
- Trichophyton mentagrophytes