

**JEE April 2019**

Roll No.	
Candidate Name	
Application No	
Test Date	12/04/2019
Test Time	2:30 PM - 5:30 PM
Subject	Paper I EH

Section : Physics

**Q.1** Two particles are projected from the same point with the same speed  $u$  such that they have the same range  $R$ , but different maximum heights,  $h_1$  and  $h_2$ . Which of the following is correct ?

- Options**
1.  $R^2 = 4 h_1 h_2$
  2.  $R^2 = 16 h_1 h_2$
  3.  $R^2 = 2 h_1 h_2$
  4.  $R^2 = h_1 h_2$

Question Type : **MCQ**  
Question ID : **41652913418**  
Option 1 ID : **41652952451**  
Option 2 ID : **41652952450**  
Option 3 ID : **41652952452**  
Option 4 ID : **41652952453**  
Status : **Not Answered**  
Chosen Option : --

**Q.2** In an amplitude modulator circuit, the carrier wave is given by,  
 $C(t) = 4 \sin(20000 \pi t)$  while modulating signal is given by,  $m(t) = 2 \sin(2000 \pi t)$ . The values of modulation index and lower side band frequency are :

- Options**
1. 0.5 and 10 kHz
  2. 0.4 and 10 kHz
  3. 0.3 and 9 kHz
  4. 0.5 and 9 kHz

Question Type : **MCQ**  
Question ID : **41652913443**  
Option 1 ID : **41652952552**  
Option 2 ID : **41652952551**  
Option 3 ID : **41652952550**  
Option 4 ID : **41652952553**  
Status : **Answered**  
Chosen Option : **1**

**Q.3** Two sources of sound  $S_1$  and  $S_2$  produce sound waves of same frequency 660 Hz. A listener is moving from source  $S_1$  towards  $S_2$  with a constant speed  $u$  m/s and he hears 10 beats/s. The velocity of sound is 330 m/s. Then,  $u$  equals :

- Options**
1. 5.5 m/s
  2. 15.0 m/s
  3. 2.5 m/s
  4. 10.0 m/s

Question Type : **MCQ**

Question ID : **41652913429**

Option 1 ID : **41652952495**

Option 2 ID : **41652952497**


Option 3 ID : **41652952494**

Option 4 ID : **41652952496**

Status : **Not Answered**

Chosen Option : --

**Q.4** Three particles of masses 50 g, 100 g and 150 g are placed at the vertices of an equilateral triangle of side 1 m (as shown in the figure). The  $(x, y)$  coordinates of the centre of mass will be :



- Options**
1.  $\left(\frac{\sqrt{3}}{4} \text{ m}, \frac{5}{12} \text{ m}\right)$
  2.  $\left(\frac{7}{12} \text{ m}, \frac{\sqrt{3}}{8} \text{ m}\right)$
  3.  $\left(\frac{7}{12} \text{ m}, \frac{\sqrt{3}}{4} \text{ m}\right)$
  4.  $\left(\frac{\sqrt{3}}{8} \text{ m}, \frac{7}{12} \text{ m}\right)$

Question Type : **MCQ**

Question ID : **41652913422**

Option 1 ID : **41652952468**

Option 2 ID : **41652952466**

Option 3 ID : **41652952467**

Option 4 ID : **41652952469**  
Status : **Marked For Review**  
Chosen Option : **3**

**Q.5** A Carnot engine has an efficiency of  $1/6$ . When the temperature of the sink is reduced by  $62^\circ\text{C}$ , its efficiency is doubled. The temperatures of the source and the sink are, respectively,

- Options
1.  $62^\circ\text{C}$ ,  $124^\circ\text{C}$
  2.  $99^\circ\text{C}$ ,  $37^\circ\text{C}$
  3.  $124^\circ\text{C}$ ,  $62^\circ\text{C}$
  4.  $37^\circ\text{C}$ ,  $99^\circ\text{C}$

Question Type : **MCQ**  
Question ID : **41652913426**  
Option 1 ID : **41652952485**  
Option 2 ID : **41652952482**  
Option 3 ID : **41652952483**  
Option 4 ID : **41652952484**  
Status : **Answered**  
Chosen Option : **2**


**Q.6** A spring whose unstretched length is  $l$  has a force constant  $k$ . The spring is cut into two pieces of unstretched lengths  $l_1$  and  $l_2$  where,  $l_1 = nl_2$  and  $n$  is an integer. The ratio  $k_1/k_2$  of the corresponding force constants,  $k_1$  and  $k_2$  will be :

- Options
1.  $n$
  2.  $\frac{1}{n^2}$
  3.  $\frac{1}{n}$
  4.  $n^2$

Question Type : **MCQ**  
Question ID : **41652913420**  
Option 1 ID : **41652952458**  
Option 2 ID : **41652952461**  
Option 3 ID : **41652952459**  
Option 4 ID : **41652952460**  
Status : **Answered**  
Chosen Option : **4**

**Q.7**

A transparent cube of side  $d$ , made of a material of refractive index  $\mu_2$ , is immersed in a liquid of refractive index  $\mu_1$  ( $\mu_1 < \mu_2$ ). A ray is incident on the face AB at an angle  $\theta$  (shown in the figure). Total internal reflection takes place at point E on the face BC.



Then  $\theta$  must satisfy :

Options

1.  $\theta < \sin^{-1} \frac{\mu_1}{\mu_2}$
2.  $\theta > \sin^{-1} \sqrt{\frac{\mu_2^2}{\mu_1^2} - 1}$
3.  $\theta < \sin^{-1} \sqrt{\frac{\mu_2^2}{\mu_1^2} - 1}$
4.  $\theta > \sin^{-1} \frac{\mu_1}{\mu_2}$

Question Type : MCQ

Question ID : 41652913437

Option 1 ID : 41652952529

Option 2 ID : 41652952527

Option 3 ID : 41652952526

Option 4 ID : 41652952528

Status : Answered

Chosen Option : 2

**Q.8** A tuning fork of frequency 480 Hz is used in an experiment for measuring speed of sound ( $v$ ) in air by resonance tube method. Resonance is observed to occur at two successive lengths of the air column,  $l_1 = 30$  cm and  $l_2 = 70$  cm. Then,  $v$  is equal to :

Options

1.  $332 \text{ ms}^{-1}$
2.  $384 \text{ ms}^{-1}$
3.  $338 \text{ ms}^{-1}$
4.  $379 \text{ ms}^{-1}$

Question Type : MCQ

Question ID : **41652913444**  
Option 1 ID : **41652952557**  
Option 2 ID : **41652952556**  
Option 3 ID : **41652952554**  
Option 4 ID : **41652952555**  
Status : **Not Answered**  
Chosen Option : --

**Q.9** The electron in a hydrogen atom first jumps from the third excited state to the second excited state and subsequently to the first excited state. The ratio of the respective wavelengths,  $\lambda_1/\lambda_2$ , of the photons emitted in this process is :

- Options
1. 20/7
  2. 27/5
  3. 7/5
  4. 9/7

Question Type : **MCQ**  
Question ID : **41652913440**  
Option 1 ID : **41652952541**  
Option 2 ID : **41652952538**  
Option 3 ID : **41652952539**  
Option 4 ID : **41652952540**  
Status : **Answered**  
Chosen Option : **2**

**Q.10** A diatomic gas with rigid molecules does 10 J of work when expanded at constant pressure. What would be the heat energy absorbed by the gas, in this process ?

- Options
1. 25 J
  2. 35 J
  3. 30 J
  4. 40 J

Question Type : **MCQ**  
Question ID : **41652913427**  
Option 1 ID : **41652952486**  
Option 2 ID : **41652952488**  
Option 3 ID : **41652952487**  
Option 4 ID : **41652952489**  
Status : **Marked For Review**  
Chosen Option : **1**

**Q.11**

Let a total charge  $2Q$  be distributed in a sphere of radius  $R$ , with the charge density given by  $\rho(r) = kr$ , where  $r$  is the distance from the centre. Two charges  $A$  and  $B$ , of  $-Q$  each, are placed on diametrically opposite points, at equal distance,  $a$ , from the centre. If  $A$  and  $B$  do not experience any force, then :

- Options
1.  $a = 8^{-1/4} R$
  2.  $a = \frac{3R}{2^{1/4}}$
  3.  $a = 2^{-1/4} R$
  4.  $a = R/\sqrt{3}$

Question Type : **MCQ**  
 Question ID : **41652913430**  
 Option 1 ID : **41652952498**  
 Option 2 ID : **41652952501**  
 Option 3 ID : **41652952499**  
 Option 4 ID : **41652952500**  
 Status : **Not Answered**  
 Chosen Option : --

**Q.12** Consider an electron in a hydrogen atom, revolving in its second excited state (having radius  $4.65 \text{ \AA}$ ). The de-Broglie wavelength of this electron is :

- Options
1.  $3.5 \text{ \AA}$
  2.  $6.6 \text{ \AA}$
  3.  $12.9 \text{ \AA}$
  4.  $9.7 \text{ \AA}$

Question Type : **MCQ**  
 Question ID : **41652913439**  
 Option 1 ID : **41652952534**  
 Option 2 ID : **41652952535**  
 Option 3 ID : **41652952537**  
 Option 4 ID : **41652952536**  
 Status : **Not Attempted and Marked For Review**  
 Chosen Option : --

**Q.13** A solid sphere, of radius  $R$  acquires a terminal velocity  $v_1$  when falling (due to gravity) through a viscous fluid having a coefficient of viscosity  $\eta$ . The sphere is broken into 27 identical solid spheres. If each of these spheres acquires a terminal velocity,  $v_2$ , when falling through the same fluid, the ratio  $(v_1/v_2)$  equals :



Options 1. 9

2.  $1/27$ 3.  $1/9$ 

4. 27

Question Type : MCQ

Question ID : 41652913425

Option 1 ID : 41652952478

Option 2 ID : 41652952481


Option 3 ID : 41652952479

Option 4 ID : 41652952480

Status : Not Answered

Chosen Option : --

- Q.14 A smooth wire of length  $2\pi r$  is bent into a circle and kept in a vertical plane. A bead can slide smoothly on the wire. When the circle is rotating with angular speed  $\omega$  about the vertical diameter AB, as shown in figure, the bead is at rest with respect to the circular ring at position P as shown. Then the value of  $\omega^2$  is equal to :



Options

1.  $\frac{\sqrt{3}g}{2r}$ 2.  $2g/(r\sqrt{3})$ 3.  $(g\sqrt{3})/r$ 4.  $2g/r$ 

Question Type : MCQ

Question ID : 41652913421

Option 1 ID : 41652952463

Option 2 ID : 41652952465

Option 3 ID : 41652952464

Option 4 ID : 41652952462

Status : Not Answered

Chosen Option : --

Q.15

A particle is moving with speed  $v = b\sqrt{x}$  along positive  $x$ -axis. Calculate the speed of the particle at time  $t = \tau$  (assume that the particle is at origin at  $t = 0$ ).

Options

1.  $\frac{b^2\tau}{4}$
2.  $\frac{b^2\tau}{2}$
3.  $b^2\tau$
4.  $\frac{b^2\tau}{\sqrt{2}}$

Question Type : **MCQ**

Question ID : **41652913417**

Option 1 ID : **41652952446**

Option 2 ID : **41652952448**

Option 3 ID : **41652952449**

Option 4 ID : **41652952447**

Status : **Answered**

Chosen Option : 1

**Q.16** The ratio of the weights of a body on the Earth's surface to that on the surface of a planet is 9 : 4. The mass of the planet is  $\frac{1}{9}$ th of that of the Earth. If 'R' is the radius of the Earth, what is the radius of the planet ? (Take the planets to have the same mass density)

Options

1.  $\frac{R}{3}$
2.  $\frac{R}{4}$
3.  $\frac{R}{9}$
4.  $\frac{R}{2}$

Question Type : **MCQ**

Question ID : **41652913423**

Option 1 ID : **41652952471**

Option 2 ID : **41652952473**

Option 3 ID : **41652952472**

Option 4 ID : **41652952470**

Status : **Answered**

Chosen Option : 3

**Q.17**



A system of three polarizers  $P_1, P_2, P_3$  is set up such that the pass axis of  $P_3$  is crossed with respect to that of  $P_1$ . The pass axis of  $P_2$  is inclined at  $60^\circ$  to the pass axis of  $P_3$ . When a beam of unpolarized light of intensity  $I_0$  is incident on  $P_1$ , the intensity of light transmitted by the three polarizers is  $I$ . The ratio  $(I_0/I)$  equals (nearly) :

- Options
1. 5.33
  2. 16.00
  3. 10.67
  4. 1.80

Question Type : **MCQ**  
Question ID : **41652913438**  
Option 1 ID : **41652952531**  
Option 2 ID : **41652952533**  
Option 3 ID : **41652952532**  
Option 4 ID : **41652952530**  
Status : **Answered**  
Chosen Option : **3**

**Q.18** A uniform cylindrical rod of length  $L$  and radius  $r$ , is made from a material whose Young's modulus of Elasticity equals  $Y$ . When this rod is heated by temperature  $T$  and simultaneously subjected to a net longitudinal compressional force  $F$ , its length remains unchanged. The coefficient of volume expansion, of the material of the rod, is (nearly) equal to :

- Options
1.  $9F/(\pi r^2 Y T)$
  2.  $6F/(\pi r^2 Y T)$
  3.  $3F/(\pi r^2 Y T)$
  4.  $F/(3\pi r^2 Y T)$

Question Type : **MCQ**  
Question ID : **41652913424**  
Option 1 ID : **41652952475**  
Option 2 ID : **41652952474**  
Option 3 ID : **41652952476**  
Option 4 ID : **41652952477**  
Status : **Not Answered**  
Chosen Option : **--**

**Q.19**

The number density of molecules of a gas depends on their distance  $r$  from the origin as,  $n(r) = n_0 e^{-\alpha r^4}$ . Then the total number of molecules is proportional to :

- Options
1.  $n_0 \alpha^{-3/4}$
  2.  $\sqrt{n_0} \alpha^{1/2}$
  3.  $n_0 \alpha^{1/4}$
  4.  $n_0 \alpha^{-3}$

Question Type : **MCQ**

Question ID : **41652913416**

Option 1 ID : **41652952444**

Option 2 ID : **41652952443**

Option 3 ID : **41652952445**

Option 4 ID : **41652952442**

Status : **Not Attempted and Marked For Review**

Chosen Option : --

**Q.20** A small speaker delivers 2 W of audio output. At what distance from the speaker will one detect 120 dB intensity sound ? [Given reference intensity of sound as  $10^{-12} \text{ W/m}^2$ ]

- Options
1. 40 cm
  2. 20 cm
  3. 10 cm
  4. 30 cm

Question Type : **MCQ**

Question ID : **41652913428**

Option 1 ID : **41652952493**

Option 2 ID : **41652952491**

Option 3 ID : **41652952490**

Option 4 ID : **41652952492**

Status : **Not Answered**


Chosen Option : --

**Q.21** Half lives of two radioactive nuclei A and B are 10 minutes and 20 minutes, respectively. If, initially a sample has equal number of nuclei, then after 60 minutes, the ratio of decayed numbers of nuclei A and B will be :

- Options
1. 3 : 8
  2. 1 : 8
  3. 8 : 1
  4. 9 : 8

Question Type : **MCQ**Question ID : **41652913441**Option 1 ID : **41652952545**Option 2 ID : **41652952542**Option 3 ID : **41652952544**Option 4 ID : **41652952543**Status : **Answered**Chosen Option : **2**


- Q.22** An electron, moving along the  $x$ -axis with an initial energy of 100 eV, enters a region of magnetic field  $\vec{B} = (1.5 \times 10^{-3} \text{T}) \hat{k}$  at S (See figure). The field extends between  $x=0$  and  $x=2$  cm. The electron is detected at the point Q on a screen placed 8 cm away from the point S. The distance  $d$  between P and Q (on the screen) is :  
(electron's charge =  $1.6 \times 10^{-19} \text{C}$ , mass of electron =  $9.1 \times 10^{-31} \text{kg}$ )



- Options**
1. 11.65 cm
  2. 12.87 cm
  3. 1.22 cm
  4. 2.25 cm

Question Type : **MCQ**Question ID : **41652913433**Option 1 ID : **41652952512**Option 2 ID : **41652952511**Option 3 ID : **41652952510**Option 4 ID : **41652952513**Status : **Not Attempted and Marked For Review**Chosen Option : **--****Q.23**

In the given circuit, the charge on  $4 \mu\text{F}$  capacitor will be :



- Options
1.  $5.4 \mu\text{C}$
  2.  $9.6 \mu\text{C}$
  3.  $13.4 \mu\text{C}$
  4.  $24 \mu\text{C}$

Question Type : **MCQ**

Question ID : **41652913431**

Option 1 ID : **41652952504**

Option 2 ID : **41652952502**

Option 3 ID : **41652952505**

Option 4 ID : **41652952503**

Status : **Answered**

Chosen Option : **1**

**Q.24** One kg of water, at  $20^\circ\text{C}$ , is heated in an electric kettle whose heating element has a mean (temperature averaged) resistance of  $20 \Omega$ . The rms voltage in the mains is  $200 \text{ V}$ . Ignoring heat loss from the kettle, time taken for water to evaporate fully, is close to :

[Specific heat of water =  $4200 \text{ J}/(\text{kg } ^\circ\text{C})$ ,  
Latent heat of water =  $2260 \text{ kJ}/\text{kg}$ ]

- Options
1. 16 minutes
  2. 22 minutes
  3. 3 minutes
  4. 10 minutes

Question Type : **MCQ**

Question ID : **41652913432**

Option 1 ID : **41652952509**

Option 2 ID : **41652952508**

Option 3 ID : **41652952506**

Option 4 ID : **41652952507**

Status : **Answered**

Chosen Option : **3**

**Q.25**

A moving coil galvanometer, having a resistance  $G$ , produces full scale deflection when a current  $I_g$  flows through it. This galvanometer can be converted into (i) an ammeter of range 0 to  $I_0$  ( $I_0 > I_g$ ) by connecting a shunt resistance  $R_A$  to it and (ii) into a voltmeter of range 0 to  $V$  ( $V = GI_0$ ) by connecting a series resistance  $R_V$  to it. Then,

Options

$$R_A R_V = G^2 \left( \frac{I_0 - I_g}{I_g} \right) \text{ and}$$

1.

$$\frac{R_A}{R_V} = \left( \frac{I_g}{I_0 - I_g} \right)^2$$

2.  $R_A R_V = G^2$  and  $\frac{R_A}{R_V} = \left( \frac{I_g}{I_0 - I_g} \right)^2$

$$R_A R_V = G^2 \left( \frac{I_g}{I_0 - I_g} \right) \text{ and}$$

3.

$$\frac{R_A}{R_V} = \left( \frac{I_0 - I_g}{I_g} \right)^2$$

4.  $R_A R_V = G^2$  and  $\frac{R_A}{R_V} = \frac{I_g}{I_0 - I_g}$

Question Type : **MCQ**

Question ID : **41652913445**

Option 1 ID : **41652952561**

Option 2 ID : **41652952558**

Option 3 ID : **41652952560**

Option 4 ID : **41652952559**


Status : **Answered**

Chosen Option : **3**

Q.26

Find the magnetic field at point P due to a straight line segment AB of length 6 cm carrying a current of 5 A. (See figure)

$$(\mu_0 = 4\pi \times 10^{-7} \text{ N-A}^{-2})$$



- Options
1.  $2.0 \times 10^{-5} \text{ T}$
  2.  $1.5 \times 10^{-5} \text{ T}$
  3.  $3.0 \times 10^{-5} \text{ T}$
  4.  $2.5 \times 10^{-5} \text{ T}$

Question Type : MCQ

Question ID : 41652913434

Option 1 ID : 41652952515

Option 2 ID : 41652952514

Option 3 ID : 41652952517

Option 4 ID : 41652952516

Status : Answered

Chosen Option : 1

Q.27 A plane electromagnetic wave having a frequency  $\nu = 23.9 \text{ GHz}$  propagates along the positive  $z$ -direction in free space. The peak value of the Electric Field is  $60 \text{ V/m}$ . Which among the following is the acceptable magnetic field component in the electromagnetic wave ?

- Options
1.  $\vec{B} = 2 \times 10^{-7} \sin(0.5 \times 10^3 z + 1.5 \times 10^{11} t) \hat{i}$
  2.  $\vec{B} = 2 \times 10^{-7} \sin(0.5 \times 10^3 z - 1.5 \times 10^{11} t) \hat{i}$
  3.  $\vec{B} = 60 \sin(0.5 \times 10^3 x + 1.5 \times 10^{11} t) \hat{k}$
  4.  $\vec{B} = 2 \times 10^{-7} \sin(1.5 \times 10^2 x + 0.5 \times 10^{11} t) \hat{j}$

Question Type : MCQ

Question ID : 41652913436

Option 1 ID : 41652952524

Option 2 ID : 41652952523

Option 3 ID : 41652952522


Option 4 ID : 41652952525

Status : Not Answered



Chosen Option : --

Q.28 Consider the LR circuit shown in the figure.  
If the switch S is closed at  $t=0$  then the amount of charge that passes through the battery between  $t=0$  and  $t=\frac{L}{R}$  is :



Options

1.  $\frac{2.7 EL}{R^2}$
2.  $\frac{EL}{2.7R^2}$
3.  $\frac{7.3 EL}{R^2}$
4.  $\frac{EL}{7.3R^2}$

Question Type : MCQ

Question ID : 41652913435

Option 1 ID : 41652952519

Option 2 ID : 41652952520


Option 3 ID : 41652952518

Option 4 ID : 41652952521

Status : Not Answered

Chosen Option : --

Q.29 Figure shows a DC voltage regulator circuit, with a Zener diode of breakdown voltage = 6V. If the unregulated input voltage varies between 10 V to 16 V, then what is the maximum Zener current ?



- Options
1. 2.5 mA
  2. 1.5 mA

3. 7.5 mA

4. 3.5 mA

Question Type : MCQ

Question ID : 41652913442

Option 1 ID : 41652952548

Option 2 ID : 41652952547


Option 3 ID : 41652952546

Option 4 ID : 41652952549

Status : Not Answered

Chosen Option : --

**Q.30** A block of mass 5 kg is (i) pushed in case (A) and (ii) pulled in case (B), by a force  $F = 20$  N, making an angle of  $30^\circ$  with the horizontal, as shown in the figures. The coefficient of friction between the block and floor is  $\mu = 0.2$ . The difference between the accelerations of the block, in case (B) and case (A) will be : ( $g = 10 \text{ ms}^{-2}$ )



- Options**
1.  $0.4 \text{ ms}^{-2}$
  2.  $3.2 \text{ ms}^{-2}$
  3.  $0.8 \text{ ms}^{-2}$
  4.  $0 \text{ ms}^{-2}$

Question Type : MCQ

Question ID : 41652913419

Option 1 ID : 41652952456

Option 2 ID : 41652952455

Option 3 ID : 41652952457

Option 4 ID : 41652952454

Status : Not Answered

Chosen Option : --

Section : Chemistry

**Q.1** Thermal decomposition of a Mn compound (X) at 513 K results in compound Y,  $\text{MnO}_2$  and a gaseous product.  $\text{MnO}_2$  reacts with  $\text{NaCl}$  and concentrated  $\text{H}_2\text{SO}_4$  to give a pungent gas Z. X, Y, and Z, respectively, are :

- Options**
1.  $\text{K}_3\text{MnO}_4$ ,  $\text{K}_2\text{MnO}_4$  and  $\text{Cl}_2$
  2.  $\text{K}_2\text{MnO}_4$ ,  $\text{KMnO}_4$  and  $\text{SO}_2$

3.  $\text{KMnO}_4$ ,  $\text{K}_2\text{MnO}_4$  and  $\text{Cl}_2$

4.  $\text{K}_2\text{MnO}_4$ ,  $\text{KMnO}_4$  and  $\text{Cl}_2$

Question Type : **MCQ**

Question ID : **41652913461**

Option 1 ID : **41652952625**

Option 2 ID : **41652952622**


Option 3 ID : **41652952623**

Option 4 ID : **41652952624**

Status : **Answered**

Chosen Option : **2**

**Q.2**  $\text{NO}_2$  required for a reaction is produced by the decomposition of  $\text{N}_2\text{O}_5$  in  $\text{CCl}_4$  as per the equation,



The initial concentration of  $\text{N}_2\text{O}_5$  is  $3.00 \text{ mol L}^{-1}$  and it is  $2.75 \text{ mol L}^{-1}$  after 30 minutes. The rate of formation of  $\text{NO}_2$  is :

- Options
1.  $4.167 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$
  2.  $1.667 \times 10^{-2} \text{ mol L}^{-1} \text{ min}^{-1}$
  3.  $8.333 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$
  4.  $2.083 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$

Question Type : **MCQ**

Question ID : **41652913474**

Option 1 ID : **41652952674**

Option 2 ID : **41652952677**

Option 3 ID : **41652952675**

Option 4 ID : **41652952676**

Status : **Not Answered**

Chosen Option : **--**

**Q.3** The pair that has similar atomic radii is :

- Options
1. Mn and Re
  2. Ti and Hf
  3. Sc and Ni
  4. Mo and W

Question Type : **MCQ**

Question ID : **41652913462**

Option 1 ID : **41652952627**

Option 2 ID : **41652952626**

Option 3 ID : **41652952628**

Option 4 ID : **41652952629**

Status : **Answered**

Chosen Option : **4**

**Q.4** The INCORRECT statement is :

Options

1. Lithium is the strongest reducing agent among the alkali metals.
2. Lithium is least reactive with water among the alkali metals.
3.  $\text{LiNO}_3$  decomposes on heating to give  $\text{LiNO}_2$  and  $\text{O}_2$ .
4.  $\text{LiCl}$  crystallises from aqueous solution as  $\text{LiCl}\cdot 2\text{H}_2\text{O}$ .

Question Type : **MCQ**  
Question ID : **41652913459**  
Option 1 ID : **41652952615**  
Option 2 ID : **41652952616**  
Option 3 ID : **41652952614**  
Option 4 ID : **41652952617**  
Status : **Answered**  
Chosen Option : **4**

**Q.5** The C – C bond length is maximum in :

- Options
1. graphite
  2.  $\text{C}_{70}$
  3.  $\text{C}_{60}$
  4. diamond

Question Type : **MCQ**  
Question ID : **41652913460**  
Option 1 ID : **41652952619**  
Option 2 ID : **41652952621**  
Option 3 ID : **41652952618**  
Option 4 ID : **41652952620**  
Status : **Answered**  
Chosen Option : **4**

**Q.6** A solution is prepared by dissolving 0.6 g of urea (molar mass =  $60 \text{ g mol}^{-1}$ ) and 1.8 g of glucose (molar mass =  $180 \text{ g mol}^{-1}$ ) in 100 mL of water at  $27^\circ\text{C}$ . The osmotic pressure of the solution is :  
( $R = 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$ )

- Options
1. 8.2 atm
  2. 2.46 atm
  3. 4.92 atm
  4. 1.64 atm

Question Type : **MCQ**  
Question ID : **41652913470**  
Option 1 ID : **41652952661**  
Option 2 ID : **41652952659**  
Option 3 ID : **41652952658**

Option 4 ID : **41652952660**  
Status : **Answered**  
Chosen Option : **3**

**Q.7** In comparison to boron, beryllium has :

- Options
1. lesser nuclear charge and lesser first ionisation enthalpy.
  2. greater nuclear charge and lesser first ionisation enthalpy.
  3. greater nuclear charge and greater first ionisation enthalpy.
  4. lesser nuclear charge and greater first ionisation enthalpy.

Question Type : **MCQ**  
Question ID : **41652913456**  
Option 1 ID : **41652952604**  
Option 2 ID : **41652952605**  
Option 3 ID : **41652952602**  
Option 4 ID : **41652952603**  
Status : **Answered**  
Chosen Option : **4**

**Q.8** The decreasing order of electrical conductivity of the following aqueous solutions is :





- 0.1 M Formic acid (A),  
0.1 M Acetic acid (B),  
0.1 M Benzoic acid (C).

- Options
1.  $A > C > B$
  2.  $C > B > A$
  3.  $A > B > C$
  4.  $C > A > B$

Question Type : **MCQ**  
Question ID : **41652913473**  
Option 1 ID : **41652952673**  
Option 2 ID : **41652952672**  
Option 3 ID : **41652952670**  
Option 4 ID : **41652952671**  
Status : **Answered**  
Chosen Option : **3**

**Q.9** What will be the major product when m-cresol is reacted with propargyl bromide ( $\text{HC} \equiv \text{C} - \text{CH}_2\text{Br}$ ) in presence of  $\text{K}_2\text{CO}_3$  in acetone ?

Options



Question Type : **MCQ**

Question ID : **41652913453**

Option 1 ID : **41652952590**

Option 2 ID : **41652952591**

Option 3 ID : **41652952592**

Option 4 ID : **41652952593**

Status : **Not Answered**

Chosen Option : --

**Q.10** The molar solubility of  $\text{Cd}(\text{OH})_2$  is  $1.84 \times 10^{-5} \text{ M}$  in water. The expected solubility of  $\text{Cd}(\text{OH})_2$  in a buffer solution of  $\text{pH} = 12$  is :

- Options
1.  $1.84 \times 10^{-9} \text{ M}$
  2.  $\frac{2.49}{1.84} \times 10^{-9} \text{ M}$
  3.  $6.23 \times 10^{-11} \text{ M}$
  4.  $2.49 \times 10^{-10} \text{ M}$

Question Type : **MCQ**

Question ID : **41652913472**

Option 1 ID : **41652952669**

Option 2 ID : **41652952668**

Option 3 ID : **41652952666**

Option 4 ID : **41652952667**

Status : **Not Answered**





Chosen Option : --

**Q.11**



Benzene diazonium chloride on reaction with aniline in the presence of dilute hydrochloric acid gives :

Options

1. 
2. 
3. 
4. 

Question Type : **MCQ**

Question ID : **41652913451**

Option 1 ID : **41652952585**

Option 2 ID : **41652952584**

Option 3 ID : **41652952582**

Option 4 ID : **41652952583**

Status : **Answered**

Chosen Option : **3**

Q.12 The correct statement is :

Options

1. leaching of bauxite using concentrated NaOH solution gives sodium aluminate and sodium silicate.
2. the Hall-Heroult process is used for the production of aluminium and iron.
3. pig iron is obtained from cast iron.
4. the blistered appearance of copper during the metallurgical process is due to the evolution of CO<sub>2</sub>.

Question Type : **MCQ**

Question ID : **41652913457**

Option 1 ID : **41652952608**

Option 2 ID : **41652952609**

Option 3 ID : **41652952607**

Option 4 ID : **41652952606**

Status : **Answered**

Chosen Option : **3**

Q.13 The primary pollutant that leads to photochemical smog is :

- Options 1. acrolein

2. nitrogen oxides
3. ozone
4. sulphur dioxide

Question Type : MCQ

Question ID : 41652913465

Option 1 ID : 41652952640

Option 2 ID : 41652952641


Option 3 ID : 41652952638

Option 4 ID : 41652952639

Status : Answered

Chosen Option : 2

Q.14 The IUPAC name for the following compound is :



- Options
1. 3-methyl-4-(3-methylprop-1-enyl)-1-heptyne
  2. 3,5-dimethyl-4-propylhept-6-en-1-yne
  3. 3-methyl-4-(1-methylprop-2-ynyl)-1-heptene
  4. 3,5-dimethyl-4-propylhept-1-en-6-yne

Question Type : MCQ

Question ID : 41652913447

Option 1 ID : 41652952567

Option 2 ID : 41652952566

Option 3 ID : 41652952569

Option 4 ID : 41652952568

Status : Answered


Chosen Option : 4

Q.15 The temporary hardness of a water sample is due to compound X. Boiling this sample converts X to compound Y. X and Y, respectively, are :

- Options
1.  $Mg(HCO_3)_2$  and  $Mg(OH)_2$
  2.  $Ca(HCO_3)_2$  and  $Ca(OH)_2$
  3.  $Mg(HCO_3)_2$  and  $MgCO_3$
  4.  $Ca(HCO_3)_2$  and  $CaO$

Question Type : **MCQ**Question ID : **41652913458**Option 1 ID : **41652952610**Option 2 ID : **41652952612**Option 3 ID : **41652952611**Option 4 ID : **41652952613**Status : **Answered**Chosen Option : **2**

**Q.16** In the following skew conformation of ethane,  $H^I - C - C - H^{II}$  dihedral angle is :



- Options
1.  $58^\circ$
  2.  $149^\circ$
  3.  $151^\circ$
  4.  $120^\circ$

Question Type : **MCQ**Question ID : **41652913446**Option 1 ID : **41652952562**Option 2 ID : **41652952564**Option 3 ID : **41652952563**Option 4 ID : **41652952565**Status : **Not Answered**Chosen Option : **--**

**Q.17** Among the following, the energy of 2s orbital is lowest in :

- Options
1. K
  2. H
  3. Li
  4. Na

Question Type : **MCQ**Question ID : **41652913468**Option 1 ID : **41652952651**Option 2 ID : **41652952650**Option 3 ID : **41652952653**Option 4 ID : **41652952652**Status : **Answered**Chosen Option : **3**

**Q.18** Which one of the following is likely to give a precipitate with  $AgNO_3$  solution ?

Options

1.  $\text{CH}_2 = \text{CH} - \text{Cl}$
2.  $\text{CCl}_4$
3.  $\text{CHCl}_3$
4.  $(\text{CH}_3)_3\text{CCl}$

Question Type : **MCQ**Question ID : **41652913455**Option 1 ID : **41652952601**Option 2 ID : **41652952598**Option 3 ID : **41652952599**Option 4 ID : **41652952600**Status : **Answered**Chosen Option : **1**





**Q.19** 25 g of an unknown hydrocarbon upon burning produces 88 g of  $\text{CO}_2$  and 9 g of  $\text{H}_2\text{O}$ . This unknown hydrocarbon contains :

- Options
1. 20 g of carbon and 5 g of hydrogen
  2. 22 g of carbon and 3 g of hydrogen
  3. 24 g of carbon and 1 g of hydrogen
  4. 18 g of carbon and 7 g of hydrogen

Question Type : **MCQ**Question ID : **41652913466**Option 1 ID : **41652952642**Option 2 ID : **41652952644**Option 3 ID : **41652952643**Option 4 ID : **41652952645**Status : **Answered**Chosen Option : **3**

**Q.20** Heating of 2-chloro-1-phenylbutane with  $\text{EtOK}/\text{EtOH}$  gives X as the major product. Reaction of X with  $\text{Hg}(\text{OAc})_2/\text{H}_2\text{O}$  followed by  $\text{NaBH}_4$  gives Y as the major product. Y is :

Options

1. 
2. 
3. 
4. 

Question Type : **MCQ**

Question ID : 41652913452

Option 1 ID : 41652952586

Option 2 ID : 41652952588

Option 3 ID : 41652952587

Option 4 ID : 41652952589

Status : Answered

Chosen Option : 1

Q.21 The compound used in the treatment of lead poisoning is :

- Options
1. D-penicillamine
  2. desferrioxime B
  3. Cis-platin
  4. EDTA

Question Type : MCQ

Question ID : 41652913463

Option 1 ID : 41652952630

Option 2 ID : 41652952633

Option 3 ID : 41652952632

Option 4 ID : 41652952631

Status : Answered

Chosen Option : 3

Q.22 Which of the given statements is INCORRECT about glycogen ?

- Options
1. It is a straight chain polymer similar to amylose.
  2. Only  $\alpha$ -linkages are present in the molecule.
  3. It is present in animal cells.
  4. It is present in some yeast and fungi.

Question Type : MCQ

Question ID : 41652913449

Option 1 ID : 41652952575

Option 2 ID : 41652952577


Option 3 ID : 41652952574

Option 4 ID : 41652952576

Status : Answered




Chosen Option : 2

Q.23 Consider the following reactions :




'A' is :

- Options
1.  $\text{CH} \equiv \text{CH}$

Question Type : **MCQ**Question ID : **41652913448**Option 1 ID : **41652952570**Option 2 ID : **41652952572**Option 3 ID : **41652952571**Option 4 ID : **41652952573**Status : **Answered**Chosen Option : **2**

**Q.24** The correct name of the following polymer is :



- Options
1. Polyisobutane
  2. Polytert-butylene
  3. Polyisoprene
  4. Polyisobutylene

Question Type : **MCQ**Question ID : **41652913450**Option 1 ID : **41652952578**Option 2 ID : **41652952580**Option 3 ID : **41652952581**Option 4 ID : **41652952579**Status : **Answered**Chosen Option : **4**

**Q.25** An 'Assertion' and a 'Reason' are given below. Choose the correct answer from the following options :

**Assertion (A)** : Vinyl halides do not undergo nucleophilic substitution easily.

**Reason (R)** : Even though the intermediate carbocation is stabilized by loosely held  $\pi$ -electrons, the cleavage is difficult because of strong bonding.

- Options
1. Both (A) and (R) are wrong statements.
  - Both (A) and (R) are correct statements and (R) is the correct explanation of (A).



- Both (A) and (R) are correct
3. statements but (R) is not the correct explanation of (A).
4. (A) is a correct statement but (R) is a wrong statement.

Question Type : **MCQ**

Question ID : **41652913454**

Option 1 ID : **41652952597**

Option 2 ID : **41652952594**

Option 3 ID : **41652952595**

Option 4 ID : **41652952596**

Status : **Answered**

Chosen Option : **3**

**Q.26** The ratio of number of atoms present in a simple cubic, body centered cubic and face centered cubic structure are, respectively :

- Options
1. 8 : 1 : 6
  2. 1 : 2 : 4
  3. 4 : 2 : 1
  4. 4 : 2 : 3

Question Type : **MCQ**

Question ID : **41652913467**

Option 1 ID : **41652952646**

Option 2 ID : **41652952647**

Option 3 ID : **41652952649**

Option 4 ID : **41652952648**

Status : **Answered**

Chosen Option : **2**

**Q.27** In which one of the following equilibria,  $K_p \neq K_c$  ?

- Options
1.  $2\text{C(s)} + \text{O}_2\text{(g)} \rightleftharpoons 2\text{CO(g)}$
  2.  $2\text{HI(g)} \rightleftharpoons \text{H}_2\text{(g)} + \text{I}_2\text{(g)}$
  3.  $\text{NO}_2\text{(g)} + \text{SO}_2\text{(g)} \rightleftharpoons \text{NO(g)} + \text{SO}_3\text{(g)}$
  4.  $2\text{NO(g)} \rightleftharpoons \text{N}_2\text{(g)} + \text{O}_2\text{(g)}$

Question Type : **MCQ**

Question ID : **41652913471**

Option 1 ID : **41652952664**

Option 2 ID : **41652952663**

Option 3 ID : **41652952662**

Option 4 ID : **41652952665**

Status : **Not Answered**

Chosen Option : **--**

**Q.28**

The coordination numbers of Co and Al in  $[\text{Co}(\text{Cl})(\text{en})_2]\text{Cl}$  and  $\text{K}_3[\text{Al}(\text{C}_2\text{O}_4)_3]$ , respectively, are :

(en = ethane-1, 2-diamine)

- Options
1. 5 and 3
  2. 3 and 3
  3. 6 and 6
  4. 5 and 6

Question Type : **MCQ**  
Question ID : **41652913464**  
Option 1 ID : **41652952637**  
Option 2 ID : **41652952635**  
Option 3 ID : **41652952634**  
Option 4 ID : **41652952636**  
Status : **Answered**  
Chosen Option : **2**

**Q.29** The **INCORRECT** match in the following is :

- Options
1.  $\Delta G^0 < 0, K > 1$
  2.  $\Delta G^0 = 0, K = 1$
  3.  $\Delta G^0 > 0, K < 1$
  4.  $\Delta G^0 < 0, K < 1$

Question Type : **MCQ**  
Question ID : **41652913469**  
Option 1 ID : **41652952654**  
Option 2 ID : **41652952657**  
Option 3 ID : **41652952656**  
Option 4 ID : **41652952655**  
Status : **Answered**  
Chosen Option : **4**

**Q.30** Among the following, the **INCORRECT** statement about colloids is :

- Options
1. They can scatter light.
  2. They are larger than small molecules and have high molar mass.
  3. The osmotic pressure of a colloidal solution is of higher order than the true solution at the same concentration.
  4. The range of diameters of colloidal particles is between 1 and 1000 nm.

Question Type : **MCQ**  
Question ID : **41652913475**

Option 1 ID : 41652952679  
 Option 2 ID : 41652952681  
 Option 3 ID : 41652952680  
 Option 4 ID : 41652952678  
 Status : Answered  
 Chosen Option : 2

Section : Mathematics

Q.1

The derivative of  $\tan^{-1}\left(\frac{\sin x - \cos x}{\sin x + \cos x}\right)$ ,

with respect to  $\frac{x}{2}$ , where  $\left(x \in \left(0, \frac{\pi}{2}\right)\right)$  is :

- Options
1. 1
  2.  $\frac{2}{3}$
  3.  $\frac{1}{2}$
  4. 2

Question Type : MCQ  
 Question ID : 41652913486  
 Option 1 ID : 41652952722  
 Option 2 ID : 41652952725  
 Option 3 ID : 41652952723  
 Option 4 ID : 41652952724  
 Status : Answered  
 Chosen Option : 3

Q.2

For an initial screening of an admission test, a candidate is given fifty problems to solve. If the probability that the candidate can solve any problem is  $\frac{4}{5}$ , then the probability that he is unable to solve less than two problems is :

- Options
1.  $\frac{201}{5} \left(\frac{1}{5}\right)^{49}$
  2.  $\frac{316}{25} \left(\frac{4}{5}\right)^{48}$
  3.  $\frac{54}{5} \left(\frac{4}{5}\right)^{49}$
  4.  $\frac{164}{25} \left(\frac{1}{5}\right)^{48}$

Question Type : MCQ  
 Question ID : 41652913502  
 Option 1 ID : 41652952786  
 Option 2 ID : 41652952788  
 Option 3 ID : 41652952787

Option 4 ID : **41652952789**  
 Status : **Not Answered**  
 Chosen Option : --

**Q.3** A value of  $\alpha$  such that

$$\int_{\alpha}^{\alpha+1} \frac{dx}{(x+\alpha)(x+\alpha+1)} = \log_e \left( \frac{9}{8} \right) \text{ is :}$$

- Options**
1.  $-2$
  2.  $\frac{1}{2}$
  3.  $-\frac{1}{2}$
  4.  $2$

Question Type : **MCQ**  
 Question ID : **41652913490**  
 Option 1 ID : **41652952740**  
 Option 2 ID : **41652952738**  
 Option 3 ID : **41652952741**  
 Option 4 ID : **41652952739**  
 Status : **Not Answered**  
 Chosen Option : --

**Q.4** Let  $\alpha \in (0, \pi/2)$  be fixed. If the integral

$$\int \frac{\tan x + \tan \alpha}{\tan x - \tan \alpha} dx =$$

$A(x) \cos 2\alpha + B(x) \sin 2\alpha + C$ , where  $C$  is a constant of integration, then the functions  $A(x)$  and  $B(x)$  are respectively :

- Options**
1.  $x + \alpha$  and  $\log_e |\sin(x + \alpha)|$
  2.  $x - \alpha$  and  $\log_e |\sin(x - \alpha)|$
  3.  $x - \alpha$  and  $\log_e |\cos(x - \alpha)|$
  4.  $x + \alpha$  and  $\log_e |\sin(x - \alpha)|$

Question Type : **MCQ**  
 Question ID : **41652913489**  
 Option 1 ID : **41652952737**  
 Option 2 ID : **41652952734**  
 Option 3 ID : **41652952736**  
 Option 4 ID : **41652952735**  
 Status : **Not Answered**  
 Chosen Option : --

**Q.5**

The angle of elevation of the top of a vertical tower standing on a horizontal plane is observed to be  $45^\circ$  from a point A on the plane. Let B be the point 30 m vertically above the point A. If the angle of elevation of the top of the tower from B be  $30^\circ$ , then the distance (in m) of the foot of the tower from the point A is :

- Options
1.  $15(3 + \sqrt{3})$
  2.  $15(5 - \sqrt{3})$
  3.  $15(3 - \sqrt{3})$
  4.  $15(1 + \sqrt{3})$

Question Type : **MCQ**  
Question ID : **41652913504**  
Option 1 ID : **41652952795**  
Option 2 ID : **41652952797**  
Option 3 ID : **41652952794**  
Option 4 ID : **41652952796**  
Status : **Marked For Review**  
Chosen Option : **2**

**Q.6** Let S be the set of all  $\alpha \in \mathbb{R}$  such that the equation,  $\cos 2x + \alpha \sin x = 2\alpha - 7$  has a solution. Then S is equal to :

- Options
1.  $\mathbb{R}$
  2.  $[1, 4]$
  3.  $[3, 7]$
  4.  $[2, 6]$

Question Type : **MCQ**  
Question ID : **41652913503**  
Option 1 ID : **41652952790**  
Option 2 ID : **41652952792**  
Option 3 ID : **41652952793**  
Option 4 ID : **41652952791**  
Status : **Not Answered**  
Chosen Option : **--**

**Q.7** A plane which bisects the angle between the two given planes  $2x - y + 2z - 4 = 0$  and  $x + 2y + 2z - 2 = 0$ , passes through the point :

- Options
1.  $(1, -4, 1)$
  2.  $(1, 4, -1)$
  3.  $(2, 4, 1)$

4.  $(2, -4, 1)$

Question Type : **MCQ**

Question ID : **41652913498**

Option 1 ID : **41652952770**

Option 2 ID : **41652952773**

Option 3 ID : **41652952771**

Option 4 ID : **41652952772**

Status : **Not Attempted and Marked For Review**

Chosen Option : --

**Q.8**

$$\lim_{x \rightarrow 0} \frac{x + 2 \sin x}{\sqrt{x^2 + 2 \sin x + 1} - \sqrt{\sin^2 x - x + 1}}$$

is :

Options

1. 6

2. 2

3. 3

4. 1

Question Type : **MCQ**

Question ID : **41652913485**

Option 1 ID : **41652952721**

Option 2 ID : **41652952719**

Option 3 ID : **41652952720**

Option 4 ID : **41652952718**

Status : **Not Answered**

Chosen Option : --

**Q.9**

A group of students comprises of 5 boys and n girls. If the number of ways, in which a team of 3 students can randomly be selected from this group such that there is at least one boy and at least one girl in each team, is 1750, then n is equal to :

Options

1. 28

2. 27

3. 25

4. 24

Question Type : **MCQ**

Question ID : **41652913481**

Option 1 ID : **41652952705**

Option 2 ID : **41652952704**

Option 3 ID : **41652952703**

Option 4 ID : **41652952702**

Status : **Answered**

Chosen Option : **3**

**Q.10**

An ellipse, with foci at  $(0, 2)$  and  $(0, -2)$  and minor axis of length 4, passes through which of the following points ?



Options

1.  $(\sqrt{2}, 2)$
2.  $(2, \sqrt{2})$
3.  $(2, 2\sqrt{2})$
4.  $(1, 2\sqrt{2})$

Question Type : **MCQ**Question ID : **41652913497**Option 1 ID : **41652952768**Option 2 ID : **41652952766**Option 3 ID : **41652952769**Option 4 ID : **41652952767**Status : **Answered**

Chosen Option : 1

**Q.11** The Boolean expression  $\sim(p \Rightarrow (\sim q))$  is equivalent to :

Options

1.  $p \wedge q$
2.  $q \Rightarrow \sim p$
3.  $p \vee q$
4.  $(\sim p) \Rightarrow q$

Question Type : **MCQ**Question ID : **41652913505**Option 1 ID : **41652952800**Option 2 ID : **41652952799**Option 3 ID : **41652952801**Option 4 ID : **41652952798**Status : **Answered**

Chosen Option : 3

**Q.12** A circle touching the  $x$ -axis at  $(3, 0)$  and making an intercept of length 8 on the  $y$ -axis passes through the point :

Options

1.  $(3, 10)$
2.  $(3, 5)$
3.  $(2, 3)$
4.  $(1, 5)$

Question Type : **MCQ**Question ID : **41652913495**Option 1 ID : **41652952758**Option 2 ID : **41652952759**Option 3 ID : **41652952761**Option 4 ID : **41652952760**Status : **Not Answered**

Chosen Option : --

**Q.13**

If  ${}^{20}C_1 + (2^2) {}^{20}C_2 + (3^2) {}^{20}C_3 + \dots + (20^2) {}^{20}C_{20} = A(2^\beta)$ , then the ordered pair  $(A, \beta)$  is equal to :

- Options
1. (420, 19)
  2. (420, 18)
  3. (380, 18)
  4. (380, 19)

Question Type : **MCQ**

Question ID : **41652913484**

Option 1 ID : **41652952717**

Option 2 ID : **41652952716**

Option 3 ID : **41652952714**

Option 4 ID : **41652952715**

Status : **Not Answered**

Chosen Option : --

Q.14 A value of  $\theta \in (0, \pi/3)$ , for which

$$\begin{vmatrix} 1 + \cos^2\theta & \sin^2\theta & 4 \cos 6\theta \\ \cos^2\theta & 1 + \sin^2\theta & 4 \cos 6\theta \\ \cos^2\theta & \sin^2\theta & 1 + 4 \cos 6\theta \end{vmatrix} = 0, \text{ is :}$$

- Options
1.  $\frac{\pi}{9}$
  2.  $\frac{\pi}{18}$
  3.  $\frac{7\pi}{24}$
  4.  $\frac{7\pi}{36}$

Question Type : **MCQ**

Question ID : **41652913479**

Option 1 ID : **41652952695**

Option 2 ID : **41652952694**

Option 3 ID : **41652952696**

Option 4 ID : **41652952697**

Status : **Answered**

Chosen Option : **4**

Q.15 The equation of a common tangent to the curves,  $y^2 = 16x$  and  $xy = -4$ , is :

- Options
1.  $x - y + 4 = 0$
  2.  $x + y + 4 = 0$
  3.  $x - 2y + 16 = 0$
  4.  $2x - y + 2 = 0$

Question Type : **MCQ**

Question ID : 41652913496

Option 1 ID : 41652952765

Option 2 ID : 41652952763

Option 3 ID : 41652952762

Option 4 ID : 41652952764

Status : Answered

Chosen Option : 3

Q.16 Let  $z \in \mathbb{C}$  with  $\text{Im}(z) = 10$  and it satisfies

$$\frac{2z - n}{2z + n} = 2i - 1 \text{ for some natural number } n.$$

n. Then :

- Options
1.  $n = 20$  and  $\text{Re}(z) = -10$
  2.  $n = 40$  and  $\text{Re}(z) = 10$
  3.  $n = 40$  and  $\text{Re}(z) = -10$
  4.  $n = 20$  and  $\text{Re}(z) = 10$

Question Type : MCQ

Question ID : 41652913477

Option 1 ID : 41652952687

Option 2 ID : 41652952688

Option 3 ID : 41652952689

Option 4 ID : 41652952686

Status : Not Answered

Chosen Option : --

Q.17 A triangle has a vertex at  $(1, 2)$  and the mid points of the two sides through it are  $(-1, 1)$  and  $(2, 3)$ . Then the centroid of this triangle is :

- Options
1.  $\left(1, \frac{7}{3}\right)$
  2.  $\left(\frac{1}{3}, 2\right)$
  3.  $\left(\frac{1}{3}, 1\right)$
  4.  $\left(\frac{1}{3}, \frac{5}{3}\right)$

Question Type : MCQ

Question ID : 41652913494

Option 1 ID : 41652952756

Option 2 ID : 41652952755

Option 3 ID : 41652952754

Option 4 ID : 41652952757

Status : Not Answered

Chosen Option : --

Q.18

If  $a_1, a_2, a_3, \dots$  are in A.P. such that  $a_1 + a_7 + a_{16} = 40$ , then the sum of the first 15 terms of this A.P. is :

- Options
1. 200
  2. 280
  3. 120
  4. 150

Question Type : **MCQ**

Question ID : **41652913483**

Option 1 ID : **41652952713**

Option 2 ID : **41652952712**

Option 3 ID : **41652952711**

Option 4 ID : **41652952710**

Status : **Answered**

Chosen Option : **3**

**Q.19** If  $[x]$  denotes the greatest integer  $\leq x$ , then the system of linear equations  $[\sin\theta]x + [-\cos\theta]y = 0$   
 $[\cot\theta]x + y = 0$

Options have infinitely many solutions if

1.  $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right)$  and has a unique

solution if  $\theta \in \left(\pi, \frac{7\pi}{6}\right)$ .

has a unique solution if

2.  $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right) \cup \left(\pi, \frac{7\pi}{6}\right)$ .

has a unique solution if  $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right)$

3. and have infinitely many solutions if

$\theta \in \left(\pi, \frac{7\pi}{6}\right)$ .

have infinitely many solutions if

4.  $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right) \cup \left(\pi, \frac{7\pi}{6}\right)$ .

Question Type : **MCQ**

Question ID : **41652913480**

Option 1 ID : **41652952701**

Option 2 ID : **41652952698**

Option 3 ID : **41652952700**

Option 4 ID : **41652952699**

Status : **Not Answered**

Chosen Option : **--**

**Q.20** A straight line L at a distance of 4 units from the origin makes positive intercepts on the coordinate axes and the perpendicular from the origin to this line makes an angle of  $60^\circ$  with the line  $x + y = 0$ . Then an equation of the line L is :

- Options**
1.  $x + \sqrt{3}y = 8$
  2.  $(\sqrt{3} + 1)x + (\sqrt{3} - 1)y = 8\sqrt{2}$
  3.  $\sqrt{3}x + y = 8$
  4.  $(\sqrt{3} - 1)x + (\sqrt{3} + 1)y = 8\sqrt{2}$

Question Type : **MCQ**

Question ID : **41652913493**

Option 1 ID : **41652952753**

Option 2 ID : **41652952750**

Option 3 ID : **41652952752**

Option 4 ID : **41652952751**

Status : **Not Answered**

Chosen Option : --

**Q.21** Let  $f(x) = 5 - |x - 2|$  and  $g(x) = |x + 1|$ ,  $x \in \mathbb{R}$ . If  $f(x)$  attains maximum value at  $\alpha$  and  $g(x)$  attains minimum value at  $\beta$ , then

$\lim_{x \rightarrow -\alpha\beta} \frac{(x - 1)(x^2 - 5x + 6)}{x^2 - 6x + 8}$  is equal to :

- Options**
1.  $1/2$
  2.  $-3/2$
  3.  $-1/2$
  4.  $3/2$

Question Type : **MCQ**

Question ID : **41652913488**

Option 1 ID : **41652952730**

Option 2 ID : **41652952733**

Option 3 ID : **41652952731**

Option 4 ID : **41652952732**

Status : **Not Answered**

Chosen Option : --

**Q.22** Let  $\alpha \in \mathbb{R}$  and the three vectors

$$\vec{a} = \alpha \hat{i} + \hat{j} + 3\hat{k}, \quad \vec{b} = 2\hat{i} + \hat{j} - \alpha\hat{k}$$

and  $\vec{c} = \alpha\hat{i} - 2\hat{j} + 3\hat{k}$ . Then the set

$$S = \{\alpha : \vec{a}, \vec{b} \text{ and } \vec{c} \text{ are coplanar}\}$$

- Options**
1. is singleton

2. is empty
3. contains exactly two positive numbers
4. contains exactly two numbers only one of which is positive

Question Type : **MCQ**  
Question ID : **41652913500**  
Option 1 ID : **41652952779**  
Option 2 ID : **41652952778**  
Option 3 ID : **41652952780**  
Option 4 ID : **41652952781**  
Status : **Answered**  
Chosen Option : **2**

**Q.23** A person throws two fair dice. He wins Rs. 15 for throwing a doublet (same numbers on the two dice), wins Rs. 12 when the throw results in the sum of 9, and loses Rs. 6 for any other outcome on the throw. Then the expected gain/loss (in Rs.) of the person is :

- Options
1.  $\frac{1}{2}$  gain
  2.  $\frac{1}{4}$  loss
  3.  $\frac{1}{2}$  loss
  4. 2 gain

Question Type : **MCQ**  
Question ID : **41652913501**  
Option 1 ID : **41652952784**  
Option 2 ID : **41652952785**  
Option 3 ID : **41652952783**  
Option 4 ID : **41652952782**  
Status : **Answered**  
Chosen Option : **3**

**Q.24** The tangents to the curve  $y = (x - 2)^2 - 1$  at its points of intersection with the line  $x - y = 3$ , intersect at the point :

- Options
1.  $\left(\frac{5}{2}, 1\right)$
  2.  $\left(-\frac{5}{2}, -1\right)$
  3.  $\left(\frac{5}{2}, -1\right)$



4.  $\left(-\frac{5}{2}, 1\right)$

Question Type : **MCQ**

Question ID : **41652913487**

Option 1 ID : **41652952728**

Option 2 ID : **41652952727**

Option 3 ID : **41652952726**

Option 4 ID : **41652952729**

Status : **Not Answered**

Chosen Option : --

**Q.25** If  $\alpha$ ,  $\beta$  and  $\gamma$  are three consecutive terms of a non-constant G.P. such that the equations  $\alpha x^2 + 2\beta x + \gamma = 0$  and  $x^2 + x - 1 = 0$  have a common root, then  $\alpha(\beta + \gamma)$  is equal to :

Options 1. 0

2.  $\alpha\beta$

3.  $\alpha\gamma$

4.  $\beta\gamma$

Question Type : **MCQ**

Question ID : **41652913478**

Option 1 ID : **41652952693**

Option 2 ID : **41652952691**

Option 3 ID : **41652952692**

Option 4 ID : **41652952690**

Status : **Not Answered**

Chosen Option : --

**Q.26** Let A, B and C be sets such that  $\phi \neq A \cap B \subseteq C$ . Then which of the following statements is not true ?

Options 1.  $B \cap C \neq \phi$

2. If  $(A - B) \subseteq C$ , then  $A \subseteq C$

3.  $(C \cup A) \cap (C \cup B) = C$

4. If  $(A - C) \subseteq B$ , then  $A \subseteq B$

Question Type : **MCQ**

Question ID : **41652913476**

Option 1 ID : **41652952682**

Option 2 ID : **41652952684**

Option 3 ID : **41652952683**

Option 4 ID : **41652952685**

Status : **Not Answered**

Chosen Option : --

**Q.27** The general solution of the differential equation  $(y^2 - x^3) dx - xy dy = 0$  ( $x \neq 0$ ) is :  
(where c is a constant of integration)

Options

1.  $y^2 - 2x^2 + cx^3 = 0$
2.  $y^2 + 2x^3 + cx^2 = 0$
3.  $y^2 + 2x^2 + cx^3 = 0$
4.  $y^2 - 2x^3 + cx^2 = 0$

Question Type : **MCQ**Question ID : **41652913492**Option 1 ID : **41652952749**Option 2 ID : **41652952748**Option 3 ID : **41652952746**Option 4 ID : **41652952747**Status : **Not Answered**

Chosen Option : --

**Q.28** If the area (in sq. units) bounded by the parabola  $y^2 = 4\lambda x$  and the line  $y = \lambda x$ ,  $\lambda > 0$ , is  $\frac{1}{9}$ , then  $\lambda$  is equal to :

- Options
1.  $2\sqrt{6}$
  2. 48
  3. 24
  4.  $4\sqrt{3}$

Question Type : **MCQ**Question ID : **41652913491**Option 1 ID : **41652952743**Option 2 ID : **41652952745**Option 3 ID : **41652952744**Option 4 ID : **41652952742**Status : **Answered**

Chosen Option : 1

**Q.29** The length of the perpendicular drawn from the point  $(2, 1, 4)$  to the plane containing the lines

$$\vec{r} = (\hat{i} + \hat{j}) + \lambda(\hat{i} + 2\hat{j} - \hat{k}) \quad \text{and}$$

$$\vec{r} = (\hat{i} + \hat{j}) + \mu(-\hat{i} + \hat{j} - 2\hat{k}) \quad \text{is:}$$

- Options
1. 3
  2.  $\frac{1}{3}$
  3.  $\sqrt{3}$
  4.  $\frac{1}{\sqrt{3}}$

Question Type : **MCQ**Question ID : **41652913499**Option 1 ID : **41652952774**

Option 2 ID : **41652952776**  
Option 3 ID : **41652952775**  
Option 4 ID : **41652952777**  
Status : **Not Answered**  
Chosen Option : --

**Q.30** The term independent of  $x$  in the expansion

of  $\left(\frac{1}{60} - \frac{x^8}{81}\right) \cdot \left(2x^2 - \frac{3}{x^2}\right)^6$  is equal to :

- Options**
1. -72
  2. 36
  3. -36
  4. -108

Question Type : **MCQ**  
Question ID : **41652913482**  
Option 1 ID : **41652952707**  
Option 2 ID : **41652952709**  
Option 3 ID : **41652952708**  
Option 4 ID : **41652952706**  
Status : **Not Answered**  
Chosen Option : --