

[CHEMISTRY]





1. A process has $\Delta H = 200 \text{ Jmol}^{-1}$ and $\Delta S = 40 \text{ JK}^{-1}\text{mol}^{-1}$. Out of the values given below, choose the minimum temperature above which the process will be spontaneous:
 (A) 12 K (B) 20 K (C) 4 K (D) 5 K

Sol. D

$$\Delta G = \Delta H - T\Delta S$$

$$T = \frac{\Delta H}{\Delta S} = \frac{200}{40} = 5 \text{ K}$$

2. Consider the following reduction processes:



The reducing power of the metals increases in the order :

- (A) $\text{Ca} < \text{Mg} < \text{Zn} < \text{Ni}$ (B) $\text{Ni} < \text{Zn} < \text{Mg} < \text{Ca}$ (C) $\text{Ca} < \text{Zn} < \text{Mg} < \text{Ni}$ (D) $\text{Zn} < \text{Mg} < \text{Ni} < \text{Ca}$

Sol. B

Higher the oxidation potential better will be reducing power.

3. Two pi and half sigma bonds are present in :

- (A) O_2^+ (B) N_2 (C) N_2^+ (D) O_2

Sol. C




$$\text{N}_2^+ \Rightarrow \text{B.O.} = 2.5 \Rightarrow [\pi\text{-Bond} = 2 \text{ \& } \sigma\text{-bond} = \frac{1}{2}]$$

$$\text{N}_2 \Rightarrow \text{B.O.} = 3.0 \Rightarrow [\pi\text{-Bond} = 2 \text{ \& } \sigma\text{-Bond} = 1]$$

$$\text{O}_2^+ \Rightarrow \text{B.O.} = 2.5 \Rightarrow [\pi\text{-Bond} = 1.5 \text{ \& } \sigma\text{-Bond} = 1]$$

$$\text{O}_2 \Rightarrow \text{B.O.} = 2 \Rightarrow [\pi\text{-Bond} = 1 \text{ \& } \sigma\text{-bond} = 1]$$

4. The decreasing order of ease of alkaline hydrolysis for the following esters is






- (A) $\text{IV} > \text{II} > \text{III} > \text{I}$ (B) $\text{III} > \text{II} > \text{I} > \text{IV}$ (C) $\text{III} > \text{II} > \text{IV} > \text{I}$ (D) $\text{II} > \text{III} > \text{I} > \text{IV}$

Sol. B

More is the electrophilic character of carbonyl group of ester faster is the alkaline hydrolysis

5. The values of K_p/K_c for the following reactions at 300 K are, respectively: (At 300 K, $RT = 24.62 \text{ dm}^3 \text{ atm mol}^{-1}$)




- (A) $1, 24.62 \text{ dm}^3 \text{ atm mol}^{-1}, 1.62 \times 10^{-3} \text{ dm}^{-2} \text{ atm}^{-2} \text{ mol}^2$


- (B) $1, 4.1 \times 10^{-2} \text{ dm}^{-3} \text{ atm}^{-1} \text{ mol}, 606 \text{ dm}^6 \text{ atm}^2 \text{ mol}^{-2}$

- (C) $1, 24.62 \text{ dm}^3 \text{ atm mol}^{-1}, 606.0 \text{ dm}^6 \text{ atm}^2 \text{ mol}^{-2}$


- (D) $24.62 \text{ dm}^3 \text{ atm mol}^{-1}, 606.0 \text{ dm}^6 \text{ atm}^2 \text{ mol}^{-2}, 1.65 \times 10^{-3} \text{ dm}^{-6} \text{ atm}^{-2} \text{ mol}^2$

Sol. A





$$\frac{k_p}{k_c} = (\text{RT})^{\Delta n_g} = (\text{RT})^0 = 1$$













$$\frac{k_p}{k_c} = (\text{RT})^1 = 24.62$$



$$\frac{k_p}{k_c} = (\text{RT})^{-2} = \frac{1}{(\text{RT})^2} = 1.65 \times 10^{-3}$$

6. Hall-Haroult's process is given by:**Sol. C**

In Hall-Heroult's process is given by

**7.** The major product formed in the reaction given below will be:**Sol. Bonus****8.** The metal used for making X-ray tube window is:

(A) Mg

(B) Na





(C) Ca

(D) Be

Sol. D

"Be" Metal is used in x-ray window is due to transparent to x-rays.

9. Which of the graphs shown below does not represent the relationship between incident light and the electron ejected from metal surface?



Sol. A






$$E = W + \frac{1}{2} mv^2$$

$$\text{K.E.} = h\nu - 4v_0$$

$$\text{K.E.} = h\nu + (-h\nu_0)$$

$$y = mx + C$$

10. The major product 'X' formed in the following reaction is:



Sol. D

NaBH_4 Reduced Ketone to 2° alcohol. Inert to Ester & $\text{C}=\text{C}$

11. Liquids A and B form an ideal solution in the entire composition range. At 350 K, the vapor pressures of pure A and pure B are 7×10^3 Pa and 12×10^3 Pa, respectively. The composition of the vapor in equilibrium with a solution containing 40 mole percent of A at this temperature is:
 (A) $x_A = 0.28; x_B = 0.72$ (B) $x_A = 0.76; x_B = 0.24$ (C) $x_A = 0.37; x_B = 0.63$ (D) $x_A = 0.4; x_B = 0.6$

Sol. **A**


$$y_A = \frac{P_A}{P_{\text{Total}}} = \frac{P_A^\circ x_A}{P_A^\circ x_A + P_B^\circ x_B}$$

$$= \frac{7 \times 10^3 \times 0.4}{7 \times 10^3 \times 0.4 + 12 \times 10^3 \times 0.6}$$

$$= \frac{2.8}{10} = 0.28$$

$$y_B = 0.72$$



12. Which dicarboxylic acid in presence of a dehydrating agent is least reactive to give an anhydride?




Sol. **B**

Adipic acid $\text{CO}_2\text{H}-(\text{CH}_2)_4-\text{CO}_2\text{H} \xrightarrow{\text{dehydrating agent}}$ 7 membered cyclic anhydride (Very unstable)

13. The major product of the following reaction is:



Sol. B




14. If dichloromethane (DCM) and water (H₂O) are used for differential extraction, which one of the following statements is correct ?

- (A) DCM and H₂O will be miscible clearly
 (B) DCM and H₂O will make turbid/ colloidal mixture
 (C) DCM and H₂O would stay as lower and upper layer respectively in the S.F.
 (D) DCM and H₂O would stay as upper and lower layer respectively in the separating funnel (S.F.)

Sol. C



15. The correct structure of product 'P' in the following reaction is :




- (A) CC(=O)NCC(=O)NCC(=O)NCC(=O)O
- (B) CC(=O)NCC(=O)NCC(=O)NCC(=O)OCC(=O)C
- (C) CC(=O)NCC(=O)NCC(=O)NCC(=O)OCC(=O)C
- (D) CC(=O)NCC(=O)NCC(=O)NCC(=O)O

Sol. D

Asn-Ser is dipeptide having following structure



P is




- 16.** Water filled in two glasses A and B have BOD values of 10 and 20, respectively. The correct statement regarding them, is :
- (A) A is suitable for drinking, whereas B is not.
 (B) A is more polluted than B.
 (C) B is more polluted than A.
 (D) Both A and B are suitable for drinking.

Sol. C

Two glasses "A" and "B" have BOD values 10 and "20", respectively. Hence glasses "B" is more polluted than glasses "A".


- 17.** Which hydrogen in compound (E) is easily replaceable during bromination reaction in presence of light ?



- (A) δ -hydrogen (B) β -hydrogen
 (C) γ -hydrogen (D) α -hydrogen

Sol. CIntermediate formed free radical $\text{CH}_3-\dot{\text{C}}\text{H}-\text{CH}=\text{CH}_2$ stable due to Resonance & hyperconjugation

- 18.** Consider the given plots for a reaction obeying Arrhenius equation ($0^\circ\text{C} < T < 300^\circ\text{C}$) : (k and E_a are rate constant and activation energy, respectively)








Choose the correct option :

- (A) I is right but II is wrong (B) Both I and II are correct
 (C) Both I and II are wrong (D) I is wrong but II is right


Sol. BOn increasing E_a , K decreases

19. The major product of the following reaction is :



Sol. **D**

Dehydrohalogenation at β – Position



20. The electronegativity of aluminium is similar to :
 (A) Boron (B) Lithium
 (C) Beryllium (D) Carbon

Sol. **C**

E.N. of Al = (1.5) \cong Be (1.5)

21. The chemical nature of hydrogen peroxide is :
 (A) Oxidising and reducing agent in acidic medium, but not in basic medium.
 (B) Oxidising and reducing agent in both acidic and basic medium.
 (C) Reducing agent in basic medium, but not in acidic medium
 (D) Oxidising agent in acidic medium, but not in basic medium

Sol. **B**

H_2O_2 act as oxidising agent and reducing agent in acidic medium as well as basic medium.

H_2O_2 Act as oxidant :-

$\text{H}_2\text{O}_2 + 2\text{H}^+ + 2\text{e}^- \rightarrow 2\text{H}_2\text{O}$ (In acidic medium)

$\text{H}_2\text{O}_2 + 2\text{e}^- \rightarrow 2\text{OH}^-$ (In basic medium)

H_2O_2 Act as reductant :-

$\text{H}_2\text{O}_2 \rightarrow 2\text{H}^+ + \text{O}_2 + 2\text{e}^-$ (In acidic medium)

$\text{H}_2\text{O}_2 + 2\text{OH}^- \rightarrow 2\text{H}_2\text{O} + \text{O}_2 + 2\text{e}^-$ (In basic medium)


22. The effect of lanthanoid contraction in the lanthanoid series of elements by and large means :
 (A) increase in atomic radii and decrease in ionic radii
 (B) increase in both atomic and ionic radii
 (C) decrease in both atomic and ionic radii
 (D) decrease in atomic radii and increase in ionic radii

Sol. **C**

Due to Lanthanoid contraction both atomic radii and ionic radii decreases gradually in the lanthanoid series.

23. The type of hybridisation and number of lone pair(s) of electrons of Xe in XeOF_4 , respectively, are :
 (A) sp^3d and 1 (B) sp^3d and 2
 (C) sp^3d^2 and 2 (D) sp^3d^2 and 1


Sol. **D**



$\text{sp}^3\text{d}^2 \Rightarrow [5\text{s-bond} + 1 \text{ l.p.}]$

- 24.** A mixture of 100 m mol of Ca(OH)_2 and 2 g of sodium sulphate was dissolved in water and the volume was made up to 100 mL. The mass of calcium sulphate formed and the concentration of OH^- in resulting solution, respectively are (Molar mass of Ca(OH)_2 , Na_2SO_4 and CaSO_4 are 74, 143 and 136 g mol⁻¹, respectively ; K_{sp} of Ca(OH)_2 is 5.5×10^{-6})
- (A) 1.9 g, 0.14 mol L⁻¹ (B) 13.6 g, 0.28 mol L⁻¹
 (C) 1.9 g, 0.28 mol L⁻¹ (D) 13.6 g, 0.14 mol L⁻¹

Sol. C




$$W_{\text{CaSO}_4} = 14 \times 10^{-3} \times 136 = 1.9 \text{ gm}$$

$$[\text{OH}^-] = \frac{28}{100} = 0.28 \text{ M}$$

- 25.** The total number of isomers for a square planar complex $[\text{M}(\text{F})(\text{Cl})(\text{SCN})(\text{NO}_2)]$ is :
- (A) 16 (B) 4
(C) 8 (D) 12

Sol. D

The total number of isomers for a square planar complex $[\text{M}(\text{F})(\text{Cl})(\text{SCN})(\text{NO}_2)]$ is 12.



- 26.** Wilkinson catalyst is :
- (A) $[(\text{Ph}_3\text{P})_3\text{RhCl}]$ (Et = C_2H_5)
 (B) $[(\text{Ph}_3\text{P})_3\text{IrCl}]$
 (C) $[(\text{Et}_3\text{P})_3\text{RhCl}]$
 (D) $[(\text{Et}_3\text{P})_3\text{IrCl}]$

Sol. A

Wilkinson catalyst is $[(\text{ph}_3\text{P})_3\text{RhCl}]$

- 27.** Which of the following is not an example of heterogeneous catalytic reaction ?
- (A) Combustion of coal (B) Hydrogenation of vegetable oils
 (C) Ostwald's process (D) Haber's process


Sol. A

Then is no catalyst is required for combustion of coal.


- 28.** Which primitive unit cell has unequal edge lengths ($a \neq b \neq c$) and all axial angles different from 90° ?
- (A) Triclinic (B) Hexagonal
 (C) Tetragonal (D) Monoclinic

Sol. A

In Triclinic unit cell

 $a \neq b \neq c$ & $\alpha \neq \beta \neq \gamma \neq 90^\circ$ **29.** The increasing order of the pKa values of the following compounds is :(A) $B < C < D < A$ (B) $D < A < C < B$ (C) $B < C < A < D$ (D) $C < B < A < D$ **Sol. C**

Acidic strength is inversely proportional to pKa.

**30.** The total number of isotopes of hydrogen and number of radioactive isotopes among them, respectively, are :

(A) 3 and 2

(B) 3 and 1

(C) 2 and 1

(D) 2 and 0

Sol. B

Total number of isotopes of hydrogen is 3

$$\Rightarrow {}^1_1\text{H}({}^2_1\text{H} \text{ or } {}^2_1\text{D})({}^3_1\text{H} \text{ or } {}^3_1\text{T})$$

and only ${}^3_1\text{H}$ or ${}^3_1\text{T}$ is an Radioactive element.