Preventing Baby Bottle Tooth Decay

A step-by-step guide for parents to protect their child's smile.

Based on Baby Bottle Tooth Decay by Cool Mountain Kids Pediatric Dentistry

Step 1: Understand Baby Bottle Tooth Decay

Baby bottle tooth decay occurs when a child's teeth are frequently exposed to sugary liquids, especially during sleep when saliva production is low. Bacteria feed on sugars, producing acids that attack tooth enamel.

- Even natural sugars in breast milk and formula can contribute if pooled around teeth during sleep.
- Decreased saliva production during sleep reduces the mouth's natural rinsing action.

↓

Step 2: Recognize Early Warning Signs

Look for white spots or chalky patches on your child's teeth, particularly along the gum line of the upper front teeth. These can progress to brown or black spots and cavities.

- Some children may experience sensitivity or discomfort.
- Regular dental checkups are crucial for early detection.

↓

Step 3: Identify Risk Factors

Be aware of common scenarios that increase risk, such as bedtime bottles with sugary liquids, prolonged bottle use beyond 12-15 months, and frequent snacking on sugary foods.

- Sippy cup use with anything other than water throughout the day can also pose a risk.
- Sharing utensils or pacifiers can transfer cavity-causing bacteria.

↓

Step 4: Start Oral Care Early

Begin cleaning your baby's gums with a damp cloth after feedings even before teeth emerge. Once the first tooth appears, use a soft-bristled brush with a smear of fluoride toothpaste twice daily.

- This helps remove sugars and bacteria and gets your baby accustomed to mouth care.

↓

Step 5: Modify Feeding and Drinking Habits

Avoid bedtime bottles with sugary liquids. Offer water if comfort is needed. Transition to a cup by age one and limit juice and sweetened beverages to mealtimes.

- Offer only water in bottles at bedtime.
- Limit juice to four ounces of 100% fruit juice with meals.

↓

Step 6: Promote Healthy Snacking

Choose nutritious snacks like cheese, vegetables, and fruits over sticky, sugary options that can cling to teeth and promote decay.

↓

Step 7: Schedule Regular Dental Visits

Schedule your child's first dental visit around their first birthday or within six months of their first tooth appearing. These visits establish a dental home and provide personalized advice.

- Ensure adequate fluoride exposure, as recommended by your dentist.

Resources & Further Reading

Related guides for this process:

- Action Sheet: Preventing Baby Bottle Tooth Decay

https://storage.googleapis.com/pediatric-dentistry-checklists-20260707-8878a5/pediatric-dentistry/Action%20Sheet/baby-bottle-tooth-decay-action-sheet.pdf

- Baby Bottle Tooth Decay Prevention Checklist

https://storage.googleapis.com/pediatric-dentistry-checklists-20260707-a10c96/pediatric-dentistry/Checklist/baby-bottle-tooth-decay-prevention-checklist.pdf

---

Cool Mountain Kids Pediatric Dentistry | 6160 Tutt Blvd #140, Colorado Springs, CO 80923
(719) 550-1010 | www.coolmountainkids.com
Protect your child's smile by understanding and preventing baby bottle tooth decay.