


FORMAL COMMENT

Response to: "Questioning the evidence for BCI-based communication in the complete locked-in state"

Ujwal Chaudhary^{1,2*}, Sudhir Pathak³, Niels Birbaumer^{1,2*}

1 Institute of Medical Psychology and Behavioral Neurobiology, Eberhard-Karls University of Tuebingen, Tuebingen, Germany, **2** Wyss Center for Bio and Neuroengineering, Geneva, Switzerland, **3** Learning Research and Development Center (LRDC), University of Pittsburgh, Pittsburgh, United States of America

* chaudharyujwal@gmail.com (UC); niels.birbaumer@uni-tuebingen.de (NB)



OPEN ACCESS

Citation: Chaudhary U, Pathak S, Birbaumer N (2019) Response to: "Questioning the evidence for BCI-based communication in the complete locked-in state". *PLoS Biol* 17(4): e3000063. <https://doi.org/10.1371/journal.pbio.3000063>

Academic Editor: Ulrich Dirnagl, Charite Universitätsmedizin Berlin, GERMANY

Received: June 27, 2018

Accepted: March 5, 2019

Published: April 8, 2019

Copyright: © 2019 Chaudhary et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Deutsche Forschungsgemeinschaft (DFG-Koselleck Grant, BI 195/69-1 and DFG, BI 195/77-1), German Ministry of Education and Research (BMBF) 16SV7701 CoMiCon, Baden-Württemberg Stiftung, LUMINOUS-H2020-FETOPEN-2014-2015-RIA (686764), and Wyss Center for Bio and Neuroengineering, Geneva.

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: ALS, amyotrophic lateral sclerosis; CLIS, completely locked-in state; CSP, common spatial pattern; EEG, electroencephalogram; fNIRS,

Patients in completely locked-in state (CLIS) have no means of communication and present a highly challenging and daunting problem for the neuroscientist [1–3]. Until today, few groups have attempted to solve this problem, and only some have reported success in advancing the goal of providing a means of communication to patients in CLIS [4–7]. In his commentary, Dr. Spüler raises doubts about all the research efforts towards this goal but primarily about the results published in 2017 by Chaudhary and colleagues. Dr. Spüler bases the commentary on 2 main calculations:

1. Absence of hemodynamic differences between “yes” and “no” thinking and
2. Chance-level classification across all the sessions in the 4 published cases with CLIS.

In this commentary, we address the issues raised by Dr. Spüler.

1. Absence of hemodynamic differences between “yes” and “no” thinking

In his commentary, Dr. Spüler claims that, in the paper by Chaudhary and colleagues [6], the change in the concentrations of oxy-hemoglobin (O₂Hb) acquired from 20 different functional near-infrared spectroscopy (fNIRS) channels were averaged, and then further averaging was performed across trials and sessions. Chaudhary and colleagues [6] presented the averaged change in relative concentration of O₂Hb separately for each of the 20 channels used during the study, as shown in their Fig 1 [6]. In none of the fNIRS literature published to date have fNIRS channels placed across such disparate regions been averaged [8–12]. The reason behind not averaging the channels is the fact that different channels represent metabolic information from the respective underlying brain region. In Chaudhary and colleagues’ paper [6], therefore, first signal acquired across different trials—i.e., “yes” and “no” thinking—were averaged separately for the different channels and were then averaged across sessions as shown in Fig 1 of Chaudhary and colleagues [6]. Fig 1 of Chaudhary and colleagues’ paper shows the averaged relative change in O₂Hb from all the 20 channels; if all 20 channels were averaged, then we would have had just 1 time-series of relative change in O₂Hb and not 20 different time-series of relative change in O₂Hb, each corresponding to a channel, as depicted by Chaudhary and colleagues. To further elucidate the difference in hemodynamic response between “yes” and “no” thinking, general linear model (GLM) analysis was performed as shown in *S1 Text*. Dr. Spüler’s claim of a lack of difference between the 2 response categories “yes” and “no” thinking is thus unfounded and not comparable to that of Chaudhary and colleagues. As reported by

functional near-infrared spectroscopy; GLM, general linear model; O₂Hb, oxy-hemoglobin.

Chaudhary and colleagues, channels were treated separately for classification and model building for online feedback session as written on page 18 and 19 of Chaudhary and colleagues' paper. According to Chaudhary and colleagues (page 18), "The mean of relative change in O₂Hb across each channel was used as a feature to train the SVM model through a 5-fold cross-validation procedure." On page 19, Chaudhary and colleagues further state that, "During an online feedback session, fNIRS data acquired online corresponding to each ISI was processed to obtain the relative change in O₂Hb, as described above, across all the channels. The mean of the relative change in O₂Hb across all the channels was used as test feature to map onto model space."

2. Chance-level classification across all the sessions

Spüler raises doubts about the classification results based on the method he employed to calculate the offline classification accuracy of each session. It is well known in the machine learning literature that application of different machine learning algorithms and features results in different outcomes, as is obvious from the result presented by Dr. Spüler. We can argue on the




Fig 1. Bar graph of offline classification accuracy results obtained from the sessions performed by the patient B and published by Chaudhary and colleagues [6]. (A) The offline classification accuracy of sessions performed by patient B using the method reported by Chaudhary and colleagues, i.e., the mean O₂Hb response was used as input feature for the linear support vector machine classifier. (B) The offline classification accuracy of sessions performed by patient B using the method suggested by Martin Spüler. (C) The offline classification accuracy of sessions performed by patient B using the CSP for feature extraction and using linear SVM classifier. CSP, common spatial pattern; O₂Hb, oxy-hemoglobin; SVM, support vector machine.

<https://doi.org/10.1371/journal.pbio.3000063.g001>

method that can and should be used for classification, but that does not invalidate the results presented by Chaudhary and colleagues [6]. It has also been argued that the sessions should be combined randomly to build a model; we argue that such a method might be valid for stable and invariant data but might be completely misleading for patients in CLIS. As reported by Chaudhary and colleagues, physiological and psychological states (arousal and attention) of the patients should be considered when performing the analysis because these patients spend long periods of the waking day sleeping and dozing, hence only sessions during which patients were vigilant should be used to build classification models to provide feedback to the patients. Therefore, combining sessions randomly to build the classification model would include time periods of an unresponsive state of the particular patient. The correlations of electroencephalogram (EEG) slow activity with performance, we reported in the original paper, underscore this point and demonstrate that such a relationship with lack of arousal and poor performance exists. These issues were thoroughly discussed in the section entitled “Slow EEG rhythms’ relationship with fNIRS classification accuracy” on page 8 as well as the discussion section on page 12 titled “BCI performance and attention-vigilance” of Chaudhary and colleagues’ paper. Thus, Dr. Spüler’s failure to replicate our results may originate from such a random treatment and random selection of sessions to calculate the classification precisions of patients’ mental answers. Assuming that we continue a session on a day when the patient is unresponsive, the EEG shows dominant slow-wave activity of 1 to 2 Hz, and we build the model to perform online feedback session—in such a case, the patient will receive wrong feedback over an extended time period not determined by his or her actual performance but by an episode of deep sleep and unresponsiveness.

Here, we present the classification result using the data from patient B from 3 different analysis methods, as follows: (a) the method described by Chaudhary and colleagues (shown in Fig 1A), (b) the method proposed by Martin Spüler (shown in Fig 1B), and (c) using common spatial pattern (CSP) [13] (details shown in S1 Text) to extract the feature and then performing classification (shown in Fig 1C).

Thus, based on the result presented in this section, it can be seen that we can implement different methods, which can have both negative and positive effect on classification result. The goal of our ongoing research is to find and apply the best strategy to improve the classification accuracy to increase the overall communication rate with patients [14], which we have stated clearly in the discussion section of Chaudhary and colleagues’ paper.

Slowing of EEG and consciousness

Slowing of EEG frequencies as found in many CLIS patients with amyotrophic lateral sclerosis (ALS) and other neurological conditions can have many reasons, such as the simple fact that these patients have to use artificial ventilation, which bypasses breathing through the nose. The neuronal epithelium involved in olfaction synchronizes high frequency (beta and gamma) in the entire brain of mammals [15]. Thus, slowing of frequencies may just indicate a side effect of bypassing airflow from nose to the tracheostoma and not any “impaired cognitive abilities.” Martin Spüler continues, in his discussion, to question the cognitive intactness of CLIS patients, claiming that alpha wave indicates “consciousness” and suggesting implicitly that their absence points to a lack of conscious processing. Such a claim is not only wrong but also medically dangerous because of attribution of conscious experience to unconscious patients and frequent misdiagnoses. In addition, alpha waves are not a unitary concept but a general term for very heterogeneous physiological phenomena: alpha of the sensorimotor cortex, for example—also called somatosensory alpha or sensorimotor rhythm [16] or mu rhythm—indicates quiescence of the motor system, with no relationship to any form of consciousness.

Auditory alpha exclusively originating from the central auditory cortex is interpreted as an inhibitory state of the central auditory analysis—again, no relationship with consciousness. An extensive literature [17] on visual, occipital alpha waves has shown that their presence depends heavily on coupling and uncoupling of the oculomotor system, which is a purely motor phenomenon independent of the conscious state of the organism. Comprehensive discussion and information of the physiological basis of alpha rhythm can be found in Andersen and Andersen [18] and Klimesch and colleagues [19]. Moreover, patients in Chaudhary and colleagues' publication participated in some of these testing procedures [20], and all showed an electrophysiological correlate of cognitive processing (not presented in Chaudhary and colleagues' 2017 paper, as that was not the goal of the paper).

Missing data link

<https://doi.org/10.5281/zenodo.1419151>

We committed an error in uploading the data of patient F: we uploaded the data of 28 sessions, although Chaudhary and colleagues contain the results of 58 sessions from patient F. Herein, we are uploading the remaining data from this patient, whose results are already included in the Chaudhary and colleagues' paper.

Supporting information

S1 Text. “Yes” and “no” thinking, GLM analysis, and classification using CSP.
(DOCX)

S1 Fig. Difference in hemodynamic response between “yes” and “no” thinking using general linear model.
(TIFF)

S2 Fig. Binary map of the *t* test between “yes” and “no” thinking.
(TIFF)

Author Contributions

Conceptualization: Ujwal Chaudhary, Niels Birbaumer.

Data curation: Ujwal Chaudhary, Sudhir Pathak.

Formal analysis: Ujwal Chaudhary, Sudhir Pathak.

Funding acquisition: Niels Birbaumer.

Investigation: Ujwal Chaudhary, Niels Birbaumer.

Methodology: Ujwal Chaudhary, Sudhir Pathak, Niels Birbaumer.

Project administration: Ujwal Chaudhary, Niels Birbaumer.

Resources: Niels Birbaumer.

Software: Ujwal Chaudhary.

Supervision: Ujwal Chaudhary, Niels Birbaumer.

Validation: Ujwal Chaudhary, Sudhir Pathak.

Visualization: Ujwal Chaudhary.

Writing – original draft: Ujwal Chaudhary.

Writing – review & editing: Ujwal Chaudhary, Niels Birbaumer.

References

1. Chaudhary U, Birbaumer N, Curado MR. Brain-Machine Interface (BMI) in paralysis. *Ann Phys Rehabil Med*. 2015; 58(1):9–13. <https://doi.org/10.1016/j.rehab.2014.11.002> PMID: 25623294
2. Chaudhary U, Birbaumer N, Ramos-Murguialday A. Brain-computer interfaces in the completely locked-in state and chronic stroke. In: *Progress in Brain Research*. 2016. p. 131–61.
3. Chaudhary U, Birbaumer N, Ramos-Murguialday A. Brain-computer interfaces for communication and rehabilitation. *Nat Rev Neurol*. 2016; 12(9):513–25. <https://doi.org/10.1038/nrneuro.2016.113> PMID: 27539560
4. Naito M, Michioka Y, Ozawa K, Ito Y, Kiguchi M, Kanazawa T. A communication means for totally locked-in ALS patients based on changes in cerebral blood volume measured with near-infrared light. *IEICE Trans Inf Syst*. 2007; E90–D(7):1028–37.
5. Gallegos-Ayala G, Furdea A, Takano K, Ruf CA, Flor H, Birbaumer N. Brain communication in a completely locked-in patient using bedside near-infrared spectroscopy. *Neurology*. 2014; 82(21):1930–2. <https://doi.org/10.1212/WNL.0000000000000449> PMID: 24789862
6. Chaudhary U, Xia B, Silvoni S, Cohen LG, Birbaumer N. Brain-Computer Interface-Based Communication in the Completely Locked-In State. *PLoS Biol*. 2017; 15(1):e1002593. <https://doi.org/10.1371/journal.pbio.1002593> PMID: 28141803
7. Guger C, Spataro R, Allison BZ, Heilinger A, Ortner R, Cho W, La Bella V. Complete locked-in and locked-in patients: command following assessment and communication with vibro-tactile P300 and motor imagery brain-computer interface tools. *Frontiers in neuroscience*. 2017; 11, p.251. <https://doi.org/10.3389/fnins.2017.00251> PMID: 28529473
8. Khan MJ, Hong KS. Passive BCI based on drowsiness detection: an fNIRS study. *Biomedical optics express*. 2015; 6(10):4063–78. <https://doi.org/10.1364/BOE.6.004063> PMID: 26504654
9. Chaudhary U, Hall M, DeCerce J, Rey G, Godavarty A. Frontal activation and connectivity using near-infrared spectroscopy: Verbal fluency language study. *Brain Res Bull*. 2011; 84(3):197–205. <https://doi.org/10.1016/j.brainresbull.2011.01.002> PMID: 21255633
10. Chaudhary U, Zhu B, Godavarty A. Frontal cortical connectivity and lateralisation of joint attention experience using near infrared spectroscopy. *Journal of Near Infrared Spectroscopy*. 2011; 19(2):105–16.
11. Hall M, Chaudhary U, Rey G, Godavarty A. Fronto-temporal mapping and connectivity using NIRS for language-related paradigms. *Journal of Neurolinguistics*. 2013; 26(1):178–94.
12. Chaudhary U, Hall M, Gonzalez J, Elbaum L, Bloyer M, Godavarty A. Motor response investigation in individuals with cerebral palsy using near infrared spectroscopy: pilot study. *Applied optics*. 2014; 53(3):503–10. <https://doi.org/10.1364/AO.53.000503> PMID: 24514139
13. Zhang S, Zheng Y, Wang D, Wang L, Ma J, Zhang J, Xu W, Li D. and Zhang D. Application of a common spatial pattern-based algorithm for an fNIRS-based motor imagery brain-computer interface. *Neuroscience letters*. 2017; 655, pp.35–40. <https://doi.org/10.1016/j.neulet.2017.06.044> PMID: 28663052
14. Tonin A, Birbaumer N, and Chaudhary U. A 20-Questions-Based Binary Spelling Interface for Communication Systems. *Brain sciences*. (2018); 8(7):126.
15. Heck DH, McAfee SS, Liu Y, Babajani-Feremi A, Rezaie R, Freeman WJ, Wheless JW, Papanicolaou AC, Ruzinko M, Sokolov Y. and Kozma R. Breathing as a fundamental rhythm of brain function. *Frontiers in neural circuits*, 2017; 10, p.115. <https://doi.org/10.3389/fncir.2016.00115> PMID: 28127277
16. Sterman MB. Effects of sensorimotor EEG feedback training on sleep and clinical manifestation of epilepsy. In: Beatty J. and Legewie H. (Eds) *Biofeedback and Behaviour*. Plenum Press: New York. (1977);167–200.
17. Mulholland T. *Biofeedback as scientific method*. Biofeedback Theory and Research. New York: Academic Press Inc. (1977).
18. Anderson SA. and Anderson P. *Physiological basis of the alpha rhythm*. New York: Appleton-Century-Crofts. (1968).
19. Klimesch W, Sauseng P, Hanslmayr S. EEG alpha oscillations: the inhibition-timing hypothesis. *Brain research reviews*. (2007); 53(1):63–88. <https://doi.org/10.1016/j.brainresrev.2006.06.003> PMID: 16887192
20. Kotchoubey B, Lang S, Mezger G, Schmalohr D, Schneck M, Semmler A, Bostanov V. and Birbaumer N. Information processing in severe disorders of consciousness: vegetative state and minimally conscious state. *Clinical Neurophysiology*. 2005; 116(10):2441–53. <https://doi.org/10.1016/j.clinph.2005.03.028> PMID: 16002333