

**S1 Table. Distribution of mosquito genera/species by season in rural and urban areas of Zanzibar.**

	<i>Ae. aegypti</i>	<i>Ae. simpsoni</i>	<i>Culex</i>	<i>An. gambiae</i> s.l.	<i>Eretmapodites</i>	Total
	N (% <sup>a</sup> )	N (% <sup>a</sup> )	N (% <sup>a</sup> )	N (% <sup>a</sup> )	N (% <sup>a</sup> )	
<b>Rural</b>	8,642 (82.6)	6 (0.1)	1,590 (15.2)	9 (0.1)	207 (2.0)	10,454
Wet	5,795 (85.1)	0 (0.0)	798 (11.8)	7 (0.1)	207 (3.0)	6,807
Dry	2,847 (78.1)	6 (0.2)	792 (21.7)	2 (0.1)	0 (0.0)	3,647
<b>Urban</b>	5,029 (64.3)	0 (0.0)	2,794 (35.7)	0 (0.0)	0 (0.0)	7,823
Wet	3,379 (92.9)	0 (0.0)	259 (7.1)	0 (0.0)	0 (0.0)	3,638
Dry	1,650 (39.4)	0 (0.0)	2,535 (60.6)	0 (0.0)	0 (0.0)	4,185
<b>Total</b>	13,671 (74.8)	6 (0.03)	4,384 (24.0)	9 (0.05)	207 (1.2)	18,277

<sup>a</sup> Percent of all mosquitoes by setting or season.