

S9 Table. Species included at the assemblage level in the energy crops and solar farm sensitivity maps.

Species ^a	Assemblage ^b
Grey partridge <i>Perdix perdix</i>	arable
Turtle dove <i>Streptopelia turtur</i>	arable
Sky Lark <i>Alauda arvensis</i>	arable
Yellow wagtail <i>Motacilla flava</i>	arable
Whinchat <i>Saxicola rubetra</i>	arable
Tree sparrow <i>Passer montanus</i>	arable
Linnet <i>Carduelis cannabina</i>	arable
Twite <i>Carduelis flavirostris</i>	arable
Reed bunting <i>Emberiza schoeniclus</i>	arable
Yellowhammer <i>Emberiza citrinella</i>	arable
Corn bunting <i>Miliaria calandra</i>	arable
Oystercatcher <i>Haematopus ostralegus</i>	farmland wader
Northern Lapwing <i>Vanellus vanellus</i>	farmland wader
Redshank <i>Tringa totanus</i>	farmland wader
Curlew <i>Numenius arquata</i>	farmland wader
Snipe <i>Gallinago gallinago</i>	farmland wader
Golden plover <i>Pluvialis apricaria</i>	upland wader
Dunlin <i>Calidris alpina</i>	upland wader
^a Data was obtained from <i>Bird Atlas 2007–11</i> [1], which is a joint project between BTO, BirdWatch Ireland and the Scottish Ornithologists' Club. ^b Areas were not categorised by land cover, so if any area had a high score based on either 'upland wader', 'farmland wader' or 'arable' assemblages then it qualified as medium sensitivity. [1] Balmer D, Gillings S, Caffrey B, Swann R, Downie I, Fuller R. <i>Bird Atlas 2007–11: the breeding and wintering birds of Britain and Ireland</i> . Thetford: BTO Books; 2013. ISBN 978-1-908581-28-0.	