

CORRECTION


# Correction: Evidence-Based Neonatal Unit Practices and Determinants of Postnatal Corticosteroid-Use in Preterm Births below 30 Weeks GA in Europe. A Population-Based Cohort Study

Alexandra Nuytten, H el ene Behal, Alain Duhamel, Pierre-Henri Jarreau, Jan Mazela, David Milligan, Ludwig Gortner, Aur elie Piedvache, Jennifer Zeitlin, Patrick Truffert, On behalf of the EPICE (Effective Perinatal Intensive Care in Europe) Research Group

The following information is missing from the Funding section: Funding for this open access publication was provided by the FP7 Post-Grant Open Access Pilot.

## Reference

1. Nuytten A, Behal H, Duhamel A, Jarreau P-H, Mazela J, Milligan D, et al. (2017) Evidence-Based Neonatal Unit Practices and Determinants of Postnatal Corticosteroid-Use in Preterm Births below 30 Weeks GA in Europe. A Population-Based Cohort Study. PLoS ONE 12(1): e0170234. doi: [10.1371/journal.pone.0170234](https://doi.org/10.1371/journal.pone.0170234) PMID: [28114369](https://pubmed.ncbi.nlm.nih.gov/28114369/)



## OPEN ACCESS

**Citation:** Nuytten A, Behal H, Duhamel A, Jarreau P-H, Mazela J, Milligan D, et al. (2017) Correction: Evidence-Based Neonatal Unit Practices and Determinants of Postnatal Corticosteroid-Use in Preterm Births below 30 Weeks GA in Europe. A Population-Based Cohort Study. PLoS ONE 12(2): e0172408. doi:10.1371/journal.pone.0172408

**Published:** February 13, 2017

**Copyright:**   2017 Nuytten et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.