

RESEARCH ARTICLE

Attitudes to mental illness among mental health professionals in Singapore and comparisons with the general population

Qi Yuan^{1*}, Louisa Picco¹, Sherilyn Chang¹, Edimansyah Abdin¹, Boon Yiang Chua¹, Samantha Ong², Kah Lai Yow³, Siow Ann Chong¹, Mythily Subramaniam¹

1 Research Division, Institute of Mental Health, Singapore, Singapore, **2** Nursing Administration, Institute of Mental Health, Singapore, Singapore, **3** Clinical and Allied Health Professionals Services, Institute of Mental Health, Singapore, Singapore

* Qi_YUAN@imh.com.sg



OPEN ACCESS

Citation: Yuan Q, Picco L, Chang S, Abdin E, Chua BY, Ong S, et al. (2017) Attitudes to mental illness among mental health professionals in Singapore and comparisons with the general population. *PLoS ONE* 12(11): e0187593. <https://doi.org/10.1371/journal.pone.0187593>

Editor: Jerson Laks, Universidade Federal do Rio de Janeiro, BRAZIL

Received: January 4, 2017

Accepted: October 23, 2017

Published: November 16, 2017

Copyright: © 2017 Yuan et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All individual data from this study resides with Office of Research, Institute of Mental Health. Data is not available for online access, however readers who wish to gain access to the data can write to the Clinical Research Committee, Institute of Mental Health/Woodbridge Hospital Secretariat at IMHRESEARCH@imh.com.sg. Access can be granted subject to the Institutional Review Board (IRB) and the research collaborative agreement guidelines. This is a requirement mandated for this research study by our IRB and funders.

Abstract

Background

Similar to the general public, mental health professionals sometimes also have negative attitudes towards individuals with mental illness; which could ultimately affect the quality of care received by the patients. This study aims to explore attitudes to mental illness among mental health professionals in Singapore; make comparisons with the general population; and investigate the significant correlates.

Methods

A cross-sectional design was used. Eligible participants were recruited from the Institute of Mental Health, Singapore. Attitudes to mental illness among the mental health professionals were measured using an adapted 26-item Attitudes to Mental Illness questionnaire (AMI). An earlier study amongst the general population in Singapore had used the same tool; however, factor analysis suggested a 20-item, 4-factor structure (AMI-SG) was the best fit. This 4-factor structure was applied among the current sample of mental health professionals to allow comparisons between the professionals and the general population.

Data were collected through an online survey tool 'Questionpro' from February to April 2016, and 379 participants were included in the current analysis. Attitudes to mental illness among these professionals were compared to those of the general population, which were captured as part of a national study conducted from March 2014 to April 2015.

Results

The 20-item, 4-factor structure AMI-SG derived from the general population was applicable among the mental health professionals in Singapore. Compared to the general population, mental health professionals had significantly more positive attitudes to mental illness; however their scores on 'social distancing' did not differ from the general population. Indian ethnicity was negatively associated with 'social distancing' and 'social restrictiveness' among the professionals; while higher education was negatively related to 'prejudice and

Funding: This study was funded by the Singapore Ministry of Health's National Medical Research Council under the Centre Grant Programme (Grant No.: NMRC/CG/004/2013). The funding recipient is SAC. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

misconception'. Compared to nurses, doctors showed significantly more positive attitudes on 'social restrictiveness' and 'prejudice and misconception'. Having family or close friends diagnosed with mental illness was negatively associated with 'social distancing' among the professionals.

Conclusion

The AMI-SG is an effective tool to measure attitudes to mental illness among mental health professionals in Singapore. Although the professionals had significantly more positive attitudes to mental illness than the general public in Singapore, their attitudes on 'social distancing' resembled closely that of the general public. Professionals tended to have more negative attitudes if they were nurses, less educated, and of Chinese ethnicity. More studies are needed to explore the underlying reasons for the differences and to generalize these findings among mental health professionals elsewhere.

Introduction

Attitudes to mental illness can encompass positive attitudes such as acceptance [1], more neutral attitudes like tolerance [2], to negative ones such as stigma [3] and even fear [4]. Previous studies suggested that negative attitudes towards people with mental illness were highly prevalent among the general public [5–8]. Similarly, mental health professionals may also have negative attitudes towards people with mental illness. A previous study in Brazil found that among a national sample of 1,414 psychiatrists, 42.9% of them were identified as having stereotype, prejudice and high social distance towards individuals with schizophrenia [9]. Nordt et al. [10] also found mental health professionals had similar level of negative attitudes about people with mental illness as the general public. Hansson et al. [11] examined attitudes of patients and mental health professionals towards people with mental illness and found that there was a high prevalence of negative attitudes towards people with mental illness among the mental health professionals, and the professionals' beliefs about people with mental illness were similar to those of the patients.

Negative attitudes among mental health professionals might affect patients in multiple ways. For example, Ellsworth [12] found that nurses' higher endorsement on restrictive attitudes were consistently related to more controlling and restricting behaviours among the nurses towards the patients. Other than the behaviours, negative attitudes have also been shown to affect the therapeutic alliance between professionals and patients [13]. A study done by Verhaeghe and Bracke [14] reported that mental health service users with more stigma experience during the service tended to show less trust and less satisfaction towards the service. Previous studies also suggested that psychiatric nurses could have negative attitudes towards specific mental disorders (e.g. borderline personality disorder [15]) and towards individuals with mental illness in general [16, 17]. However, nurses' attitudes are one of the most important factors in facilitating progress movement within the one-to-one therapeutic relationship [18, 19]. In all these circumstances, attitudes among the professionals could ultimately lower the quality of care received by the patients.

Despite some research highlighting the harmful impact these negative attitudes can have towards people with mental illness, others also suggested that mental health professionals, due to their knowledge and daily contact with individuals with mental illness, would have more

positive attitudes towards mental illness. Mårtensson et al. [20] found that compared to the general Swedish population, mental health nurses in Sweden had more positive attitudes to mental illness. Similar findings were also reported by Lauber et al. [21] who suggest that psychiatrists have significantly more positive attitudes to mental illness compared to the general population. Compared with non-mental health professionals, mental health professionals also possess more positive attitudes towards individuals with mental illness [22–24]. However, these studies all have their own limitations, such as small sample size [21–24] and poor analysis strategy (i.e. comparisons without controlling for potential confounding) [20, 23, 24]. Moreover, due to the inconsistent results of such comparisons, more studies are still needed in this area.

Unlike various studies comparing attitudes of mental health professionals with other healthcare professionals [22–24] or general public [10, 11, 20, 21], fewer studies have been done to explore correlates of attitudes to mental illness among mental health professionals. For example, Mårtensson et al. [20] found that nursing staff had more positive attitudes towards mental illness if their knowledge about mental illness was less stigmatized (i.e. higher endorsement on the six stigma related statements of the Mental Health Knowledge Schedule [25]), and if they had a friend with mental illness currently or in the past. Another study by Hansson and colleagues [11] suggested that mental health professionals' attitudes towards mental illness might be affected by their work setting characteristics, with staff working within inpatient services having more negative attitudes than those working in out-patient services. Cross-cultural studies on attitudes to mental illness among mental health professionals also suggest that cultural differences might contribute to the differences in attitudes. A previous study which compared Swiss mental health professionals with counterparts in Brazil showed that professionals in Switzerland had significantly higher levels of social distance, and age was a statistically significant predictor (i.e. younger age associated with less social distance) [26]. Another study among mental health nurses across 5 European countries (i.e. Finland, Lithuania, Ireland, Italy and Portugal) also suggested the attitudes differed by country of origin [27]. To the best of our knowledge, there is only one study which was conducted among Singapore mental health professionals to explore their attitudes to mental illness [28]. However, that study was conducted more than 10 years ago and the study sample only included nurses. Moreover, the bivariate analysis strategy used in that study could not exclude the potential confounding effects of other variables.

In order to address these gaps, the current study aims to 1) explore attitudes to mental illness among mental health professionals in Singapore; 2) compare the attitudes of mental health professionals with the local general population; 3) explore the correlates of attitudes to mental illness among mental health professionals.

Methods

Participants and procedure

Data relating to attitudes of mental health professionals were extracted from a study which aimed to explore associative stigma and positive mental health among staff working at the Institute of Mental Health (IMH), the only tertiary psychiatric service provider in Singapore [29]. This was a cross-sectional, online survey which used convenience sampling. Inclusion criteria comprised being: 1) Singapore citizens, Permanent Residents or Non-residents; 2) doctors, nurses, or allied health staff (i.e. psychologists, pharmacists, occupational therapists, physiotherapists, case managers or medical social workers) working at IMH during recruitment; 3) aged 21 years and above and; 4) able to complete the online survey in English. Given that we also aimed to compare the attitudes to mental illness of the mental health professionals with

that of the general population in Singapore, data from non-residents or work permit holders were excluded from the final analysis, to ensure comparability.

Data were collected from February to April 2016. Email invitations were sent to all nurses, doctors, and allied health staff through relevant hospital group email lists informing them of the study. By clicking the link enclosed in the email, potential participants were directed to an online portal including the screening questions. They were used to ensure the respondents met the inclusion criteria. For those who were screened out, an automatic email was sent notifying them that they were not eligible for this study. The remaining participants were then directed to the online consent form. Clicking the 'agree' button indicated an individual's willingness to participate in the study. The questionnaire was administered through an online survey tool 'Questionpro', and took around 10–15 minutes to complete. Upon completion, a SGD \$20 Starbucks voucher was given to the participants to compensate for their time and inconvenience. In total, 470 participants were recruited during the 2-month recruitment period, among which 8 cases were removed due to unreliable data or the respondents not meeting the inclusion criteria and another 83 cases were excluded from analysis due to them being non-residents. In all, 379 mental health professionals were included in the current study. Ethical approval was granted by the National Healthcare Group Domain Specific Review Board, Singapore.

Measurements

Attitudes to mental illness were measured by an adapted 26-item version Attitudes to Mental Illness questionnaire (AMI). The original AMI questionnaire has 27 items in total, and it was developed by the UK Department of Health [30, 31], based on the 40-item Community Attitudes toward the Mentally Ill Scale [32]. An earlier study conducted by the same study team had also used the AMI questionnaire among a national representative sample of the general population in Singapore [33]. Factor analysis of this 26-item AMI questionnaire revealed a 20-item, 4-factor structure which was the best fit for the general population in Singapore, and comprised: 'social distancing' (3 items), 'tolerance/support for community care' (9 items), 'social restrictiveness' (3 items) and 'prejudice and misconception' (5 items) [33]. Given the current study aimed to compare attitudes to mental illness between mental health professionals and the general population in Singapore, whilst the 26-item version of the AMI questionnaire was administered, analysis was based on the 20-item, 4-factor version. Henceforth, to differentiate these two versions, the 20-item version is referred to as the Attitudes to Mental Illness questionnaire—Singapore version (AMI-SG). The 20 items of the AMI-SG can be found in the [S1 Appendix](#).

The participants were required to rate items on a 5-point Likert scale ranging from '1 = strongly agree' to '5 = strongly disagree'. To enable easier interpretations, the AMI-SG items were reverse scored (changed to '1 = strongly disagree' to '5 = strongly agree') as per an earlier national study conducted in Singapore [33]. The total score of each factor was then summed and used in the multivariate regression analyses. For tolerance/support for community care, the reverse-scoring item 'Increased spending on mental health services is a waste of money' was reverse scored again, and then added up with the scores of the remaining items within this factor. As a result, more positive attitudes towards people with mental illness were characterized as—lower 'social distancing', 'social restrictiveness' and 'prejudice and misconception' and higher 'tolerance/support for community care' scores.

Socio-demographic information including age, gender, ethnicity, marital status, education level, residency status, and information on employment (i.e. position and years worked in

IMH), and whether any close family and friends had been diagnosed with mental illness was also collected.

Attitudes to mental illness among general population

Data on attitudes to mental illness among the Singapore general population (i.e. socio-demographic characteristics including age, gender, ethnicity, marital status, education level, and AMI-SG factor scores) were extracted from a previous national mental health literacy study conducted from March 2014 to April 2015 [33]. Since the main purpose of the current study was to explore attitudes to mental illness among mental health professionals, and the fact that detailed information of this national study could be found elsewhere [33]; in the current study, this extracted data were purely used in the comparison analysis.

Statistical analysis

Descriptive analyses were performed for socio-demographic and other dependent variables. Continuous variables were listed as mean and standard deviation (SD); for categorical variables, they were presented as frequency and percentage. Before further analysis, a confirmatory factor analysis (CFA) was conducted, to test the applicability of the 20-item, 4-factor AMI-SG derived from the general population in Singapore, among the current sample of mental health professionals, whilst also ensuring the construct validity [34] of AMI-SG among the mental health professionals before further comparisons. CFA was performed through the 'lavaan' package under R software [35], and adjusted for categorical variables with the estimator of 'Weighted Least Square Means and Variance Adjusted (WLSMV)' [36]. In the current study, an acceptable model was defined as 1) the comparative fit index (CFI) > 0.90; 2), the Tucker-Lewis index (TLI) > 0.90, and 3) the root mean square error of approximation (RMSEA) < 0.08 [37]; while for a good model, these indices should be 'CFI > 0.95, TLI > 0.95 and RMSEA < 0.06' [38]. The internal consistency indicators (Cronbach's alpha) were also calculated for each factor.

To compare the AMI-SG factor scores between the general population and mental health professionals in Singapore, the two datasets were combined with the included variables being recoded to ensure consistency across the two datasets. Given the study among the general population adopted a disproportionate stratified sampling design with 12 strata defined according to ethnicity (Chinese, Malay, Indian, Others) and age group (18–34, 35–49, 50–65 years) [33]; during the combination, a separate strata number (i.e. 13) and a weight equals to '1' (to indicate no sampling weight) were assigned to the mental health professionals sample. Multivariate regression was conducted with each of the AMI-SG factors being the dependent variable and mental health professionals or general population as the independent variable, after controlling for socio-demographic characteristics including age, gender, ethnicity, marital status and education level. This analysis was conducted via the 'PROC SURVEYREG' syntax in SAS9.3 [39]. The mean AMI-SG factor scores of the general population were also extracted from Yuan et al. [33].

Lastly, multivariate linear regression was performed to examine the significant correlates (e.g. socio-demographic characteristics such as age, gender and ethnicity, position, years worked at IMH and whether any close family and friends had been diagnosed with mental illness) of each of the AMI-SG factor scores (dependent variables) among the mental health professionals working at IMH. The descriptive analysis and this multivariate linear regression analysis were also conducted using SAS 9.3 [39]. For all regression analyses, a two-sided p-value below 0.05 was considered as statistically significant.

Results

The sample characteristics are presented in Table 1. The sample (n = 379) comprised 51 doctors, 137 nurses, and 191 allied health staff. The average age of the study sample was 37.4 (SD = 11.0) years, with the majority being female (65.7%), Chinese (70.5%), having a bachelor's degree or above (84.2%), and having worked at IMH for at least one year (91.6%). Only 30.9% participants reported that they had family or close friends who have been diagnosed with mental illness.

The CFA results confirmed that the 4-factor structure of AMI-SG derived from the general population in Singapore [33] was acceptable among mental health professionals working at IMH. The fit indices were $\chi^2_{(df)} = 485.086_{(164)}$ (p = 0.000), CFI = 0.930, TLI = 0.919, RMSEA = 0.072. The absolute value of the factor loading for each item varied from 0.451 to 0.956. The internal reliability statistics for the four factors, namely 'social distancing', 'tolerance/support for community care', 'social restrictiveness' and 'prejudice and misconception', were 0.791, 0.735, 0.663, and 0.671, respectively; where a cut-off of 0.6 was deemed as acceptable [40]. The mean AMI-SG factors scores among the mental health professionals were 7.52 for 'social distancing', 40.48 for 'tolerance/support for community care', 5.16 for 'social restrictiveness', and 11.20 for 'prejudice and misconception' (Table 2).

Table 3 shows the comparisons of AMI-SG factor scores between mental health professionals and the general population in Singapore. The results suggest that, after controlling for socio-demographic characteristics including age, gender, ethnicity, marital status and education level, mental health professionals had significantly more 'tolerance/support for

Table 1. Socio-demographic characteristics of the sample (n = 379).

Characteristics	Groups	Mean/Frequency	SD/%
Age (years)		37.4	11.0
Gender	Female	249	65.7
	Male	130	34.3
Ethnicity	Chinese	267	70.5
	Indian	53	14.0
	Malay	36	9.5
	Others (including Filipino and Myanmar)	23	6.1
Marital status	Ever married	216	57.0
	Never married	163	43.0
Education level	Secondary/ITE/'O' level	18	4.8
	A' level/diploma	42	11.1
	Bachelor	173	45.7
	Master or above	146	38.5
Nationality	Singapore Citizen	320	84.4
	Permanent Resident	59	15.6
Position	Doctors	51	13.5
	Allied health staff	191	50.4
	Nurses	137	36.2
Years in IMH	Less than 1 year	32	8.4
	1–5 years	148	39.1
	6–10 years	88	23.2
	More than 10 years	111	29.3
Family or close friends diagnosed with mental illness	Yes	117	30.9
	No	262	69.1

<https://doi.org/10.1371/journal.pone.0187593.t001>

Table 2. AMI-SG factor scores among mental health professionals and general population in Singapore.

	Mental Health Professionals		General Population	Score Range
	Mean	SD	Mean	
Social Distancing (3 items)	7.52	2.97	8.07	3–15
Tolerance/Support for community care (9 items)	40.48	4.46	14.81	9–45
Social Restrictiveness (3 items)	5.16	2.12	7.21	3–15
Prejudice & Misconception (5 items)	11.20	3.94	15.36	5–25

<https://doi.org/10.1371/journal.pone.0187593.t002>

community care’, less ‘social restrictiveness’, and less ‘prejudice and misconception’, compared to the general population. However, ‘social distancing’ scores did not differ between the two groups.

Multivariate linear regression results for all 4 AMI-SG factors among the mental health professionals working at IMH are presented in Table 4. The results revealed that Indians and those from ‘other’ ethnic groups (including Filipino and Myanmar) as well as those who had family or close friends diagnosed with mental illness, showed significantly less ‘social distancing’ towards individuals with mental illness; while mental health professionals who were ever married showed significantly more ‘social distancing’ towards people with mental illness. ‘Tolerance/support for community care’ was only correlated with education level, with individuals with Secondary/ITE/’O’ level education having significantly lower level of ‘tolerance/support for community care’ for mental illness. ‘Social restrictiveness’ was negatively associated with being Indian and being a doctor. While ‘prejudice and misconception’ toward people with mental illness was positively associated with lower education level and being a permanent resident, as well as negatively associated with being either a doctor or allied health staff.

Discussion

The CFA analysis confirmed that the 4-factor structure of AMI-SG derived from the Singapore general population was applicable among mental health professionals in Singapore. It suggested that AMI-SG had good construct validity in measuring attitudes to mental illness among mental health professionals. Compared to the general population, the professionals had significantly more positive attitudes to mental illness—less ‘social restrictiveness’ and ‘prejudice and misconception’; and more ‘tolerance/support to community care’. This finding is consistent with those reported by the majority of the studies in the literature [20, 21, 41, 42]. However, for ‘social distancing’, no statistically significant difference was identified. Lauber et al. [21] reported that ‘although accepting mental health facilities in the community,

Table 3. Comparison of AMI-SG factor scores between mental health professionals and general population in Singapore.

	Social Distancing			Tolerance/Support for community care			Social Restrictiveness			Prejudice & Misconception		
	β	95% CI		β	95% CI		β	95% CI		β	95% CI	
Mental health professionals	-0.258	-0.629	0.113	26.345*	25.790	26.900	-1.191*	-1.488	-0.893	-2.220*	-2.686	-1.755
General population	Ref			Ref			Ref			Ref		

Ref—reference group;

* p<.0001;

Controlled for socio-demographic characteristics including age, gender, ethnicity, marital status and education level;

<https://doi.org/10.1371/journal.pone.0187593.t003>

Table 4. Correlates of AMI-SG factor scores among mental health professionals.

	Social Distancing				Tolerance/Support for community care				Social Restrictiveness				Prejudice and Misconception			
	β	95% CI	p		β	95% CI	p		β	95% CI	p		β	95% CI	p	
Age	-0.014	-0.051	0.022	0.444	0.005	-0.051	0.061	0.851	-0.007	-0.033	0.019	0.593	-0.009	-0.049	0.032	0.674
Gender																
Female	-0.661	-1.332	0.009	0.053	0.453	-0.573	1.480	0.386	-0.223	-0.700	0.254	0.359	0.113	-0.624	0.851	0.762
Male	Ref				Ref				Ref				Ref			
Ethnicity																
Indian	-1.326	-2.280	-0.372	0.007	-0.653	-2.128	0.822	0.385	-0.811	-1.492	-0.129	0.020	-0.211	-1.260	0.837	0.692
Malay	-0.607	-1.811	0.596	0.322	0.898	-0.974	2.770	0.346	-0.519	-1.381	0.344	0.238	0.385	-0.955	1.724	0.573
Others	-1.705	-3.201	-0.210	0.026	-1.293	-3.579	0.993	0.267	-0.304	-1.368	0.761	0.575	0.707	-0.962	2.377	0.405
Chinese	Ref				Ref				Ref				Ref			
Marital status																
Ever married	0.795	0.104	1.486	0.024	-0.900	-1.964	0.164	0.097	0.128	-0.363	0.619	0.610	0.042	-0.717	0.801	0.913
Never married	Ref				Ref				Ref				Ref			
Education level																
Secondary/ITE/'O' level	0.028	-1.579	1.635	0.973	-4.582	-7.035	-2.130	0.0003	0.122	-1.019	1.263	0.834	5.070	3.303	6.837	<.0001
A' level/diploma	-0.566	-1.856	0.725	0.389	-1.693	-3.666	0.280	0.092	0.562	-0.368	1.491	0.236	4.120	2.695	5.545	<.0001
Bachelor	-0.045	-0.798	0.708	0.906	-0.907	-2.062	0.248	0.124	0.064	-0.470	0.597	0.815	1.537	0.711	2.364	0.0003
Masters or above	Ref				Ref				Ref				Ref			
Residency																
Permanent Resident	0.329	-0.660	1.317	0.513	0.939	-0.567	2.444	0.221	0.169	-0.540	0.877	0.639	1.450	0.360	2.540	0.009
Singapore Citizen	Ref				Ref											
Position																
Doctors	-0.470	-1.692	0.751	0.449	0.568	-1.303	2.439	0.551	-1.431	-2.299	-0.563	0.001	-2.338	-3.677	-0.999	0.0007
Allied health staff	-0.537	-1.435	0.361	0.240	0.085	-1.307	1.477	0.905	-0.427	-1.065	0.211	0.189	-1.880	-2.863	-0.897	0.0002
Nurses	Ref				Ref				Ref				Ref			
Years in IMH																
Less than 1 year	-0.656	-2.098	0.787	0.372	-0.630	-2.809	1.550	0.570	-0.128	-1.147	0.892	0.806	0.523	-1.054	2.100	0.515
1–5 years	-0.502	-1.488	0.484	0.317	-0.137	-1.645	1.372	0.859	-0.678	-1.379	0.022	0.058	-0.636	-1.735	0.462	0.256
6–10 years	0.071	-0.872	1.013	0.883	-0.788	-2.231	0.655	0.283	-0.363	-1.035	0.308	0.288	-0.596	-1.637	0.445	0.261
More than 10 years	Ref				Ref				Ref				Ref			
Family or close friends diagnosed with mental illness																
Yes	-0.947	-1.620	-0.275	0.006	0.813	-0.218	1.844	0.122	-0.303	-0.779	0.172	0.210	-0.321	-1.056	0.413	0.390
No	Ref				Ref				Ref				Ref			
R Square	0.096				0.081				0.102				0.385			

Ref—reference group;

<https://doi.org/10.1371/journal.pone.0187593.t004>

psychiatrists also agree mental health facilities are downgrading a residential area', and he named it as a minor form of the 'not in my back yard' phenomenon. Our study, on the other hand, provided more solid evidence for this phenomenon, suggesting that although mental health professionals tends to have more 'tolerance/support for community care', less 'social restrictiveness' and less 'prejudice and misconception', their desire for closeness or intimacy towards individuals with mental illness ('social distancing') resemble closely that of the general public.

Another interesting finding was how having 'family or close friends diagnosed with mental illness' affected the attitudes of the mental health professionals. Various studies have suggested that social contact is one of the most effective interventions to reduce mental health related stigma and discrimination among adults [43, 44]. This was also observed among the local

general population, i.e. a previous national study suggested that social contact was negatively associated with personal stigma and social distance [6]. In our study, the results suggested that among professionals, having family or close friends diagnosed with mental illness (personal contact experience) predicted significantly less social distance towards those with mental illness. This finding, together with the difference in attitudes to mental illness between mental health professionals and the general population suggests that although both professional contact and personal contact belongs to social contact; the mechanism of how they work might be totally different. This assumption is supported by the intergroup contact hypothesis by Allport [45] where professional and personal contact represent the typical contact experience between groups with unequal group status (i.e. professionals and patients) and equal group status (i.e. friend and friend). However, due to the limitation of observational study design; more studies, especially qualitative and longitudinal studies, are needed to further test this assumption and to explore the exact underlying mechanism for such differences.

The demographic correlates of AMI-SG domains among mental health professionals included ethnicity, marital status, education level, and residency status. Indian ethnicity was negatively associated with social distancing and social restrictiveness. This is consistent with findings from a multi-national study suggesting that pharmacy students from India had lower level of social distancing compared to students from Australia, Belgium, Estonia and Latvia [46]. Permanent residents showed more prejudice and misconception towards mental illness compared to citizens. One possible explanation for this finding could be the different training received by permanent residents and local mental health professionals. Permanent residents would normally be trained in their own countries (e.g. India and China), which usually adopt a different system compared to that in Singapore (e.g. the undergraduate training in Singapore has an emphasis on communication skills [47], while this is not the case in India and China [48]). Such differences might lead to different values and perspectives among the professionals, and in turn result in different views towards individuals with mental illness. Since ethnicity and residency status both are relevant to culture, their correlations with AMI-SG indicate the potential role of culture as an underlying factor that can influence attitudes to mental illness. Mental health professionals who were ever married tended to show more 'social distancing' towards mental illness compared to those who were never married. However, since the potential effect of marriage on professionals' attitudes was not tested among previous studies [10, 11, 20, 21], more empirical studies are still needed to confirm the relationship and explore the potential explanations. Regarding the finding on education level, for future attitude campaigns among mental health professionals, these should be designed and tailored towards those with lower education levels. Other factors like age and gender which were found to be significant predictors among the general population [33] did not have any effects on the AMI-SG domains among the mental health professionals.

The different occupations of the mental health professionals also contributed to differences in AMI-SG domain scores, with doctors showing significantly less 'social restrictiveness' and 'prejudice and misconception', while allied health staff also showed less 'prejudice and misconception' compared to nurses. This is quite surprising yet understandable. Two reasons might have led to such a phenomenon. First of all, the different roles of mental health professionals were determined largely by their training. Although all participants were working in a tertiary mental health hospital, a previous study suggested that more than half of the nurses working at IMH were not registered as psychiatric nurses [28]. Instead they received general nursing training before they started working at IMH, and then they received on-site training. While for doctors at IMH, they are normally trained specifically under psychiatry which focuses more on the mechanism of mental disorders and associated treatments. This enables doctors to have a better understanding of mental illness and the behaviours of the patients. Such

difference allows the doctors to make more objective judgement over the patients and how they might perform in society. Thus it's not surprising that doctors had less 'social restrictiveness' and less 'prejudice and misconception' towards individuals with mental illness. Another difference between doctors and nurses is that for doctors, their contact with patients is usually formalized and lasts for a certain period of time in the consultation room; this also applies to their contact with inpatients. For allied health staff, they are more supportive in nature compared to doctors and nurses; thus have relatively less direct contact with patients. Even for those whom need to interact with the patients, they mainly deal with short-stay patients whom are usually non-severe cases [28]. While for nurses, they work more in the inpatient settings (i.e. short- and long-stay patients) where they are heavily involved in the daily care of more severe patients; and these severe patients, compared to outpatients, are more likely to be less cooperative. According to Allport [45], intergroup contact would lead to favourable outcomes when the participants have equal status, common goals and intergroup cooperation. In this case, compared to doctors, nurses' contact experiences are more likely to violate these conditions; which in turn result in less improvement in their attitudes. This is consistent with assumption from researchers, indicating that increased personal and professional contact is associated with more positive attitudes to people with mental illness [20]; however, contact with individuals with more severe, long-term and recurrent mental illness might lead to comparatively more negative attitudes [11]. This explanation is supported by the findings from a previous study on attitudes towards people with mental illness among nurses working at IMH, which found nurses working in the short-stay wards had more positive attitudes than those working in the long-stay wards [28]. In this case, it is important to determine the level of involvement with the inpatient care of people with mental illness which would produce the most optimum shift in nurses' attitudes towards mental illness. For example, one could explore for nurses who are working in the long-stay wards, if providing more support to share their care responsibilities or having more rotations between short- and long-stay wards would improve their attitudes.

The current study has several strengths. Firstly, it is the first study comparing the attitudes to mental illness between the general public and the mental health professionals in Singapore. Secondly, the utilization of online tools during the data collection reduced efforts in data entry, and it also obviated the associated data entry errors. Moreover, the absence of interviewers during the data collection process also helped to avoid the social desirability bias. Lastly, the CFA analysis demonstrated that, while the assessment tool was originally designed for use in general population, its psychometrics confirm its applicability among mental health professionals.

These findings should be interpreted with the following limitations in mind. Firstly, the study participants were recruited through a convenience sample drawn from one single mental health institution in Singapore. Although it is the only tertiary mental health provider, the sample might not be representative of all mental health professionals in Singapore and was restricted to doctors, nurses and allied health staff and therefore this would affect the generalizability of the study findings. Secondly, since the email invitations were sent out through hospital group emails lists, it was not possible to track the exact number of staff who accessed that email; thus we are unable to provide the response rate. Thirdly, although we compared the attitudes to mental illness between mental health professionals and the general population, the two studies were conducted at different time periods through different data collection strategies which might affect their comparativeness. However, the socio-demographic characteristics were controlled for in the analysis which excluded some of the potential confounding effects. Lastly, although we examined the correlates of AMI-SG factors among mental health

professionals, the cross-sectional design did not allow any casual relationships to be established. More studies are needed to further explore this topic.

Conclusion

The current study confirmed that the AMI-SG is an effective tool to measure attitudes to mental illness among mental health professionals in Singapore, and it had the same factor structure as that among the general population. Although mental health professionals had significantly more positive attitudes to mental illness compared to the general population, their attitudes on 'social distancing' did not differ from the attitudes of the general population. Multivariate regression analysis suggested that attitudes to mental illness among the mental health professionals were negatively associated with Chinese ethnicity, being ever married, less educated, being a nurse, and positively associated with the status of having family or close friends diagnosed with mental illness. For future studies, researchers could 1) use more representative samples (i.e. samples across institutions providing psychiatric care); 2) explore the mechanism of how social contact affect attitudes to mental illness differently between the mental health professionals and the general public in the local context; 3) investigate the underlying mechanism of how culture might affect attitudes to mental illness among the mental health professionals; 4) explore the most appropriate level of involvement for mental health nurses in the inpatient care of individuals with mental illness in the long-stay wards.

Supporting information

S1 Appendix. Attitude to mental illness questionnaire–Singapore version (AMI-SG). (DOCX)

Acknowledgments

The authors would like to thank the study participants for devoting their time to complete the study, and thank Dr. Daniel Poremski from the Research Division, Institute of Mental Health for his valuable comments on certain points in the discussion.

Author Contributions

Conceptualization: Qi Yuan, Louisa Picco.

Data curation: Qi Yuan, Boon Yiang Chua.

Formal analysis: Qi Yuan.

Funding acquisition: Siow Ann Chong.

Investigation: Louisa Picco, Sherilyn Chang, Boon Yiang Chua, Samantha Ong, Kah Lai Yow.

Methodology: Qi Yuan, Edimansyah Abdin.

Project administration: Louisa Picco, Sherilyn Chang.

Resources: Siow Ann Chong, Mythily Subramaniam.

Software: Qi Yuan, Boon Yiang Chua.

Supervision: Siow Ann Chong, Mythily Subramaniam.

Validation: Louisa Picco, Edimansyah Abdin, Mythily Subramaniam.

Visualization: Qi Yuan.

Writing – original draft: Qi Yuan.

Writing – review & editing: Louisa Picco, Sherilyn Chang, Edimansyah Abdin, Boon Yiang Chua, Samantha Ong, Kah Lai Yow, Siow Ann Chong, Mythily Subramaniam.

References

1. Angermeyer MC, Holzinger A, Carta MG, Schomerus G. Biogenetic explanations and public acceptance of mental illness: systematic review of population studies. *The British Journal of Psychiatry*. 2011; 199(5):367–72. <https://doi.org/10.1192/bjp.bp.110.085563> PMID: 22045945
2. Parra F. Social tolerance of the mentally ill in the Mexican American community. *International Journal of Social Psychiatry*. 1985; 31(1):37–45. <https://doi.org/10.1177/002076408503100104> PMID: 3972491
3. Corrigan PW, Watson AC. Understanding the impact of stigma on people with mental illness. *World psychiatry*. 2002; 1(1):16–20. PMID: 16946807
4. Brockington IF, Hall P, Levings J, Murphy C. The community's tolerance of the mentally ill. *The British Journal of Psychiatry*. 1993; 162(1):93–9.
5. Angermeyer MC, Dietrich S. Public beliefs about and attitudes towards people with mental illness: a review of population studies. *Acta Psychiatrica Scandinavica*. 2006; 113(3):163–79. <https://doi.org/10.1111/j.1600-0447.2005.00699.x> PMID: 16466402
6. Subramaniam M, Abdin E, Picco L, Pang S, Shafie S, Vaingankar J, et al. Stigma towards people with mental disorders and its components—a perspective from multi-ethnic Singapore. *Epidemiology and psychiatric sciences*. 2016:1–12.
7. Stuart H, Arboleda-Florez J. Community attitudes toward people with schizophrenia. *The Canadian Journal of Psychiatry*. 2001; 46(3):245–52. <https://doi.org/10.1177/070674370104600304> PMID: 11320678
8. Angermeyer MC, Breier P, Dietrich S, Kenzine D, Matschinger H. Public attitudes toward psychiatric treatment. *Social psychiatry and psychiatric epidemiology*. 2005; 40(11):855–64. <https://doi.org/10.1007/s00127-005-0958-x> PMID: 16215879
9. Loch AA, Guarniero FB, Lawson FL, Hengartner MP, Rössler W, Gattaz WF, et al. Stigma toward schizophrenia: do all psychiatrists behave the same? Latent profile analysis of a national sample of psychiatrists in Brazil. *BMC psychiatry*. 2013; 13(1):92.
10. Nordt C, Rössler W, Lauber C. Attitudes of mental health professionals toward people with schizophrenia and major depression. *Schizophrenia bulletin*. 2006; 32(4):709–14. <https://doi.org/10.1093/schbul/sbj065> PMID: 16510695
11. Hansson L, Jormfeldt H, Svedberg P, Svensson B. Mental health professionals' attitudes towards people with mental illness: Do they differ from attitudes held by people with mental illness? *International Journal of Social Psychiatry*. 2013; 59(1):48–54. <https://doi.org/10.1177/0020764011423176> PMID: 21954319
12. Ellsworth RB. A behavioral study of staff attitudes toward mental illness. *Journal of Abnormal Psychology*. 1965; 70(3):194.
13. Nakash O, Nagar M, Levav I. Predictors of mental health care stigma and its association with the therapeutic alliance during the initial intake session. *Psychotherapy Research*. 2015; 25(2):214–21. <https://doi.org/10.1080/10503307.2014.885147> PMID: 24527724
14. Verhaeghe M, Bracke P. Stigma and trust among mental health service users. *Archives of psychiatric nursing*. 2011; 25(4):294–302. <https://doi.org/10.1016/j.apnu.2011.02.001> PMID: 21784287
15. Markham D, Trower P. The effects of the psychiatric label 'borderline personality disorder' on nursing staff's perceptions and causal attributions for challenging behaviours. *British Journal of Clinical Psychology*. 2003; 42(3):243–56.
16. Ebrahimi H, Namdar H, Vahidi M. Mental illness stigma among nurses in psychiatric wards of teaching hospitals in the north-west of Iran. *Iranian journal of nursing and midwifery research*. 2012; 17(7):534. PMID: 23922602
17. Delaney KR. Psychiatric mental health nurses: Stigma issues we fail to see. *Archives of psychiatric nursing*. 2012; 26(4):333–5. <https://doi.org/10.1016/j.apnu.2012.04.006> PMID: 22835753
18. Forchuk C, Westwell J, Martin ML, Azzopardi WB, Kosterewa-Tolman D, Hux M. Factors Influencing Movement of Chronic Psychiatric Patients From the Orientation to the Working Phase of the Nurse-Client Relationship on an Inpatient Unit. *Perspectives in Psychiatric Care*. 1998; 34(1):36–44. PMID: 9847826
19. Forsyth A. The effects of diagnosis and non-compliance attributions on therapeutic alliance processes in adult acute psychiatric settings. *Journal of Psychiatric and Mental Health Nursing*. 2007; 14(1):33–40. <https://doi.org/10.1111/j.1365-2850.2007.01036.x> PMID: 17244003

20. Mårtensson G, Jacobsson JW, Engström M. Mental health nursing staff's attitudes towards mental illness: an analysis of related factors. *Journal of psychiatric and mental health nursing*. 2014; 21(9):782–8. <https://doi.org/10.1111/jpm.12145> PMID: 24654776
21. Lauber C, Anthony M, Ajdacic-Gross V, Rössler W. What about psychiatrists' attitude to mentally ill people? *European Psychiatry*. 2004; 19(7):423–7. <https://doi.org/10.1016/j.eurpsy.2004.06.019> PMID: 15504649
22. Gateshill G, Kucharska-Pietura K, Wattis J. Attitudes towards mental disorders and emotional empathy in mental health and other healthcare professionals. *The Psychiatrist*. 2011; 35(3):101–5.
23. Kopera M, Suszek H, Bonar E, Myszk M, Gmaj B, Ilgen M, et al. Evaluating explicit and implicit stigma of mental illness in mental health professionals and medical students. *Community mental health journal*. 2015; 51(5):628–34. <https://doi.org/10.1007/s10597-014-9796-6> PMID: 25535045
24. Smith AL, Cashwell CS. Social distance and mental illness: Attitudes among mental health and non-mental health professionals and trainees. *The Professional Counselor: Research and Practice*. 2011; 1(1):13–20.
25. Evans-Lacko S, Little K, Meltzer H, Rose D, Rhydderch D, Henderson C, et al. Development and psychometric properties of the Mental Health Knowledge Schedule. *Canadian journal of psychiatry Revue canadienne de psychiatrie*. 2010; 55(7):440–8. Epub 2010/08/14. <https://doi.org/10.1177/070674371005500707> PMID: 20704771.
26. Hengartner MP, Loch AA, Lawson FL, Guarniero FB, Wang Y-P, Rössler W, et al. Attitudes of mental health professionals towards persons with schizophrenia: a transcultural comparison between Switzerland and Brazil. *Archives of Clinical Psychiatry (São Paulo)*. 2012; 39(4):115–21.
27. Chambers M, Guise V, Välimäki M, Botelho MAR, Scott A, Staniulienė V, et al. Nurses' attitudes to mental illness: A comparison of a sample of nurses from five European countries. *International Journal of Nursing Studies*. 2010; 47(3):350–62. <https://doi.org/10.1016/j.ijnurstu.2009.08.008> PMID: 19804882
28. Tay SE, Pariyasami S, Ravindran K, Ali MI, Rowsudeen MT. Nurses' attitudes toward people with mental illnesses in a psychiatric hospital in Singapore. *J Psychosoc Nurs Ment Health Serv*. 2004; 42(10):40–7. Epub 2004/11/17. PMID: 15543671.
29. Picco L, Yuan Q, Vaingankar JA, Chang S, Abdin E, Chua HC, et al. Positive mental health among health professionals working at a psychiatric hospital. *PloS one*. 2017; 12(6):e0178359. <https://doi.org/10.1371/journal.pone.0178359> PMID: 28591203
30. UK Department of Health. Attitudes to Mental Illness 2014 Research Report. Department of Health, United Kingdom, 2014.
31. Mehta N, Kassam A, Leese M, Butler G, Thornicroft G. Public attitudes towards people with mental illness in England and Scotland, 1994–2003. *The British Journal of Psychiatry*. 2009; 194(3):278–84. <https://doi.org/10.1192/bjp.bp.108.052654> PMID: 19252160
32. Taylor SM, Dear MJ. Scaling community attitudes toward the mentally ill. *Schizophrenia Bulletin*. 1981; 7(2):225. PMID: 7280561
33. Yuan Q, Abdin E, Picco L, Vaingankar JA, Shahwan S, Jeyagurunathan A, et al. Attitudes to Mental Illness and Its Demographic Correlates among General Population in Singapore. 2016; 11(11): e0167297.
34. Boelen PA, van den Hout MA, van den Bout J. The factor structure of posttraumatic stress disorder symptoms among bereaved individuals: A confirmatory factor analysis study. *Journal of Anxiety Disorders*. 2008; 22(8):1377–83. <https://doi.org/10.1016/j.janxdis.2008.01.018> PMID: 18342486
35. Rosseel Y. lavaan: An R Package for Structural Equation Modeling. 2012. 2012; 48(2):36. Epub 2012-05-24. <https://doi.org/10.18637/jss.v048.i02>
36. Rosseel Y. The lavaan tutorial. Department of Data Analysis Ghent University. 2013.
37. Kline RB. Principles and practice of structural equation modeling: Guilford publications; 2015.
38. Lt Hu, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural equation modeling: a multidisciplinary journal*. 1999; 6(1):1–55.
39. Inc. SI. SAS® 9.3 Software. SAS Institute Inc Cary, NC; 2011.
40. Moss S, Prosser H, Costello H, Simpson N, Patel P, Rowe S, et al. Reliability and validity of the PAS-ADD Checklist for detecting psychiatric disorders in adults with intellectual disability. *Journal of Intellectual Disability Research*. 1998; 42(2):173–83.
41. Wahl O, Aroesty-Cohen E. Attitudes of mental health professionals about mental illness: a review of the recent literature. *Journal of Community Psychology*. 2010; 38(1):49–62.
42. Stuber JP, Rocha A, Christian A, Link BG. Conceptions of mental illness: Attitudes of mental health professionals and the general public. *Psychiatric services*. 2014; 65(4):490–7. <https://doi.org/10.1176/appi.ps.201300136> PMID: 24430508

43. Thornicroft G, Mehta N, Clement S, Evans-Lacko S, Doherty M, Rose D, et al. Evidence for effective interventions to reduce mental-health-related stigma and discrimination. *The Lancet*. 2016; 387 (10023):1123–32.
44. Corrigan PW, Morris SB, Michaels PJ, Rafacz JD, Rüsch N. Challenging the public stigma of mental illness: a meta-analysis of outcome studies. *Psychiatric Services*. 2012.
45. Allport GW. *The nature of prejudice: Basic books*; 1979.
46. Bell J, Aaltonen S, Airaksinen M, Volmer D, Gharat M, Muceniec R, et al. Determinants of mental health stigma among pharmacy students in Australia, Belgium, Estonia, Finland, India and Latvia. *International journal of social psychiatry*. 2009.
47. Kua EH. Focus on psychiatry in Singapore. *The British Journal of Psychiatry*. 2004; 185(1):79–82.
48. Zisook S, Balon R, Björkstén KS, Everall I, Dunn L, Ganadjian K, et al. Psychiatry residency training around the world. *Academic Psychiatry*. 2007; 31(4):309–25. <https://doi.org/10.1176/appi.ap.31.4.309> PMID: 17626194