

1 Honey bee and native solitary bee foraging behavior in a crop with dimorphic  
2 parental lines

3

4 María Cecilia Estravis Barcala<sup>1,2,¶</sup>; Florencia Palottini<sup>1,2,¶</sup>, Walter Marcelo Farina<sup>1,2,\*</sup>.

5

6

7

8 <sup>1</sup>Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de  
9 Biodiversidad y Biología Experimental, Laboratorio de Insectos Sociales.

10

11 <sup>2</sup>CONICET-Universidad de Buenos Aires, Instituto de Fisiología, Biología Molecular y Neurociencias  
12 (IFIBYNE), Buenos Aires, Argentina.

13

14

15

16 \*Correspondence author

17 E-mail: [walter@fbmc.fcen.uba.ar](mailto:walter@fbmc.fcen.uba.ar)

18

19 ¶ These authors contributed equally to this work.

20

21

22 **S1 Table. Set of variables considered in the generalized linear model explaining the**  
 23 **foraging behavior of *Apis mellifera* and *Melissodes* spp.**

24

Section	Variable	Statistics value	P
Number of inflorescences visited	Visitor x Parental line	LR= 0.4928	0.4827
	Parental line	LR= 0.01598	0.89939
	Visitor	LR= 37.5487	8.92e-10
Resource exploited	Visitor x Parental line	LR= 30.752	2.932e-08
Resource exploited on MF	Visitor x Time of day	LR=3.6081	0.0575
	Visitor	LR=37.741	8.079e-10
	Time of day	LR=50.438	1.23e-12

25

26