

S6 Table. Correlations between T cell phenotype, time after transplantation and EBV load of kidney transplant recipients

T cell subpopulations	Time post-transplant (years)		EBV load in Log copies/mL	
	Rho	P-value ^a	Rho	P-value ^a
% CD95 ⁺ CD4 ⁺ T cells	-0.4620	0.1735	0.1438	0.6900
% Naive CD4 ⁺ T cells (CD45RA ⁺ CCR7 ⁺)	0.3161	0.3707	0.0187	0.9651
% Effector memory CD4 ⁺ T cells (CD45RA ⁻ CCR7 ⁻)	0.1337	0.7127	0.0750	0.8386
% HLA-DR ⁺ CD8 ⁺ T cells	0.3830	0.2726	0.4627	0.1779
% HLA-DR ⁺ CD38 ⁺ CD8 ⁺ T cells	-0.2188	0.5284	0.6003	0.0721
% Naive CD8 ⁺ T cells (CD45RA ⁺ CCR7 ⁺)	0.1398	0.6990	-0.2126	0.4936
% central memory CD8 ⁺ T cells (CD45RA ⁻ CCR7 ⁺)	-0.1022	0.7166	0.0509	0.8880
% Terminally differentiated effector memory CD8 ⁺ T cells (CD45RA ⁺ CCR7 ⁻)	0.2979	0.3992	0.3939	0.2580

EBV, Epstein-Barr virus. ^aBonferroni significance threshold ≤ 0.0041