


A gyrokinetic model for the tokamak periphery

R. Jorge , B. Frei, P. Ricci, A. Baillod, N. F. Loureiro



**SWISS PLASMA
CENTER**

The Tokamak Periphery = Edge + SOL



- ❑ Core boundary conditions
- ❑ Heat exhaust
- ❑ Plasma fueling and ashes removal
- ❑ Impurity control

Properties of Periphery Turbulence

❑ Low-frequency $\omega \ll \Omega_i$

❑ Large Scale Fluctuations

$$k_{\perp} \rho_i \ll 1 \quad \frac{e\phi}{T_e} \sim 1$$

❑ Small Scale Fluctuations

$$k_{\perp} \rho_i \sim 1 \quad \frac{e\phi}{T_e} \ll 1$$

❑ Wide range of temperatures
and densities

$$T_e \sim 1 - 10^3 \text{ eV} \quad n \sim 10^{18} - 10^{20} \text{ m}^{-3}$$



Properties of Periphery Turbulence

❑ Low-frequency $\omega \ll \Omega_i$

❑ Large Scale Fluctuations

$$k_{\perp} \rho_i \ll 1 \quad \frac{e\phi}{T_e} \sim 1$$



Gyrokinetic Theory

$$\epsilon = k_{\perp} \rho_i \frac{e\phi}{T_e} \ll 1$$

❑ Small Scale Fluctuations

$$k_{\perp} \rho_i \sim 1 \quad \frac{e\phi}{T_e} \ll 1$$

❑ Wide range of temperatures

and densities

$$T_e \sim 1 - 10^3 \text{ eV} \quad n \sim 10^{18} - 10^{20} \text{ m}^{-3}$$



Arbitrary Collisionality

First Heroic Steps On The Way To An Edge Model

	Qin et al. 2007¹	Hahm et al. 2009²	Dimits et al. 2012³
Small Scale Fluctuations	EM $O(\epsilon^2)$	EM $O(\epsilon^2)$	EM $O(\epsilon^2)$
Large Scale Fluctuations	EM $O(\epsilon)$	ES $O(\epsilon^2)$	ES $O(\epsilon^2)$
Poisson's Eq.	$\int (\dots) d^3v$	Long-Wavelength Limit	$\int (\dots) d^3v$
Collisions	No	No	No

¹Qin et al., Physics of Plasmas **14**, 056110 (2007)

²Hahm et al., Physics of Plasmas **16**, 022305 (2009)

³Dimits, Physics of Plasmas **19**, 022504 (2012)

Our Model

Retain

- ❑ Both large scale (full-F) and small scale fluctuations
- ❑ Second order
- ❑ Full Coulomb collisions
- ❑ Numerical efficiency

How we proceed



Single Particle Dynamics - Ordering Assumptions

Magnetized Plasma

$$\frac{\omega}{\Omega_i} \sim \epsilon \quad k_{\perp} \rho_i \sim \epsilon \quad \frac{k_{\parallel}}{k_{\perp}} \sim \epsilon \quad \frac{\rho_i}{L_p} \sim \epsilon$$

Fluctuation Scales and Amplitude

$$k_{\perp} \rho_i \frac{e\phi}{T_e} = \epsilon \ll 1$$


Electromagnetic

$$v_{\parallel} A_{\parallel} \sim \phi$$

Arbitrary Collisionalities
(still magnetized)

$$\frac{\nu_{ei}}{\Omega_i} \lesssim \epsilon$$

Single Particle Dynamics – Hamiltonian Perturbation Theory




Particle Lagrangian

$$L(\mathbf{x}, \mathbf{v})$$

2-Step Phase-Space



Reduction



Lagrangian

$$\Gamma(\mathbf{R}, v_{\parallel}, \mu)$$

(cyclic coordinate θ , conserved μ)

Single Particle Dynamics – First Step

$$\text{Large Scales } \epsilon \sim k_{\perp} \rho_i \ll 1, \quad \frac{e\phi}{T_e} \sim 1$$

Start from

$$L = q\mathbf{A} \cdot \dot{\mathbf{x}} - q\phi - \frac{mv^2}{2}$$



Split between parallel and perpendicular velocity

Describe **guiding center** motion $O(\epsilon^2)$

$$L = q\mathbf{A}^* \cdot \dot{\mathbf{R}} - q\phi^* - \frac{mv_{\parallel}^2}{2} + \mu \frac{m\dot{\theta}}{q}$$

Single Particle Dynamics – Second Step

$$\text{Small Scales } \epsilon \sim \frac{e\phi}{T_e} \ll 1, k_{\perp}\rho_i \sim 1$$

Start from

$$L = q\mathbf{A}^* \cdot \dot{\mathbf{R}} - q\phi^* - \frac{mv_{\parallel}^2}{2} + \mu \frac{m\dot{\theta}}{q}$$

Introduce small scale fluctuations

Describe **gyro center** motion $O(\epsilon^2)$

$$L = q\mathbf{A}^* \cdot \dot{\mathbf{R}} - q\phi^* - \frac{mv_{\parallel}^2}{2} + \mu \frac{m\dot{\theta}}{q} - q \langle \phi - v_{\parallel} A_{\parallel} \rangle - q \frac{\partial \langle (\phi - v_{\parallel} A_{\parallel})^2 \rangle}{\partial \mu}$$

Single Particle Dynamics – Equations of Motion

$$\dot{\mathbf{R}} = v_{\parallel} \mathbf{b} + \mathbf{v}_{\mathbf{E} \times \mathbf{B}} + \text{Other Drifts}$$



Including large scale $\frac{\nabla \phi(\mathbf{R}) \times \mathbf{B}}{B^2}$

and small scale $\frac{\nabla \langle \phi(\mathbf{x}) \rangle \times \mathbf{B}}{B^2}$ fluctuations

Curvature Drift

Polarization Drift

Non-Linear Drifts

Single Particle Dynamics – Equations of Motion

$$\dot{\mathbf{R}} = v_{\parallel} \mathbf{b} + \mathbf{v}_{\mathbf{E} \times \mathbf{B}} + \text{Other Drifts}$$

$$\dot{v}_{\parallel} = qE_{\parallel} + \mu \nabla_{\parallel} B + \text{Non-Linear Forces}$$



Including large scale $\nabla_{\parallel} \phi(\mathbf{R})$

and small scale $\nabla_{\parallel} \langle \phi(\mathbf{x}) \rangle$ fluctuations

Single Particle Dynamics – Equations of Motion

$$\dot{\mathbf{R}} = v_{\parallel} \mathbf{b} + \mathbf{v}_{\mathbf{E} \times \mathbf{B}} + \text{Other Drifts}$$

$$\dot{v}_{\parallel} = qE_{\parallel} + \mu \nabla_{\parallel} B + \text{Non-Linear Forces}$$

$$\dot{\mu} = 0$$



Conserved Adiabatic Invariant

From Single Particle to Particle Distribution

Gyrokinetic Equation
$$\frac{\partial F}{\partial t} + \dot{\mathbf{R}} \cdot \nabla F + \dot{v}_{\parallel} \frac{\partial F}{\partial v_{\parallel}} = \langle C(F) \rangle$$

Challenges

- ❑ 5-D + time
- ❑ Full Coulomb Collisions
- ❑ Coupling to Maxwell's equations (integro-differential system)

These challenges can be successfully approached by
using a moment hierarchy

Our Goal – Turn Gyrokinetic Eq. Into a Hierarchy of Fluid-Like Eqs.

Expand GK Equation into a Set of 3D

Moment Hierarchy Equations

$$\frac{dn}{dt} = \dots$$

$$\frac{dv}{dt} = \dots$$

$$\frac{dT}{dt} = \dots$$

...



Retain necessary kinetic effects and no more

Advantages of a Moment Hierarchy Model

Set of fluid-like equations with reasonable computational cost

Tune the number of moments according to the desired level of accuracy (function of collisionality)

Reduce to the fluid model at high collisionality

3 Steps To Build a Moment Hierarchy Model

1. Choose an orthogonal polynomial basis for F

$$F = F_M \sum_{p,j} N^{pj}(\mathbf{R}) H_p(v_{||}) L_j(\mu)$$

2. Project kinetic equation onto basis

$$\int (\text{GK Eq.}) H_p L_j dv_{||} d\mu$$

3. Obtain evolution equation for the basis coefficients

$$\frac{\partial N^{pj}}{\partial t} = \dots$$

From Gyrokinetic Equation to Moment Hierarchy

$$\int (\text{GK Eq.}) H_p L_j dv_{\parallel} d\mu$$



Spatial evolution of
Moments + Fields

Fluid Operator
(density, velocity,
temperature)

$$\boxed{\frac{\partial N^{pj}}{\partial t}} + \boxed{\nabla \cdot \dot{\mathbf{R}}^{pj}} - \boxed{\frac{\sqrt{2p}}{v_{th}} \dot{v}_{\parallel}^{p-1j}} + \boxed{\mathcal{F}^{pj}} = \boxed{C^{pj}}$$

Time Evolution

Forces included at $p > 0$

Collisions

Example – 1D Linear Gyrokinetic Moment Hierarchy

Phase Mixing
(coupling with other moments)

Collisions

$$\boxed{\frac{\partial N^{pj}}{\partial t}} + \boxed{\frac{1}{2} \frac{\partial N^{p+1j}}{\partial z} + p \frac{\partial N^{p-1j}}{\partial z}} = \boxed{2\delta_{p,1} K_j(k_{\perp} \rho) E_{\parallel}} + \boxed{C^{pj}}$$

Time Evolution

Electric Field Drive

Example – 1D Linear Gyrokinetic Moment Hierarchy

Phase Mixing
(coupling with other moments)

Collisions

$$\boxed{\frac{\partial N^{pj}}{\partial t}} + \boxed{\frac{1}{2} \frac{\partial N^{p+1j}}{\partial z} + p \frac{\partial N^{p-1j}}{\partial z}} = \boxed{2\delta_{p,1} K_j(k_{\perp} \rho) E_{\parallel}} + \boxed{C^{pj}}$$

Time Evolution

Kernel $K_j(x) = \frac{1}{j!} \left(\frac{x}{2}\right)^{2j} e^{-x^2/4}$



Analytical Closed formula for the Gyroaverage operation

Full finite Larmor radius effects

Projection of the Full Coulomb Collision Operator

$$C^{pj} = \int \langle C(F) \rangle H_p L_j dv_{\parallel} d\mu$$

$$\text{with } C(F) = \frac{\partial}{\partial \mathbf{v}} \cdot [\mathbf{H}(F)F] + \frac{\partial^2}{\partial \mathbf{v} \partial \mathbf{v}} : [\mathbf{G}(F)F]$$

- ❑ Bilinear
- ❑ Tensorial Nature
- ❑ Gyroaveraging Operation
- ❑ No parallel/perpendicular velocity symmetries

$$H(F) \sim \int \frac{F(\mathbf{v}')}{|\mathbf{v} - \mathbf{v}'|} d\mathbf{v}'$$

Not immediate...

Collision Operator - Spherical Harmonic Decomposition

Expand the Rosenbluth potentials

$$H(F) \sim \int \frac{F(\mathbf{v}')}{|\mathbf{v} - \mathbf{v}'|} d\mathbf{v}'$$

Electrostatic potential of a charge distribution F in velocity space

in a Taylor series

$$\frac{1}{|\mathbf{v} - \mathbf{v}'|} = \begin{cases} \sum_l \frac{(-\mathbf{v}')^l}{l!} \cdot \partial_{\mathbf{v}}^l \left(\frac{1}{v} \right), & v' \leq v \\ \sum_l \frac{(-\mathbf{v})^l}{l!} \cdot \partial_{\mathbf{v}'}^l \left(\frac{1}{v'} \right) & v < v' \end{cases}$$


with spherical harmonics as coefficients

$$(-1)^l v^{l+1} \left(\frac{\partial}{\partial \mathbf{v}} \right)^l \frac{1}{v} \sim \sum_{m=-l}^l Y_{lm}(\varphi, \theta) \hat{\mathbf{e}}^{lm}$$

Basis Tensors $\hat{\mathbf{e}}^{lm}$

Spherical Harmonics

Gyroangle



Collision Operator – Expansion in Spherical Harmonics

$$C[F] \sim C[Y_{lm}] \sim Y_{lm}$$

Gyroaveraging Procedure

Large Scales $k_{\perp} \rho_i \ll 1$

$$\langle Y_{lm}(\varphi, \theta) \rangle \longrightarrow H_p(v_{\parallel}) L_j(\mu)$$

Small Scales $\frac{e\phi}{T_e} \ll 1$

$$\langle Y_{lm}(\varphi, \theta) e^{in\theta} \rangle \longrightarrow f(n) K_j(k_{\perp} \rho) H_p(v_{\parallel}) L_j(\mu)$$

Moments of the Collisional Operator

$$C^{pj} = \int \langle C(F) \rangle H_p L_j dv_{||} d\mu$$

After integration



$$C^{pj} = \text{Kernel} \times f(n) \times \text{moments } N^{pj} \text{ of } F$$

Physics That We Are Able To Capture – High Collisionality

$$F = F_M(1 + \delta F) \quad \text{with} \quad \delta F = \sum_{p,j} N^{pj} H_p L_j$$

Semi-collisional closure

$$\delta F \sim \frac{\lambda_{\text{mfp}}}{L_{\parallel}}$$



Drift-Reduced Braginskii
equations retrieved

$$N^{30} \sim q_{\parallel} = -\chi_{\parallel} \nabla_{\parallel} T$$

Jorge et al., JPP **83**, 6 (2018)

Towards a Numerical Implementation

Numerical and theoretical investigation of linear modes

GK non-linear model reduced
to the drift-kinetic linear limit


$$\frac{\partial N^{pj}}{\partial t} = \sum_{s,t} D_{st}^{pj} N^{st}$$

- ☐ Compute collisional/hierarchy coefficients D_{st}^{pj}
- ☐ Solve time evolution problem
- ☐ Perform eigenvalue analysis
- ☐ Compare with collisionless result


Electron Plasma Waves –Damping at Arbitrary Collisionality

- One spatial and two velocity dimensions
- Electron perturbations only
- Time-evolution of $N^{00} \sim \phi$


Collisional and Landau Damping computed for the first time with the full Coulomb collision operator



Parameter scan
➔
Three collision operators




Electron Plasma Waves – Entropy Mode



Zero-frequency
entropy mode

- Electron-ion collisions known to yield long-time zero-frequency behaviour
- Full Coulomb collisions reduce the entropy mode damping rate

Jorge et al., to be submitted to JPP




First Study on Coulomb Collision Effects on the Drift-Wave Instability

- Two spatial and two velocity dimensions in slab geometry
- Electron and ion perturbations
- Finite background density gradient


❑ Important deviations between full Coulomb collision operator and presently used collision operators

Jorge *et al.*, PRL **121**, 165001 (2018)

Eigenmode Spectra



Instability Growth Rate



Summary

	This Work	Qin et al. 2007 ¹	Hahm et al. 2009 ²	Dimits et al. 2012 ³	Madsen et al. 2013 ⁴	Mandell et al. 2018 ⁵
Large Scales	EM $O(\epsilon^2)$	EM $O(\epsilon)$	ES $O(\epsilon^2)$	ES $O(\epsilon^2)$	ES $O(\epsilon)$	ES $O(\epsilon)$
Small Scales	EM $O(\epsilon^2)$	EM $O(\epsilon^2)$	EM $O(\epsilon^2)$	EM $O(\epsilon^2)$	EM $O(\epsilon)$	ES $O(\epsilon)$
Collisions	Yes	No	No	No	No	Simplified
Poisson's Eq.	Moments	$\int (\dots) d^3v$	Long-Wavelength Limit	$\int (\dots) d^3v$	Long-Wavelength Limit, Padé	Moments
$B_{ }^*$	Exact	Exact	Exact	Exact	$O(\epsilon)$	B

¹ Qin et al., Physics of Plasmas **14**, 056110 (2007)

² Hahm et al., Physics of Plasmas **16**, 022305 (2009)

³ Dimits, Physics of Plasmas **19**, 022504 (2012)

⁴ Madsen, Physics of Plasmas **20**, 072301 (2013)

⁵ Mandell et al., J. Plasma Phys **84**, 905840108 (2018)