Set up Default SIGs

Attach default SIGs to products and save time when dispensing by using the **k. sig** button in the Dispense Form. From **Dispense Pro**:

1. Click the Maintenance menu.

2. Select E. Default SIGS.

The Default SIGS window is displayed.

ocked	Name	Default SIGS	
1	! DISPENSE UPDATED 01-07-20		Change
N	2IN1-SMART-GLUCOSE TEST-STRP 50×1		
N	3TC 0-SOLN 10MG/ML 240MLX1		Delete
N	3TC TAB 150MG 60		Delete
N	3TC TAB 300MG 30		Ded
N	A-CITALOPRAM TAB 20MG 28		<u>r</u> ina
N	ABACAVIR/LAMIVDNE-MYLAN TAB 600/300MG 30		
N	ABACAVIR/LAMIVUDINE-GH TAB 600/300MG 30		
N	ABACAVIR/LAMIVUDINE-SUN TAB 600/300MG 30		
N	ABACAVIR/LAMVDNE-APOTEX TAB 600/300MG 30		
N	ABBOCILLIN-V DULCET TAB-CHEW 125MG 25 #D		
N	ABBOCILLIN-V SUSP 125MG/5ML 100ML 1 #DIS		
N	ABBOCILLIN-V SUSP 150MG/5ML 100MLX1		
N	ABBOCILLIN-VK CAP 250MG 25 #DISC		
N	ABBOCILLIN-VK-FILMTAB TAB 250MG 25		
N	ABBOCILLIN-VK-FILMTAB TAB 500MG 25		
N	ABELCET VIAL 100MG-20ML 10		
Y	ABILIFY TAB 10MG 30		
Y	ABILIFY TAB 15MG 30		
N	ABILIFY TAB 20MG 30		
N	ABILIFY TAB 30MG 30		Next
N	ABILIFY TAB 5MG 30		
N	ABILIFY-MAINTENA INJ 400MG 1		Prev
N	ABILIFY-MAINTENA INJ 300MG 1		
N	ABISART TAB 150MG 30		Cancel

3. Click Find to search for the product.

The **Search** window is displayed.

Search				
Enter Product Name				
<u> </u>				
Okay Cancel				

4. Enter the name of the product you wish to add a default SIG too. The product must be marked as dispensable. Refer to <u>Mark Products as Dispensable (https://help.minfos.com.au/en/support/solutions/articles/51000081665)</u> for further guidance.

5. Click Okay.

6. Highlight the correct product and click Change to update or add a default SIG.

7. The Default SIG Edit window is displayed.

	Default SIG Edit				
	Please enter the default SIG you would like to appear for this drug. This is activated by pressing [ALT + K] in dispense.				
	EGO-DERMAID CRM 1% 30G 1				
A					
	Okay Cancel				

A. Enter the previously added SIG code, e.g. AAA3D

8. Press Tab to update the SIG.

Default SIG Edit					
Please enter the default SIG you would like to appear for this drug. This is activated by pressing [ALT + K] in dispense.					
EGO-DERMAID CRM 1% 30G 1 Appy to the affected area three times a day.					
AAA3D					
	Okay Cancel				

The Default SIG Edit window will update with the directions.

9. Click Okay to save the default SIG.

You are returned to the **Default SIGS** window, and the Default SIG has been updated.

10. Repeat Steps 3 to 9 for any other products that require a default SIG.

11. Click Cancel to close the Default SIGs window.

The

<u>k</u>. Sig

option will be available when dispensing for any product with a Default SIG attached.