

Fig. 4. HSUR 1 down-regulates mature miRNAs in a sequence-specific and binding-dependent manner. **(A)** Partial sequences of HSUR 1 (positions 40 to 62) and its mutants (in red) H1Mt and H1m20. Bold nucleotides are perfectly conserved (Fig. 1A). **(B)** Coimmunoprecipitation of miRNAs with α Sm, as in Fig. 1C, from extracts of Jurkat T cells stably expressing HSURs 2 to 7 and either wild-type HSUR 1 (Wt, lanes 1 to 5), no HSUR 1 (Δ H1, lanes 6 to 10), mutant HSUR 1 H1Mt (lanes 11 to 15), or mutant HSUR 1 H1m20 (lanes 16 to 20). **(C)** miRNA levels in Jurkat T cells fluorescence-activated cell sorted for GFP after transient transfection with empty vector (GFP) or with plasmids expressing GFP and the following: HSUR 3 (GFP-H3), Wt HSUR 1 (GFP-H1), H1Mt (GFP-H1Mt), H1m20 (GFP-H1m20), or H1M1 (GFP-H1M1).

between HSURs 1 and 2 and miR-16 and miR-142-3p requires further investigation.

It is not yet clear how down-regulation of miR-27 benefits HVS. Down-regulation of the same host miRNA has been reported for another herpesvirus, murine cytomegalovirus, upon infection of cell lines and primary macrophages apparently also at the posttranscriptional level (23). Only a few targets of miRNA-27, including the transcription factors FOXO1, RUNX1 and PAX3, have been validated (18, 24, 25). Thus, identification of additional targets of miR-27 in T cells transformed with HVS is needed, as well as elucidation of the molecular mechanism by which association with HSUR 1 leads to miR-27 decay.

References and Notes

1. A. Ensser, B. Fleckenstein, *Adv. Cancer Res.* **93**, 91 (2005).
2. B. Biesinger, J. J. Trimble, R. C. Desrosiers, B. Fleckenstein, *Virology* **176**, 505 (1990).
3. S. I. Lee, S. C. Murthy, J. J. Trimble, R. C. Desrosiers, J. A. Steitz, *Cell* **54**, 599 (1988).
4. D. A. Wassarman, S. I. Lee, J. A. Steitz, *Nucleic Acids Res.* **17**, 1258 (1989).
5. J. C. Albrecht, *J. Virol.* **74**, 1033 (2000).
6. A. Ensser, A. Pfänder, I. Müller-Fleckenstein, B. Fleckenstein, *J. Virol.* **73**, 10551 (1999).
7. S. C. Murthy, J. J. Trimble, R. C. Desrosiers, *J. Virol.* **63**, 3307 (1989).
8. H. L. Cook, H. E. Mischo, J. A. Steitz, *Mol. Cell. Biol.* **24**, 4522 (2004).
9. X. C. Fan, V. E. Myer, J. A. Steitz, *Genes Dev.* **11**, 2557 (1997).
10. V. E. Myer, S. I. Lee, J. A. Steitz, *Proc. Natl. Acad. Sci. U.S.A.* **89**, 1296 (1992).
11. H. L. Cook *et al.*, *Curr. Biol.* **15**, 974 (2005).
12. P. Landgraf *et al.*, *Cell* **129**, 1401 (2007).
13. G. D. Cimino, H. B. Gamper, S. T. Isaacs, J. E. Hearst, *Annu. Rev. Biochem.* **54**, 1151 (1985).
14. M. Lagos-Quintana, R. Rauhut, W. Lendeckel, T. Tuschl, *Science* **294**, 853 (2001).
15. V. N. Kim, J. Han, M. C. Siomi, *Nat. Rev. Mol. Cell Biol.* **10**, 126 (2009).
16. H. W. Hwang, E. A. Wentzel, J. T. Mendell, *Science* **315**, 97 (2007).
17. Materials and methods are available as supporting material on Science Online.

18. I. K. Gutilla, B. A. White, *J. Biol. Chem.* **284**, 23204 (2009).
19. T. Ideue, K. Hino, S. Kitao, T. Yokoi, T. Hirose, *RNA* **15**, 1578 (2009).
20. M. S. Ebert, J. R. Neilson, P. A. Sharp, *Nat. Methods* **4**, 721 (2007).
21. A. Cimmino *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **102**, 13944 (2005).
22. Q. Liu *et al.*, *Nucleic Acids Res.* **36**, 5391 (2008).
23. A. H. Buck *et al.*, *RNA* **16**, 307 (2010).
24. O. Ben-Ami, N. Pencovich, J. Lotem, D. Levanon, Y. Groner, *Proc. Natl. Acad. Sci. U.S.A.* **106**, 238 (2009).
25. C. G. Crist *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **106**, 13383 (2009).
26. We thank A. Giraldez for initial bioinformatic searches, R. C. Desrosiers for plasmids and cell lines, R. Jobava

for technical assistance, K. Tycowski and K. Riley for critical commentary, and A. Miccinello for editorial assistance. This work was supported by grant CA16038 from the NIH. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. J.S. is an investigator at the Howard Hughes Medical Institute.

Supporting Online Material

www.sciencemag.org/cgi/content/full/328/5985/1563/DC1
 Materials and Methods
 Figs. S1 to S11
 Table S1
 References
 18 January 2010; accepted 14 April 2010
 10.1126/science.1187197

MicroRNA-33 and the SREBP Host Genes Cooperate to Control Cholesterol Homeostasis

S. Hani Najafi-Shoushtari,^{1,2} Fjoralba Kristo,³ Yingxia Li,⁴ Toshi Shioda,¹ David E. Cohen,⁴ Robert E. Gerszten,^{3,5} Anders M. Näär,^{1,2*}

Proper coordination of cholesterol biosynthesis and trafficking is essential to human health. The sterol regulatory element-binding proteins (SREBPs) are key transcription regulators of genes involved in cholesterol biosynthesis and uptake. We show here that microRNAs (miR-33a/b) embedded within introns of the SREBP genes target the adenosine triphosphate-binding cassette transporter A1 (ABCA1), an important regulator of high-density lipoprotein (HDL) synthesis and reverse cholesterol transport, for posttranscriptional repression. Antisense inhibition of miR-33 in mouse and human cell lines causes up-regulation of ABCA1 expression and increased cholesterol efflux, and injection of mice on a western-type diet with locked nucleic acid-antisense oligonucleotides results in elevated plasma HDL. Our findings indicate that miR-33 acts in concert with the SREBP host genes to control cholesterol homeostasis and suggest that miR-33 may represent a therapeutic target for ameliorating cardiometabolic diseases.

Cholesterol and other lipids play key roles in many physiological processes in metazoans, and aberrant cholesterol/lipid homeostasis has been linked to a number of diseases,

including atherosclerosis, metabolic syndrome, and type II diabetes, underscoring the importance of understanding fully how cholesterol/lipid homeostasis is regulated (1, 2).

Downloaded from https://www.science.org at University of California Berkeley on July 26, 2022

The sterol regulatory element-binding protein (SREBP) transcription factors are critical regulators of cholesterol/lipid homeostasis, which act by controlling the expression of many cholesterologenic and lipogenic genes [e.g., low-density lipoprotein (LDL) receptor, 3-hydroxy-3-methylglutaryl coenzyme A reductase, and fatty acid synthase] (3–6). Although much is known about SREBP-dependent transcriptional mechanisms governing the biosynthesis and uptake of cholesterol and fatty acids, it is unclear how this regulatory circuit is coordinated with opposing pathways that mediate cholesterol/lipid efflux or degradation to achieve appropriate cholesterol/lipid levels to satisfy cellular and physiological demands.

In addition to classical transcription regulators, a class of noncoding RNAs termed microRNAs (miRNAs) has emerged as important modulators of numerous cellular processes that affect organism growth, development, homeostasis, and disease (7–12). Indeed, recent studies have revealed that a liver-restricted miRNA, miR-122, regulates cholesterol/lipid metabolism in mice and nonhuman primates, although the mechanism remains unclear (13–16). Data from these studies not only emphasize the important roles played by miRNAs in normal physiology but also point to the feasibility of antisense-based therapeutic targeting of miRNAs to treat human disease.

During our investigations of gene regulation by SREBPs (17), we noted the intriguing presence of a highly conserved miRNA family, miR-33, within intronic sequences of the SREBP genes in organisms from *Drosophila* to humans (Fig. 1, A and B, and fig. S1). Two isoforms of miR-33 exist in humans: miR-33b, which is present in intron 17 of the SREBP-1 gene on chromosome 17, and miR-33a, which is located in intron 16 of the SREBP-2 gene on chromosome 22 (Fig. 1, A and B). In mice, however, there is only one miR-33 isoform (which is conserved with human miR-33a), located within intron 15 of the mouse SREBP-2 gene, whereas intron 17 of the mouse SREBP-1 gene lacks sequence homology to the human intronic sequences harboring miR-33b (fig. S1).

Similar to miR-33, many mammalian miRNAs are located within introns of protein-coding genes rather than in their own unique transcription units (18). Intronic miRNAs are typically coordinately expressed and processed with the precursor mRNA in which they reside (19–21). Accord-

ingly, the mature forms of miR-33a/b appear to be coexpressed with the SREBP host genes in a number of human and mouse tissues examined (Fig. 1, C to E, and figs. S2 to S4).

We wished to determine the potential function(s) of miR-33a/b and whether they exhibit functional association with the SREBP host genes. miRNAs have been shown to target mRNAs for posttranscriptional repression by base-pairing with mRNA sequences typically located in the 3' untranslated regions (3'UTRs) and causing translational inhibition or mRNA cleavage (22). We initially employed commonly used bioinformatics tools that predict miRNA targets largely based on the ability of the miRNA sequence to undergo specific base-pairing with its putative 3'UTR target, known as “seed pairing” (22). The most prominent predicted conserved target for miR-33a/b among vertebrates is the adenosine triphosphate-binding cassette A1 (ABCA1) cholesterol transporter (table S1 and fig. S5A). ABCA1 is a key mediator of intracellular cholesterol efflux from liver to apolipoprotein A-I (apoA-I) for generation of high-density lipoprotein (HDL) (1, 23, 24). It is also important for HDL-cholesterol trafficking from peripheral tissues (e.g., macrophages) by the reverse cholesterol transport (RCT) pathway back to the liver for processing and excretion into bile and feces (25).

As SREBPs promote cholesterol uptake and synthesis through transactivation of the LDL receptor and cholesterol biosynthesis genes, we hypothesized that miR-33-mediated inhibition of ABCA1 and cholesterol efflux could potentially act in cooperation with SREBPs to boost intracellular cholesterol levels. To test this hypothesis, we first carried out RNA interference-mediated knockdown of components of the miRNA biogenesis pathway to determine whether ABCA1 protein expression is regulated by miRNAs. Indeed, transfection with siRNAs directed against the Drosha and Dicer miRNA processing enzymes resulted in a marked increase in ABCA1 protein expression in several human and mouse cell lines, including human HepG2 liver carcinoma cells, IMR-90 normal human fibroblasts, and the mouse macrophage cell line J774 (Fig. 2A). These results are thus consistent with regulation of ABCA1 by miRNAs. To evaluate the specific effect of miR-33 on ABCA1 expression, we transfected cells with synthetic miR-33 precursor oligonucleotides (pre-miR-33a/b) to increase the intracellular levels of miR-33a and b, respectively. These studies show that ABCA1 expression is repressed by excess miR-33, especially in mouse J774 macrophages and the human IMR-90 fibroblasts (Fig. 2B). To further determine whether miR-33a/b are specifically involved in regulating




Fig. 1. SREBPs are host genes to conserved intronic miRNAs, miR-33a/b, which are coexpressed with SREBPs. Human SREBP-1 (A) and SREBP-2 (B) genes harbor related intronic miRNAs (miR-33b and miR-33a, respectively). The sequences encoding the pre-miRNAs are shown, with the mature miRNA sequences highlighted in red. (C to E) Expression profile of miR-33a/b and SREBP host genes in selected human tissues. Error bars represent experimental SD.

¹Massachusetts General Hospital Cancer Center, Charlestown, MA 02129, USA. ²Department of Cell Biology, Harvard Medical School, Boston, MA 02115, USA. ³Center for Immunology and Inflammatory Diseases, Massachusetts General Hospital, Charlestown, MA 02129, USA. ⁴Department of Medicine, Division of Gastroenterology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA 02115, USA. ⁵Cardiovascular Research Center, Massachusetts General Hospital, Charlestown, MA 02129, USA.

*To whom correspondence should be addressed. E-mail: naar@helix.mgh.harvard.edu

ABCA1 expression, we transfected the three cell lines with antisense oligonucleotides directed against miR-33a and/or b (anti-miR-33a/b). Introduction of anti-miR-33a/b indeed resulted in strongly increased levels of ABCA1 in the three cell lines (Fig. 2C). Taken together, these findings demonstrate that ABCA1 protein expression is regulated by miR-33a/b.

To determine whether miR-33a/b specifically target ABCA1 posttranscriptionally through its 3'UTR, we fused a fragment of the human ABCA1 3'UTR harboring the predicted miR-33 target sequences to a Luciferase reporter plasmid (fig. S5B). As shown in Fig. 2D, the Luciferase-ABCA1 3'UTR reporter was expressed at markedly lower levels in human embryonic kidney (HEK) 293 cells as compared with the Luciferase vector without insert. Introduction of excess wild-type human miR-33a/b precursor oligonucleotides resulted in further repression of the wild-type ABCA1 3'UTR reporter (Fig. 2E). By contrast, miR-33a/b precursors mutated in the seed base-pairing sequence had no effect, as compared with precursor control (PC) (Fig. 2E and fig. S5C). Together, these data show that miR-33a/b are capable of repressing expression from the ABCA1 3'UTR. Importantly, cotransfection with miR-33a/b antisense oligonucleotides caused complete derepression of the Luciferase-ABCA1 3'UTR reporter, consistent with regulation of ABCA1 3'UTR by endogenous miR-33a/b (Fig. 2F). Finally, mutations in the seed base-pairing sequences of the predicted miR-33 target sites in the ABCA1 3'UTR reporter abolished regulation by miR-33a/b precursors as well as anti-miR-33a/b (Fig. 2G). Together, these results suggest that miR-33a/b inhibit expression of human ABCA1 by targeting the ABCA1 3'UTR for translation repression or mRNA degradation.

Reverse cholesterol transport from atherogenic macrophages/foam cells is of clinical importance because of the prominent role played by foam cells in cardiovascular disease, and a number of studies have highlighted the biomedical implications of ABCA1-dependent cholesterol efflux and HDL biogenesis for atherosclerosis (23, 26–28). We have investigated the potential biomedical and physiological relevance of the miR-33/SREBP/ABCA1 cholesterol regulatory circuit using the mouse J774 macrophage model (29). SREBPs are regulated in a classic negative feedback manner by cholesterol (3, 5), and we find that depletion of cholesterol by lovastatin/ β -cyclodextrin treatment results in increased expression of both miR-33a and the mSREBP-2 host gene in J774 macrophages, with a concomitant decrease in ABCA1 protein expression (Fig. 3, A to C). The strong suppression of ABCA1 protein levels in response to cholesterol depletion is at least partially reversed by miR-33a antisense oligonucleotides (Fig. 3D), consistent with the notion that miR-33a mediates cholesterol-regulated posttranscriptional control of ABCA1 levels in mouse macrophages. We also found that the high protein levels of ABCA1 observed in J774 macrophages cultured in the presence of





Fig. 2. miR-33 regulates the cholesterol transporter ABCA1. (A) Transfection of siRNA against Drosha and Dicer results in increased protein levels of ABCA1 in human HepG2 hepatoma cells (top row), human IMR-90 fibroblasts (middle row), and the mouse macrophage cell line J774 (bottom row). (B) Introduction of miR-33a or miR-33b precursors into the three cell lines represses ABCA1 protein levels. (C) miR-33a/b antisense oligonucleotides (A33a/b) increase ABCA1 levels. (D) Schematic representation of the Luciferase reporter vector harboring the ABCA1 3'UTR fragment (green) containing the two predicted miR-33 target sites. (E) Insertion of the ABCA1 3'UTR fragment into a Luciferase reporter results in decreased Luciferase expression in HEK293 cells. (F) Cotransfection with wild-type, but not mutated, miR-33a/b precursors causes further repression of the Luciferase-ABCA1 3'UTR reporter. A scrambled sequence was used as PC. (G) miR-33a/b antisense oligonucleotides result in derepression of the Luciferase-ABCA1 3'UTR reporter. A scrambled sequence was used as AC. Error bars represent SD. *, $P < 0.05$.


Fig. 3. Reciprocal regulation by cholesterol and SREBP/miR-33. Lovastatin/ β -cyclodextrin-mediated depletion of cholesterol in J774 mouse macrophages results in increased expression of (A) SREBP-2 and (B) miR-33a, and (C) in decreased ABCA1 protein levels. (D) Cholesterol-depleted J774 mouse macrophages exhibit up-regulation of ABCA1 protein in response to miR-33a antisense inhibition. (E) Transfection of J774 macrophages cultured in the presence of serum/cholesterol with miR-33a precursor results in decreased levels of ABCA1. (F) Introduction of miR-33a antisense oligonucleotides into J774 macrophages loaded with radio-labeled cholesterol results in increased cholesterol efflux, whereas miR-33a precursors inhibit cholesterol efflux. Error bars represent SD. **, $P < 0.01$.



serum/cholesterol are substantially suppressed by additional miR-33a precursor (Fig. 3E), in keeping with miR-33 regulation of ABCA1. By contrast with the apparent regulation of ABCA1 levels in

J774 cells by miR-33a, expression of the intracellular cholesterol transporter NPC1 is unaffected by miR-33a precursor or antisense treatment (fig. S6), demonstrating specific regulation of ABCA1

Fig. 4. Regulation of HDL by miR-33a in vivo. **(A)** Tail-vein injection of mice fed a western-type diet with LNA-antisense oligonucleotides directed against mouse miR-33a results in elevated total serum cholesterol. **(B)** Fast protein liquid chromatography analysis of pooled serum from five mice on a western-type diet revealed an increase in the magnitude of the HDL-cholesterol peak in the LNA-antimiR-treated animals. Elution of lipoprotein standards is indicated by the labels. Plasma triglycerides **(C)**, glucose **(D)**, AST **(E)**, and ALT **(F)** were unaffected by the treatments. Error bars represent SEM. **, $P < 0.01$.



by miR-33a. Our data together show that mouse miR-33a and the SREBP-2 host gene are coregulated by cholesterol and suggest the existence of a reciprocal regulatory network of SREBP, miR-33, ABCA1, and cholesterol in macrophages.

Next, we examined the effects of miR-33 manipulations on cholesterol efflux from J774 macrophages. Cells were first labeled with [³H]-cholesterol and then transfected with miR-33a precursor, miR-33a antisense, or control oligonucleotides, followed by measurements of the level of [³H]-cholesterol efflux to serum/apoA-I. As shown in Fig. 3F, introduction of miR-33a antisense oligonucleotides into mouse macrophages caused a marked (two-fold) increase in [³H]-cholesterol efflux, whereas miR-33a precursor treatment led to a significant decrease in [³H]-cholesterol efflux, as compared with control oligonucleotides [antisense control (AC)/PC] or untransfected cells (UT). These findings demonstrate that miR-33 posttranscriptional repression of ABCA1 results in retention of intracellular cholesterol in macrophages and are in keeping with our hypothesis that miR-33 regulates cholesterol trafficking by controlling ABCA1 levels.

Based on our in vitro findings, we hypothesized that manipulation of ABCA1 levels by miR-33 antisense approaches might lead to augmented HDL-cholesterol levels in a mammalian in vivo model. Previous studies in mice and nonhuman primates have supported the feasibility of locked nucleic acid (LNA) antisense approaches to modulate miRNA pathways with important physiological consequences in vivo (13, 14). We therefore performed

tail-vein injections of mice on a western-type diet with phosphate-buffered saline (vehicle), LNA-miR-33a antisense, or LNA-control oligonucleotides. After three injections over 5 days, mice were killed and serum was collected. We find that plasma HDL-cholesterol concentrations were significantly increased in LNA-miR-33a antisense-treated animals as compared with LNA-control-treated mice (Fig. 4, A and B, and table S2). By contrast, there were no significant effects on plasma concentrations of LDL cholesterol, triglycerides, or glucose levels (Fig. 4, C to F, and table S2). In these experiments, there was also no apparent hepatotoxicity [as measured by plasma aspartate aminotransferase/alanine aminotransferase (AST/ALT)] (Fig. 4, E and F, and table S2). The observed elevation in plasma HDL cholesterol in response to LNA-miR-33a antisense treatment is consistent with regulation of ABCA1-dependent cholesterol efflux by miR-33a in vivo. Hepatic ABCA1 is a major contributor to HDL production in normal mice (23, 24); however, we find only modest effects of miR-33a LNA-antimiR on hepatic ABCA1 mRNA/protein as compared with LNA-control (fig. S7). Based on these findings, we speculate that liver might not be the sole target tissue in mediating the effect of miR-33a LNA-antimiR on HDL production, with possible additional contribution of miR-33a regulation of reverse cholesterol transport from extrahepatic tissues/cells.

We have shown that members of the mammalian SREBP family of transcription factors, key regulators of cholesterologenic and lipogenic

genes, are hosts to conserved miRNAs (miR-33a/b) that may function in concert with the SREBP host gene products to govern intracellular cholesterol levels and cholesterol homeostasis in vertebrates (fig. S8). Indeed, our data reveal that miR-33 exerts posttranscriptional control of the ABCA1 cholesterol transporter, with important consequences for cholesterol trafficking in vitro and HDL synthesis in vivo. Our findings also point to the possibility of antisense therapeutic targeting of miR-33a/b as a strategy to increase HDL in individuals suffering from cardiometabolic diseases.

References and Notes

1. F. R. Maxfield, I. Tabas, *Nature* **438**, 612 (2005).
2. D. E. Moller, K. D. Kaufman, *Annu. Rev. Med.* **56**, 45 (2005).
3. M. S. Brown, J. L. Goldstein, *J. Lipid Res.* **50**, (Suppl), S15 (2009).
4. T. F. Osborne, P. J. Espenshade, *Genes Dev.* **23**, 2578 (2009).
5. M. S. Brown, J. L. Goldstein, *Cell* **89**, 331 (1997).
6. J. D. Horton, J. L. Goldstein, M. S. Brown, *J. Clin. Invest.* **109**, 1125 (2002).
7. A. H. Williams, N. Liu, E. van Rooij, E. N. Olson, *Curr. Opin. Cell Biol.* **21**, 461 (2009).
8. M. V. Iorio, C. M. Croce, *J. Clin. Oncol.* **27**, 5848 (2009).
9. D. P. Bartel, *Cell* **116**, 281 (2004).
10. L. He, G. J. Hannon, *Nat. Rev. Genet.* **5**, 522 (2004).
11. J. Krützfeldt, M. Stoffel, *Cell Metab.* **4**, 9 (2006).
12. G. Stefani, F. J. Slack, *Nat. Rev. Mol. Cell Biol.* **9**, 219 (2008).
13. J. Elmén *et al.*, *Nature* **452**, 896 (2008).
14. J. Elmén *et al.*, *Nucleic Acids Res.* **36**, 1153 (2008).
15. C. Esau *et al.*, *Cell Metab.* **3**, 87 (2006).
16. J. Krützfeldt *et al.*, *Nature* **438**, 685 (2005).
17. F. Yang *et al.*, *Nature* **442**, 700 (2006).
18. A. Rodríguez, S. Griffiths-Jones, J. L. Ashurst, A. Bradley, *Genome Res.* **14**, (10A), 1902 (2004).
19. D. Wang *et al.*, *PLoS ONE* **4**, e4421 (2009).
20. S. Baskerville, D. P. Bartel, *RNA* **11**, 241 (2005).
21. Y. K. Kim, V. N. Kim, *EMBO J.* **26**, 775 (2007).
22. D. P. Bartel, *Cell* **136**, 215 (2009).
23. A. R. Tall, L. Yuan-Charvet, N. Terasaka, T. Pagler, N. Wang, *Cell Metab.* **7**, 365 (2008).
24. X. Wang, D. J. Rader, *Curr. Opin. Cardiol.* **22**, 368 (2007).
25. D. J. Rader, E. T. Alexander, G. L. Weibel, J. Billheimer, G. H. Rothblat, *J. Lipid Res.* **50**, (Suppl), S189 (2009).
26. R. R. Singaraja *et al.*, *J. Clin. Invest.* **110**, 35 (2002).
27. X. Wang *et al.*, *J. Clin. Invest.* **117**, 2216 (2007).
28. M. Van Eck *et al.*, *Arterioscler. Thromb. Vasc. Biol.* **26**, 929 (2006).
29. M. de la Llera-Moya *et al.*, *Arterioscler. Thromb. Vasc. Biol.* **30**, 796 (2010).
30. We thank S. Vasudevan and A. Walker for critical reading of the manuscript and K. Coser for technical assistance. This work was supported by the following funding sources: NIH R01GM071449 and R21DK084459 (A.M.N.), R01DK56626, R01DK48873, and P30 DK34854 (D.E.C.), American Heart Association Established Investigator Award (R.E.G.), the Fondation Leducq (R.E.G.), and Massachusetts Biomedical Research Corporation Tosteson Fellowship Award (S.H.N.-S.). A.M.N. and Massachusetts General Hospital have filed a patent application relating to the therapeutic use of antisense oligonucleotides directed against miR-33.

Supporting Online Material

www.sciencemag.org/cgi/content/full/science.1189123/DC1
 Materials and Methods
 Figs. S1 to S8
 Tables S1 and S2
 References

4 March 2010; accepted 5 May 2010
 Published online 13 May 2010;
 10.1126/science.1189123
 Include this information when citing this paper.

Downloaded from https://www.science.org at University of California Berkeley on July 26, 2022

MicroRNA-33 and the SREBP Host Genes Cooperate to Control Cholesterol Homeostasis

S. Hani Najafi-Shoushtari, Fjoralba Kristo, Yingxia Li, Toshi Shioda, David E. Cohen, Robert E. Gerszten, and Anders M. Näär

Science, 328 (5985), • DOI: 10.1126/science.1189123

miR-33 in Cholesterol Control

With the well-established link between serum cholesterol levels and cardiovascular disease and the availability of effective cholesterol-lowering drugs, cholesterol screening has rapidly become a routine part of health care. Yet, much remains to be learned about how cholesterol levels are regulated at the cellular level (see the Perspective by Brown *et al.*). Now, Najafi-Shoushtari *et al.* (p. 1566, published online 13 May) and Rayner *et al.* (p. 1570, published online 13 May) have discovered a new molecular player in cholesterol control—a small noncoding RNA that, intriguingly, is embedded within the genes coding for sterol regulatory element-binding proteins (SREBPs), transcription factors already known to regulate cholesterol levels. This microRNA, called miR-33, represses expression of the adenosine triphosphate-binding cassette transporter A1, a protein that regulates synthesis of high-density lipoprotein (HDL, or “good” cholesterol) and that helps to remove “bad” cholesterol from the blood. Reducing the levels of miR-33 in mice boosted serum HDL levels, suggesting that manipulation of this regulatory circuit might be therapeutically useful.

View the article online

<https://www.science.org/doi/10.1126/science.1189123>

Permissions

<https://www.science.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of service](#)