## WEBBERVILLE 16% CALF GROWER RUM MEDICATED

For Starting Dairy Calves

For prevention and control of coccidiosis caused by Eimeria bovis and Eimeria zuemii in calves (excluding yeal).

ACTIVE DRUG INGREDIENT  Monensin (as Monensin Sodium)	36.6 g/ton
	16.00%
Crude Protein, minimum	2.00%
Crude Fat, minimum	5.75%
Crude Fiber, maximum.	6.90%
Acid Detergent Fiber (ADF), maximum.	
Calcium (Ca), minimum	0.50%
Calcium (Ca), maximum	1.00%
Selenium (Se), minimum	0.3
Vitamin A, minimum	2,500
Vitamin E, minimum	25.8

INGREDIENTS

Grain Products, Plant Protein Products, Processed Grain By-Products, Calcium Carbonate, Molasses Products, Salt, Magnesium Oxide, Ferrous Sulfate, Cobalt Carbonate, Copper Sulfate, Calcium Iodate, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Calcium Sulfate, Vitamin A Supplement, Vitamin D<sub>3</sub> Supplement, Vitamin E Supplement, Sodium Selenite

## **FEEDING DIRECTIONS**

Hand feed continuously to calves (excluding veal calves) to provide .14 to 1.00 mg monensin/lb bodyweight. Feed with good quality hay and provide plenty of fresh, clean water.



CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions.



Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing.



This feed contains copper. Do not feed to copper sensitive species.

WARNING: A withdrawal period has not been established for preruminating calves. Do not use in calves to be processed for veal.



NET WT. 50 lb (22.68 KG) or BULK