



**Chapter 11**  
**Infinite Sequences and Series**

 CENGAGE

Stewart, Calculus: Early Transcendentals, 8th Edition. © 2016 Cengage. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

1

---

---

---

---

---

---

---

---

**11.9 Representations of Functions as Power Series**

 CENGAGE

Stewart, Calculus: Early Transcendentals, 8th Edition. © 2016 Cengage. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

2

---

---

---

---

---

---

---

---

**Representations of Functions as Power Series (1 of 2)**

We start with an equation that we have seen before:

$$1 \quad \frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots = \sum_{n=0}^{\infty} x^n \quad |x| < 1$$

We have obtained it by observing that the series is a geometric series with  $a = 1$  and  $r = x$ .

But here our point of view is different. We now regard Equation 1 as expressing the function  $f(x) = \frac{1}{(1-x)}$  as a sum of a power series.

 CENGAGE

Stewart, Calculus: Early Transcendentals, 8th Edition. © 2016 Cengage. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

3

---

---

---

---


---

---

---

---

### Representations of Functions as Power Series (2 of 2)



$$f(x) = \frac{1}{1-x} \text{ and some partial sums}$$

Figure 1

CENGAGE

Stewart, Calculus: Early Transcendentals, 8th Edition, © 2016 Cengage. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

4

---

---

---

---

---

---

---

---

---

---

### Example 1

Express  $\frac{1}{(1+x^2)}$  as the sum of a power series and find the interval of convergence.

**Solution:**

Replacing  $x$  by  $-x^2$  in Equation 1, we have

$$\begin{aligned} \frac{1}{1+x^2} &= \frac{1}{1-(-x^2)} \\ &= \sum_{n=0}^{\infty} (-x^2)^n \end{aligned}$$

CENGAGE

Stewart, Calculus: Early Transcendentals, 8th Edition, © 2016 Cengage. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

5

---

---

---

---

---

---

---

---

---

---

### Example 1 – Solution

$$\begin{aligned} &= \sum_{n=0}^{\infty} (-1)^n x^{2n} \\ &= 1 - x^2 + x^4 - x^6 + x^8 - \dots \end{aligned}$$

Because this is a geometric series, it converges when  $| -x^2 | < 1$ , that is,  $x^2 < 1$ , or  $|x| < 1$ .

Therefore the interval of convergence is  $(-1, 1)$ . (Of course, we could have determined the radius of convergence by applying the Ratio Test, but that much work is unnecessary here.)

CENGAGE

Stewart, Calculus: Early Transcendentals, 8th Edition, © 2016 Cengage. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

6

---

---

---

---

---

---

---

---

---

---

## Differentiation and Integration of Power Series



Stewart, Calculus: Early Transcendentals, 8th Edition, © 2016 Cengage. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

7

---

---

---

---

---

---

---

---

### Differentiation and Integration of Power Series (1 of 2)

The sum of a power series is a function  $f(x) = \sum_{n=0}^{\infty} c_n (x-a)^n$  whose domain is the interval of convergence of the series.

We would like to be able to differentiate and integrate such functions, and the following theorem says that we can do so by differentiating or integrating each individual term in the series, just as we would for a polynomial.

This is called **term-by-term differentiation and integration**.



Stewart, Calculus: Early Transcendentals, 8th Edition, © 2016 Cengage. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

8

---

---

---

---

---

---

---

---

### Differentiation and Integration of Power Series (2 of 2)

**2 Theorem** If the power series  $\sum c_n (x-a)^n$  has radius of convergence  $R > 0$ , then the function  $f$  defined by

$$f(x) = c_0 + c_1(x-a) + c_2(x-a)^2 + \dots = \sum_{n=0}^{\infty} c_n (x-a)^n$$

is differentiable (and therefore continuous) on the interval  $(a-R, a+R)$  and

$$(i) f'(x) = c_1 + 2c_2(x-a) + 3c_3(x-a)^2 + \dots = \sum_{n=1}^{\infty} n c_n (x-a)^{n-1}$$

$$(ii) \int f(x) dx = C + c_0(x-a) + c_1 \frac{(x-a)^2}{2} + c_2 \frac{(x-a)^3}{3} + \dots$$

$$= C + \sum_{n=0}^{\infty} c_n \frac{(x-a)^{n+1}}{n+1}$$

The radii of convergence of the power series in Equations (i) and (ii) are both  $R$ .



Stewart, Calculus: Early Transcendentals, 8th Edition, © 2016 Cengage. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

9

---

---

---

---

---

---

---

---

**Example 4**

We have seen the Bessel function

$$J_0(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{2^{2n} (n!)^2}$$

is defined for all  $x$ .

Thus, by Theorem 2,  $J_0$  is differentiable for all  $x$  and its derivative is found by term-by-term differentiation as follows:

$$J_0'(x) = \sum_{n=0}^{\infty} \frac{d}{dx} \frac{(-1)^n x^{2n}}{2^{2n} (n!)^2} = \sum_{n=1}^{\infty} \frac{(-1)^n 2nx^{2n-1}}{2^{2n} (n!)^2}$$



Stewart, Calculus: Early Transcendentals, 8th Edition, © 2016 Cengage. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

---



---



---



---



---



---



---