

PLEASE READ THE LABEL COMPLETELY BEFORE USE

WARNING

KEEP OUT OF REACH OF CHILDREN

TALON[®] WB

RODENTICIDE

Active Ingredient: TALON WB contains 0.05 g/kg BRODIFACOUM in the form of a bait.

Ready to use block bait for use in damp or dry situations in and around industrial, commercial, public services, agricultural and domestic buildings.

Pack size: 1 kg, 2.4 kg and 10 kg

Registered pursuant to the ACVM Act 1997, No. V9229.

See www.foodsafety.govt.nz/industry/acvm/ for registration conditions.

Approved pursuant to the HSNO Act 1996, Approval Code. HSR001594.

Syngenta Crop Protection Limited

Tower 2, Level 7, 110 Symonds Street, Auckland

In a transport emergency dial 111, Police or Fire Brigade. For specialist advice in an emergency only, call 0800 734 607 (24 hours).

UN-Free

This product **MUST NOT** be used for any purpose or in any manner contrary to the controls and classifications of this registration.

10 kg pack only:

MUST only be sold by persons approved by the ACVM group and a register of sales MUST be kept for a minimum of 3 years.

HSNO Classifications 6.1E, 6.9B, 9.1D

WARNING

May be harmful if swallowed.

May cause damage to the blood and hematopoietic system from repeated oral exposure at high doses.

Harmful to aquatic organisms.

Harmful to terrestrial vertebrates.

PRECAUTIONS

DO NOT store near food or feed.

Avoid skin contact.

For use only in bait stations.

DO NOT lay in bush areas where stock or feral animals have access to baits.

DO NOT lay baits in places accessible to children, pets and domestic animals.

Wash hands and exposed skin after handling.

Dispose of carcasses by burial or by wrapping in paper, placing in plastic bag and putting in garbage.

Dispose of empty container and unused or untaken baits by wrapping in paper, placing in plastic bag and putting in garbage.

DO NOT contaminate drains, sewers, streams or ponds with the pellet bait or its used container.

Container disposal: Empty containers should be burnt if circumstances, especially wind direction, permit. Otherwise dispose of empty container by wrapping in paper and placing in garbage.

Product disposal: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

FIRST AID

If swallowed DO NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

Treatment of Domestic Animals: Depending on the amount eaten, this bait may kill or injure dogs, cats, pigs, poultry and wildlife. DO NOT place baits in locations which are accessible to pets, domestic animals, livestock or birds. DO NOT contaminate dams, waterways and drains with the bait or its used container. Vitamin K1 is an effective antidote against accidental poisoning of domestic animals by TALON WB.

Practising veterinarians have been provided with information on Vitamin K1 and should be consulted.

Note to Physicians and Veterinarians: Brodifacoum, the active constituent of TALON WB, is a long lasting anticoagulant chemical, which if ingested by humans, domestic animals or pets, can reduce the clotting power of the blood and haemorrhage may result. Vitamin K1 (Phytomenadione) should be administered urgently by medical, appropriate para-medical personnel or veterinarians. If administered intravenously the injection must be given slowly.

GENERAL INFORMATION

TALON WB is formulated for use indoors or outdoors. It will control rats and mice, including those that may be resistant to Warfarin. Pre-baiting is not required as individual rodents can readily consume a lethal dose at a single feeding. Most deaths occur 4 to 7 days after consumption of bait. Wax blocks are particularly suitable for use in damp situations and sewers, and crop stores where the risk of contamination should be minimised.

STORAGE AND DISPOSAL

Containers which have held bait should not be used for any other purpose. Store in the closed, original container in a cool, dry place out of the reach of children. DO NOT store in direct sunlight. Dispose of empty containers and unused or untaken baits by burying in an approved dump. Look for dead rats or mice, burn or bury them. Ensure baits are removed when rodent activity has ceased. See Safety Data Sheet for additional storage information. Stores containing quantities of 10,000 kg or more require signage and secondary containment.

SAFETY DATA SHEET

If additional hazard information is required, refer to the Safety Data Sheet. For a copy, visit our website at www.syngenta.co.nz

MAF approved 1.4.92 for use in Dairy Factories Category B and C.

NZFSA Approved Type C (All animal product except dairy).

DISCLAIMER

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warrant is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

®TALON, the SYNGENTA logo and the CP frame are trademarks of a Syngenta Group Company



Distributed by and registered to Syngenta Crop Protection Limited, Tower 2, Level 7, 110 Symonds Street, Auckland.

DIRECTIONS FOR USE

Situation	Pest	Rate	Critical Comments
Industrial, commercial, public services, agricultural and domestic buildings	Rats and mice	1 to 3 blocks/ bait station	<p>Set out baits in places frequented by mice and rats.</p> <p>For rats: Place 1 to 3 blocks at intervals of 5 to 9 m in infested areas.</p> <p>For mice: Place single blocks at 2 to 3 m intervals in infested areas.</p> <p>DO NOT exceed 3 m between bait stations for mice or 9 m for rats.</p> <p>DO NOT place bait stations in the open.</p> <p>Ensure baits are inaccessible to children, pets, domestic animals and wildlife.</p> <p>If baits are placed outside where members of the public ordinarily have access signs must be erected at every normal point of entry before the bait is laid and must remain for 12 months after removal of the bait.</p>

Baiting strategy

- Eliminate as far as practical, all alternative food sources.
- Locate baits under cover, along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail in place if relevant. Use with an appropriate bait station is recommended.
- Record the number and location of bait stations for the purpose of replenishment and later removal.
- Inspect baits regularly.

For rats: Use 1 to 3 blocks depending on the level of infestation.

For mice: Use single blocks.

- Replace taken baits every 7 days until no more baits are taken. If taken quickly, increase number of baits. Baiting for at least 2 weeks will be necessary to reduce rat/mouse numbers.
- Ensure baits are removed when rodent activity has ceased.

Note: One feed is sufficient to kill rats and mice. Rats and mice will continue to eat the bait after a lethal dose has been eaten.