

# The first gapless, reference-quality, fully annotated genome from a Southern Han Chinese individual




JOHNS HOPKINS UNIVERSITY CENTER FOR COMPUTATIONAL BIOLOGY

**CCB** 

Kuan-Hao Chao<sup>1,2</sup>, Aleksey V Zimin<sup>2,3</sup>, Mihaela Pertea<sup>2,3</sup>, Steven L Salzberg<sup>1,2,3,4</sup> <sup>1</sup>Department of Computer Science, <sup>2</sup>Center for Computational Biology, <sup>3</sup>Department of Biomedical Engineering, <sup>4</sup>Department of Biostatistics, Johns Hopkins University

github.com/JHUCCB/ChineseHanSouthGenome



G3 doi.org/10.1093/g3journal/jkac321

kh.chao@cs.jhu.edu, salzberg@jhu.edu




# Introduction


- Used <u>HiFi + ONT long reads</u> to assemble the genome of a Southern Han Chinese male
- Filled in gaps using T2T-CHM13 as a guide. *Han1* is gap-free
- Han1 contains <u>3,099,707,698 bases</u>
- Annotated Han1 using Liftoff: **60,708 putative genes** (20,003 protein coding genes)
- The first gene-level comparison between two finished, annotated individual human

## Han1 / T2T-CHM13 Assembly Comparison

Collinearity (MUMmer4)



• Inversion on β defensin gene cluster (Plotsr)



## **Genome Assembly**

#### Sequence data

Sequence type	Basecaller	Number of reads	N50 read length (bp)	Mean read length (bp)	Maximum length	Total length (Gbp)	Genome coverage
PacBio HiFi	ccs v4.0.0	5,570,675	21,499	21,989	50,388	122.50	39.45x
ONT Ultralong	Guppy v4.0.11	<b>3,110,293</b> (9.11% > 100 Kb)	83,822	34,136	2,495,296	106.17	34.75x

#### **Draft HiFi and ONT Assemblies**

Assembler	Sequencing data	Assembled sequence (bp)	Contig N50	Number of contigs	Quality value
Hifiasm v0.16.1-r375	PacBio HiFi	3,110,501,483	95,769,069	182	57.8 <sup>1</sup>
Flye v2.5	ONT Ultralong	2,974,205,132	40,850,737	1,658	25.6 <sup>1</sup>

#### Han1 Assembly

- **Primary:** Hifiasm assembly
- Secondary: Flye Assembly
- **Reference:** T2T-CHM13
- Fixing mis-assembled contigs
- **Closing Gaps** on primary assembly
- 1. HiFi reads 2. Fly assembly 3. T2T-CHM13

# **Gene Annotation**

### GRCh38 / T2T-CHM13 / Han1 Gene Count Comparison

• Two-pass annotation lift-over from T2T-CHM13 to Han1 using Liftoff


Gene biotype	Count in GRCh38	Count in T2T-CHM13	Count in Han1
protein coding	19,871	20,022	20,003
lncRNA	17,793	18,389	18,321
pseudogene	15,357	16,027	15,881
miRNA	1,914	2,046	2,058
transcribed pseudogene	1,221	1,262	1,232
rRNA	38	765	658
other	2,506	2,629	2,556
Total	58,700	61,140	60,708


### Han1 / T2T-CHM13 Gene Content Comparison

• LiftoffTools: <sup>(1)</sup> Variant

#### Mapped genes with CDS features

Chromosome	Han1 total (bp)	Non-HG00621 sequence Ra (inserted from CHM13) (bp)	tio of source sequences (HG00621:CHM13)	
1	249,525,787	119,184	0.9995	
2	242,739,747	2,482,037	0.9898	
3	200,211,729	377,991	0.9981	
4	192,045,028	518,393	0.9973	
5	181,667,637	494,129	0.9973	
6	170,861,069	314,798	0.9982	
7	160,865,769	107,243	0.9993	
8	145,880,131	791,768	0.9946	
9	148,018,047	35,504,706	0.7601	
10	135,316,043	585,347	0.9957	
11	135,129,219	874,841	0.9935	
12	134,132,185	102,971	0.9992	
13	111,903,191	10,782,722	0.9036	
14	101,435,482	5,090,291	0.9498	
15	101,210,777	12,429,469	0.8772	
16	95,412,483	13,280,238	0.8608	
17	83,450,189	1,080,955	0.9870	
18	78,996,361	210,798	0.9973	
19	61,978,944	1,089,081	0.9824	
20	65,189,243	963,587	0.9852	
21	45,827,290	5,613,897	0.8775	
22	50,610,422	5,397,082	0.8934	
Х	154,227,164	7,056,525	0.9542	
Y	53,057,190	14,607,629	0.7247	
Μ	16,571	0	1.0000	
Total	3,099,707,698	119,875,682	0.9614	






Conserved genes (82%)		Altered genes (18%)						
Identical	Synonymous	Nonsynonymous	Start lost	Stop gained	Truncated	Frameshift	In-frame insertion	In-frame deletion
14130	2442	3200	21	31	1+39 (5'+3')	143	110	92

• LiftoffTools: <sup>(2)</sup> Synteny

#### • LiftoffTools: <sup>(3)</sup> Clusters



#### **Homozygous Altered Genes**

Gene name	RefSeq protein length –	Protein length		Protein length ratio (vs RefSeq)		Amino acid identity score (vs RefSeq)	
	protein length	Han1	<b>T2T-CHM13</b>	Han1	<b>T2T-CHM13</b>	Han1	T2T-CHM13
MUC19	6,985	1,241	5,332	0.178	0.763	0.278	0.943
AQP12A	295	163	295	0.553	1.000	0.551	0.986
RETNLB	111	111	14	1.000	0.126	1.00	0.295
TCP11X1	407	407	201	1.000	0.494	0.997	0.417

# **Comparing Han1 with Published Assemblies**

Genome	Ethnicity	Contig N50 (Mb)	Number of contigs	Number of gaps	Assembly size (Gb)
Han1 <sup>1</sup>	Southern Han Chinese	148.02	25	0	3.10
HG00621 (hifiasm) <sup>2</sup>	Southern Han Chinese	95.77	182	157	3.11
T2T-CHM13v2.0 <sup>3</sup>	Northern European	150.62	25	0	3.12
HJ-H1 <sup>4</sup>	Northern Han Chinese	28.15	1,330	427	3.07
HJ-H2 <sup>4</sup>	Northern Han Chinese	25.90	896	390	2.91
NH1 <sup>4</sup>	Northern Han Chinese	3.60	11,019	8,484	2.89
HX1 <sup>4</sup>	Southern Han Chinese	8.33	5,843	4,025	2.93
YH2.0 <sup>5</sup>	Southern Han Chinese	0.02	361,157	235,514	2.91
TJ1.p0 <sup>6</sup>	Tujia	13.67	1430	907	2.87
TJ1.p1 <sup>6</sup>	Tujia	13.70	1426	873	2.87
ZF1 <sup>7</sup>	Tibetan	23.62	1384	1360	2.85
GRCh38.p14 <sup>8</sup>	Mixed	57.88	994	804	3.10

DEFB126	111	111	82	1.000	0.739	0.936	0.576
TPSB2	275	275	166	1.000	0.604	0.927	0.584
PBOV1	135	136	135	1.010	1.000	1.000	0.875
GOLGA6L10	536	550	522	1.026	0.974	0.945	1.000
KLHDC7B	1,235	1,211	1,215	0.981	0.984	0.978	0.981
NBPF19	3,843	3,283	3,772	0.854	0.982	0.927	0.961
RP1L1	2,400	2,417	2,464	1.007	1.027	0.989	0.970
TMEM82	343	343	344	1.000	1.000	1.000	0.994
KIR2DL3	341	341	341	1.00	1.000	1.000	0.956

# Conclusions

• First gapless, fully annotated Southern Han Chinese Genome

• First gene content comparison between two individuals

Acknowledgements: supported by NIH under grants R01-HG006677 and R35-GM130151 and by NSF under grants IOS-1744309 and DBI-1759518.