

Urological Associates Medication Instructions

For a period of 5 7 10 days before the procedure, you must **stop** the use of aspirin or medications containing aspirin. Studies indicate and our personal experience shows that taking aspirin increases bleeding during and after procedures or surgeries because it is such a powerful anticoagulant.

Below is a list of **most** of the medications that contain aspirin that are available with and without a prescription.

PLEASE CHECK WITH YOUR DOCTOR IF YOU ARE TAKING COUMADIN, PLAVIX OR WARFARIN.

Advil	Extra Strength Bufferin
Alka-Seltzer Effervescent Pain Reliever and Antacid	Florinaal
Alka-Seltzer Plus Cold Medicine	Fish Oil
Aleve	4 Way Cold Tablets
Anacin Analgesic (capsules & tablets)	Goody's Headache Powders
Anacin maximum Strength Analgesic (capsules and tablets)	Halfprin
Arthritis Strength Bufferin	Ibuprofen
Arthrotec	Lortab ASA
Ascriptin	Motrin
Ascriptin A/D	Micrainin
Aspergum	Midol
Aspirin suppositories	Momentum
Bayer Aspirin	Norwich Aspirin
Bayer Children's Chewable Aspirin	Nuprin
Bayer children's Cold Tablets	Orudis
Bayer time Release Aspirin	P-A-C Analgesic
Bufferin	Panagesic
Congespirin	Pepto Bismol
Cope	Percodan
Coricidin "D" Decongestant tablets	Plavix
Coricidin Demilet tablets	Quiet World Analgesic Sleep-aid
Coricidin Medilets tablets	Rixiprin
Coricidin tablets	St Joseph's Aspirin for kids
Coumadin	St Joseph's cold tablets for kids
Darvon Compound-65	Sine-Off Sinus Medicine
Dristan Decongestant/Antihistamine/Analgesic Tablets	Sloprin
Easprin	Triminacin tablets
Ecotrin	Vanquish
Empirin	Vio-Med tablets
Entab-650 tablets	Vitamin E
Excedrin	Voltaren
	Zorprin