

Low-Purine Diet

Purines (PYUR-eenz) are found in many foods, especially organ meats, anchovies, mackerel, and sardines. Purines make up 15 percent of the uric acid found in the body.

Too much uric acid can lead to problems, including kidney stones and gout. For this reason, a low-purine diet is recommended for people who have kidney stones, gout, and sometimes for people who have had an organ transplant.

The diet is often used with medication to lower uric acid levels. Some people can lower their uric acid levels through diet alone.

Not everyone needs to follow a rigid diet to treat gout, but avoiding foods that are high in purines may help. Check with your doctor or dietitian to see if you should follow this diet.

Important Points to Keep in Mind

- Avoid high-purine foods. See attached lists.
- Avoid or limit alcohol. Alcohol increases purine production, leading to higher uric acid levels in your blood and urine.
- Limit meat to 3 ounces per meal.
- Limit high-fat foods such as salad dressings, ice cream, fried foods, gravies, and dressings. Fat holds onto uric acid in your kidneys.
- Eat enough carbohydrates. They help your body get rid of extra uric acid.
- If you are overweight, lose weight gradually. Rapid weight loss can increase uric acid levels.
- Drink 8 to 12 cups of fluid every day to help reduce kidney stone formation.
- Don't take baker's or brewer's yeast as a supplement.

Low-purine Diet

Breads, Cereals, Rice, and Pasta	Foods to Choose	Foods to Avoid
<ul style="list-style-type: none"> • 6-11 servings each day • Serving size= 1 slice bread, 1 cup ready-to-eat cereal, 1/2 cup cooked cereal, rice or pasta 	<ul style="list-style-type: none"> • All enriched breads, cereals, rice, noodles, pasta, and potatoes • Limit to 2 servings per week: whole-grain breads and cereals, wheat germ, bran and oatmeal 	<ul style="list-style-type: none"> • Limit high-fat breads like pancakes, French toast, biscuits, muffins, and French fries

Fruit	Foods to Choose	Foods to Avoid
<ul style="list-style-type: none"> • 2-4 servings each day • Serving size= 1 medium-size piece of fresh fruit, 1/2 cup canned fruit, 3/4 cup fruit juice 	<ul style="list-style-type: none"> • All fruit and juices 	<ul style="list-style-type: none"> • Limit avocados (high in fat)

Milk and Dairy Products	Foods to Choose	Foods to Avoid
<ul style="list-style-type: none"> • 2 servings each day • Serving size= 1 cup milk or yogurt 	<ul style="list-style-type: none"> • Skim or low-fat milk • Low-fat yogurt 	<ul style="list-style-type: none"> • Whole milk, cream, and sour cream

Vegetables	Foods to Choose	Foods to Avoid
<ul style="list-style-type: none"> • 3 servings each day • Serving size= 1 cup raw, 1/2 cup cooked or chopped 	<ul style="list-style-type: none"> • All • Limit to 2 servings per week: mushrooms, dried peas and beans, spinach, asparagus, cauliflower 	<ul style="list-style-type: none"> • Limit high-fat cooking, including au gratin, fried foods, and cream sauces

Meats, Poultry, Fish, Dried Beans, Peas, Eggs, and Cheese	Foods To Choose	Foods to Avoid
<ul style="list-style-type: none"> • Total of 6 ounces daily • Serving size= 2-3 ounces cooked (count 1 egg, 1/2 cup cooked beans, 2 tablespoons peanut butter, or 1 ounce of cheese as 1 ounce of meat). 	<ul style="list-style-type: none"> • Beef, lamb, veal, pork, poultry, fish, eggs, peanut butter, nuts, and low-fat cheese • Limit to 2 servings per week: dried peas and beans 	<ul style="list-style-type: none"> • Sweetbreads, anchovies, sardines, liver, kidneys, brains, meat extracts, herring, mackerel, scallops, gravies, goose, heart, mincemeat, and mussels

Fats, Condiments, and Drinks	Foods To Choose	Foods to Avoid
	<ul style="list-style-type: none"> • Salt, herbs, spices, and condiments • Carbonated drinks, coffee, cocoa, and tea • Soups made with low-fat milk and vegetable-based broth • Limit to 3 teaspoons daily: butter, margarine, oils and mayonnaise 	<ul style="list-style-type: none"> • Meat gravies, baker's and brewer's yeast, alcohol, and meat stock-based soups (such as bouillon, broth, and consommé)

Snacks, Sweets, and Desserts	Food to Choose	Foods to Avoid
	<ul style="list-style-type: none"> • Gelatin, ice milk, vanilla wafers, angel food cake • Low-fat frozen yogurt 	<ul style="list-style-type: none"> • High-fat desserts such as ice cream, cookies, cakes, pies, doughnuts, and chocolate • Mincemeat pie

Sample Menu

--	--

Breakfast	<ul style="list-style-type: none"> • 1/2 cup orange juice • 1/2 cup farina • Berries • White toast with jelly and 1 teaspoon margarine • 1 cup 2 percent milk • Coffee
Lunch	<ul style="list-style-type: none"> • Hamburger (3 ounces) on bun • Baked potato with 1 teaspoon margarine • 1 cup fruit salad • 2 percent milk
Dinner	<ul style="list-style-type: none"> • Tossed salad with fat-free salad dressing • 1/2 cup chicken breast (3 ounces) with no skin • 1/2 cup rice • 1/2 cup broccoli • Dinner roll with 1 teaspoon margarine • 1/2 cup sherbet • Iced tea

For help in finding a doctor or health service that suits your needs, call the UPMC Referral Service at 412-647-UPMC (8762) or 1-800-533-UPMC (8762). Select option 1.

UPMC is an equal opportunity employer. UPMC policy prohibits discrimination or harassment on the basis of race, color, religion, ancestry, national origin, age, sex, genetics, sexual orientation, gender identity, marital status, familial status, disability, veteran status, or any other legally protected group status. Further, UPMC will continue to support and promote equal employment opportunity, human dignity, and racial, ethnic, and cultural diversity. This policy applies to admissions, employment, and access to and treatment in UPMC programs and activities. This commitment is made by UPMC in accordance with federal, state, and/or local laws and regulations.

Medical information made available on UPMC.com is not intended to be used as a substitute for professional medical advice, diagnosis, or treatment. You should not rely entirely on this information for your health care needs. Ask your own doctor or health care provider any specific medical questions that you have. Further, UPMC.com is not a tool to be used in the case of an emergency. If an emergency arises, you should seek appropriate emergency medical services.

For UPMC Mercy Patients: As a Catholic hospital, UPMC Mercy abides by the Ethical and Religious Directives for Catholic Health Care Services, as determined by the United States Conference of Catholic Bishops. As such, UPMC Mercy neither endorses nor provides medical practices and/or procedures that contradict the moral teachings of the Roman Catholic Church.

© UPMC 2017
Pittsburgh, PA, USA www.upmc.com