Cosentyx (Secukinumab) Infusion/Injection Order

INFUSACARE
www.infusacare.com

Patient Name: DOB: □Male ☐Female

Diagnosis (please provide ICD10 code):

□ Other:

□ NKDA □ Allergies:

□ New Therapy Order □ Continuation of Therapy Date of last dose:

Ordering Provider Name:

Provider NPI: Phone: Fax:

Practice Address: City: State: Zip Code:

Pre-Medication

□ Acetaminophen _____mg by □ PO
□ Cetirizine _____mg by □ PO
□ Diphenhydramine _____mg by □ PO/ □ IVP
□ Solu-Medrol _____mg by □ PO/ □ IVP
□ Solu-Cortef _____mg by □ PO/ □ IVP
□ Other:

Required Documentation:

□ Patient Demographics
□ Patient Insurance (med and pharm card copies, front and back)
□ Progress Notes/Labs supporting diagnosis
□ Quantiferon lab results
□ Other:

Cosentyx may be a weight based drug: Patient Height (cm): Patient Weight (kg):

Cosentyx Medication Order:

Plaque Psoriasis

- Subcutaneous Dosage for Adults
  - □ Induction: 300mg on Weeks 0, 1, 2, 3, 4, and every 4 weeks thereafter
  - □ Maintenance Only: 300mg every 4 weeks
- Subcutaneous Dosage for Pediatric Patients 6 Years and Older
  - Dosage (Please Choose One)
    - □ Patients less than 50kg at the time of dosing, 75mg
    - □ Patients more than 50kg at the time of dosing, 150mg
  - Frequency (Please Choose One)
    - □ Induction: Weeks 0, 1, 2, 3, 4, and every 4 weeks thereafter
    - □ Maintenance Only: Every 4 weeks

Psoriatic Arthritis

- Subcutaneous Dosage for Adults
  - □ Induction: 150mg on Weeks 0, 1, 2, 3, 4, and every 4 weeks thereafter
  - □ Maintenance Only: 150mg every 4 weeks
  - □ Maintenance Only: 300mg every 4 weeks
- Intravenous Dosage for Adults (Do NOT exceed 300mg)
  - □ Induction: 6mg/kg on Initially and 1.75mg/kg every 4 weeks thereafter
  - □ Maintenance Only: 1.75mg/kg every 4 weeks
- Subcutaneous Dosage for Pediatric Patients 2 Years and Older
  - Dosage (Please Choose One)
    - □ Patients between 15kg and 50kg at the time of dosing, 75mg
    - □ Patients more than 50kg at the time of dosing, 150mg
  - Frequency (Please Choose One)
    - □ Induction: Weeks 0, 1, 2, 3, 4, and every 4 weeks thereafter
    - □ Maintenance Only: Every 4 weeks

Ankylosing Spondylitis

- Subcutaneous Dosage for Adults
  - □ Induction: 150mg on Weeks 0, 1, 2, 3, 4, and every 4 weeks thereafter
  - □ Maintenance Only: 150mg every 4 weeks
  - □ Maintenance Only: 300mg every 4 weeks
- Intravenous Dosage for Adults (Do NOT exceed 300mg)
  - □ Induction: 6mg/kg on Initially and 1.75mg/kg every 4 weeks thereafter
  - □ Maintenance Only: 1.75mg/kg every 4 weeks

Non-Radiographic Axial Spondylarthritis

- Subcutaneous Dosage for Adults
  - □ Induction: 150mg on Weeks 0, 1, 2, 3, 4, and every 4 weeks thereafter
  - □ Maintenance Only: 150mg every 4 weeks
- Intravenous Dosage for Adults (Do NOT exceed 300mg)
  - □ Induction: 6mg/kg on Initially and 1.75mg/kg every 4 weeks thereafter
  - □ Maintenance Only: 1.75mg/kg every 4 weeks

Enthesitis-Related Arthritis

- Subcutaneous Dosage for Patients 4 Years and Older
  - Dosage (Please Choose One)
    - □ Patients between 15kg and 50kg at the time of dosing, 75mg
    - □ Patients more than 50kg at the time of dosing, 150mg
  - Frequency (Please Choose One)
    - □ Induction: Weeks 0, 1, 2, 3, 4, and every 4 weeks thereafter
    - □ Maintenance Only: Every 4 weeks

Hidradenitis Suppurativa

- Subcutaneous Dosage for Adults
  - □ Induction: 300mg on Weeks 0, 1, 2, 3, 4, and every 4 weeks thereafter
  - □ Maintenance Only: 300mg every 4 weeks

Additional Instructions/Notes:

Patient IV Access

□ PIV
□ Use patient's SL/DL PICC, flush per protocol
□ Access/deaccess patient's Port, flush per protocol
Imaging results of proper placement required for PICC/Port

OrderingProvider:

Signature: _____ Date: _____
Provider: _____ Phone: _____ Fax: _____
Best Contact Person in Office: _____ Phone: _____

Locations:

□ *Skokie,IL*: 4711 Golf Rd, Ste 900. P: 847-324-6800. F: 224-251-7141
□ *Libertyville, IL*: 1900 Hollister Dr, Ste 210. P: 847-324-6800. F: 224-251-7141
□ *Orland Park, IL*: 16612 107th Ct, Ste B. P: 847-324-6800. F: 224-251-7141
□ *Houston, TX*: 8303 SW Freeway, Ste 111. P: 346-738-9600. F: 346-613-8400