PACROMA Clinical Picture

Case study contributed by:

Dr. Ravi Saini

MD Skin STD Aids and Sexual Disorder Consultant Member of European Academy of Dermatology and Venereology

Vitiligo on the side of the neck

Case history

- The 11-year-old boy has a hyperpigmented linear patch on the left side of his neck that has been present for more than a year.
- He has not experienced itching or any other associated symptoms.

Physical examination

- There is a linear hyperpigmented patch extending from the left ear down to the clavicle, measuring 10mm x 4.5mm.
- The left ear lobule is also affected.

Past and family medical history

- The patient's maternal uncle has vitiligo on the dorsal side of both hands and feet.
- The patient has no other relevant medical history and is fully vaccinated.

Laboratory analysis

 All blood tests, including the typhoid profile were within normal limits.



Issue 4

Diagnosis

Vitiligo


Recommended treatment


Topical Pacroma for 7 months

Follow-up

- The patient showed 50% improvement in 3 months and achieved comparable remission in 7 months
- No side effects were reported, and the treatment was welltolerated by the child

50% improvement in 3 months and comparable remission in 7 months





After Treatment

Discussion

Pimecrolimus can be a preferred treatment choice, especially for children^{1,2}

As effective as clobetasol propionate to restore skin color in lesions of vitiligo in children

Onset of regimentation within the first to third months

Low systemic immunosup-pressive potential (lacks epidermal atrophy and telangiectasia)

Good tolerability profile

References: 1. Sendur N et al. Journal of Dermatological Treatment. 2006; 17: 338-342 2. Sharquie KE et al. Journal of Cosmetics, Dermatological Sciences and Applications. 2015; 5:107-115

In Vitiligo & Mild to Moderate AD

PACROMA 10g XL

Pimecrolimus 1% w/w Cream

PURE PROVEN & SAFE





ajanta pharma limited

Ajanta House, Charkop, Kandivii West, Mumbai 400 067, India