



Holo-K™

Ultra-micronized nutrient

This product is intended for use according to an approved organic system plan. All materials comply with the USDA National Organic Program (NOP) rules and other international bio-organic regulations for certified organic crop production. Verify with your certifier before using.

0-0-12

Guaranteed analysis:

Soluble Potash (K₂O) 12.0%
Sulfur (S) 2.0%
2% Combined Sulfur (S)

Derived from: Potassium sulfate

Net Contents:

- | | |
|---|--|
| <input type="checkbox"/> 1 gallon (3.78 L)
10.6 lbs (4.8 kg) | <input type="checkbox"/> 55 gallon (208.2 L)
583 lbs (264.4 kg) |
| <input type="checkbox"/> 5 gallon (18.92 L)
53 lbs (24 kg) | <input type="checkbox"/> 275 gallon (1041 L)
2915 lbs (1322 kg) |
| <input type="checkbox"/> 30 gallon (113.56 L)
318 lbs (144.2 kg) | |

Density: 10.6 lbs. per gallon at 68° F

F2315

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

PURPOSE: Holo-K™ is a source of soluble potash and sulfur.

APPLICATION RATES AND TIMING

Holo-K™ is intended for foliar, soil or fertigation application and should only be used in accordance with the manufacturer's instructions. Consult your crop consultant for specific application rates.

Soil: Apply up to 2 gallons per acre.

Foliar: Apply up to 1 gallon per acre at appropriate plant growth stages.

Fertigation: Apply up to 1 gallon per acre at appropriate plant growth stages.

Handling: Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 35°F. Crystallization or sedimentation may occur below 35°F or above 95°F. Shake or stir before use to dissolve any residue on the bottom of the container. Do not pre-mix or store in diluted form. Always use product within 8 hours of diluting. All foliar sprays perform best when applied early morning or late afternoon.

When tank mixing: A jar test is recommended: Prepare a small amount of the proposed mix and allow it to stand in a glass jar overnight observing for signs of incompatibility. Ideally, a sample of the test mix should then be sprayed on a small area of the target crop and observed for 3 to 4 days for any phytotoxic effects. Agitation is the key to usability of this product. AEA products involve tiny particles which are far too small to clog filters; however, if they are inadequately diluted or agitated before or during application, there is a possibility of problems with spray equipment. Consider using an educator nozzle to agitate prior to use. Compatibility and/or performance cannot be guaranteed when combined with other products.

Disclaimer: AEA products are sold "as is" and all warranties, expressed or implied, are disclaimed. In no event shall AEA or its affiliates be liable for any damages of any kind for the storage or use of this product. The buyer assumes all responsibility for the use of any AEA products.



Advancing Eco Agriculture, LLC
4551 Parks West Road, Middlefield, OH 44062
www.advancingecoag.com • 800-495-6603