



## HumaCarb™

This product is intended for use according to an approved organic system plan. All materials comply with the USDA National Organic Program (NOP) rules for certified organic crop production. Verify with your organic certifier before using.

### Guaranteed Analysis:

#### CONTAINS NONPLANT FOOD INGREDIENTS

Soil Amending Ingredients:

10% Humic acid (derived from humalite)

90% Total Other Ingredients

#### Net Contents:

- |   |  |
|---|--|
| <input type="checkbox"/> 1 gallon (3.78L)<br>9.2 lbs. (4.17 kg)   | <input type="checkbox"/> 55 gallon (208.2L)<br>506 lbs. (229 kg)   |
| <input type="checkbox"/> 5 gallon (18.92L)<br>46 lbs. (20.8 kg)   | <input type="checkbox"/> 275 gallon (1041L)<br>2530 lbs. (1147 kg) |
| <input type="checkbox"/> 30 gallon (113.56L)<br>276 lbs. (125 kg) |  |

**Density:** 9.20 lbs. per gallon at 68°F

F2315

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

**Purpose:** HumaCarb™ is a soil conditioner and may aid in the uptake of micronutrients.

#### APPLICATION RATES AND TIMING

This product is intended for soil applications and should only be used in accordance with the manufacturer's directions. Check with your crop consultant for specific application rates.

**Side dress or fertigation:** Mix with liquid nitrogen or phosphorous fertilizer at 3% of the solution.

**Small Grains:** Apply up to 2 quarts per acre at late tilling or as needed.

**Vegetables and Fruits:** Apply up to 1 quart per acre per week, starting at transplant or greening.

**Handling:** Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 35°F. Crystallization or sedimentation may occur below 35°F or above 95°F. Shake or stir before use to dissolve any residue on the bottom of the container. Do not pre-mix or store in diluted form. Always use product within 8 hours of diluting.

**When tank mixing:** A jar test is recommended: Prepare a small amount of the proposed mix and allow it to stand in a glass jar overnight observing for signs of incompatibility. Ideally, a sample of the test mix should then be sprayed on a small area of the target crop and observed for 3 to 4 days for any phytotoxic effects. Agitation is the key to usability of this product. AEA products involve tiny particles which are far too small to clog filters; however, if they are inadequately diluted or agitated before and during application there is a possibility of problems with spray equipment. Consider using an eductor nozzle to agitate prior to use. Compatibility and/or performance cannot be guaranteed when combined with other products.

*Disclaimer: AEA products are sold "as is" and all warranties, expressed or implied, are disclaimed. In no event shall AEA or its affiliates be liable for any damages of any kind for the storage or use of this product. The buyer assumes all responsibility for the use of any AEA products.*



**Advancing Eco Agriculture, LLC**

4551 Parks West Road, Middlefield, OH 44062  
www.advancingecoag.com • 800-495-6603