



0-0-1.5

**Guaranteed analysis:**

Soluble Potash (K <sub>2</sub> O).....	1.5%
Boron (B).....	0.04%
Cobalt (Co).....	0.003%
Manganese (Mn) .....	0.08%
0.08% Water Soluble Manganese (Mn)	
Zinc (Zn) .....	0.08%
0.08% Water Soluble Zinc (Zn)	

**Derived from:** kelp, disodium octaborate, zinc sulfate, manganese sulfate, cobalt sulfate

**Net Contents:**

- 1 gallon (3.78L)  
9.3 lbs (4.2 kg)

**Density:** 9.3 lbs. per gallon at 68°F

F2315

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

*Disclaimer: AEA products are sold "as is" and all warranties, expressed or implied, are disclaimed. In no event shall AEA or its affiliates be liable for any damages of any kind for the storage or use of this product. The buyer assumes all responsibility for the use of any AEA products.*

# NutraLive Vitality™

All materials comply with the USDA National Organic Program.

## APPLICATION RATE AND TIMING

NutraLive Vitality™ is a blend of potassium and micro-nutrients intended for the home garden. This product is designed for soil or foliar applications and should only be used in accordance with the manufacturer's instructions.

### Home Gardens, Vegetables and Fruits:

Soil Drench: Mix up to 1/2 oz. per gallon of water and apply over 100 sq. ft. every 7-10 days.

Foliar: Mix up to 1/2 oz. per gallon of water and mist plant leaves every 7-10 days. Expected coverage area is 100 sq. ft. of leaf surface.

Irrigation: Apply at a rate of up to 1/2 oz. per gallon of water and apply over 100 sq. ft. every 7-10 days.

**Handling:** Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 35°F. Crystallization or sedimentation may occur below 35°F or above 95°F. Shake or stir before use to dissolve any residue on the bottom of the bottle. Do not pre-mix or store in diluted form. Always use product within 8 hours of diluting.

**When tank mixing:** A jar test is recommended: Prepare a small amount of the proposed mix and allow it to stand in a glass jar over night observing for signs of incompatibility. Ideally, a sample of the test mix should then be sprayed on a small area of the target plants and observed for 3 to 4 days for any phytotoxic effects.

Agitation is the key to usability of this product. AEA products involve tiny particles which are far too small to clog filters; however, if they are inadequately diluted or agitated before and during application there is a possibility of problems with spray equipment. Consider using an eductor nozzle to agitate prior to use. Compatibility and/or performance cannot be guaranteed when combined with other products.