



SeaCrop®

sea water mineral concentrate

Guaranteed analysis:

Magnesium (Mg)3%
Boron (B)0.02%

Derived from: Concentrated sea water

Density: 9.36 lbs. per gal at 68°

Net Contents:

- | | |
|---|--|
| <input type="checkbox"/> 1 gallon (3.78L)
9.36 lbs | <input type="checkbox"/> 55 gallon (208.2L)
514.8 lbs |
| <input type="checkbox"/> 5 gallon (18.92L)
46.8 lbs | <input type="checkbox"/> 275 gallon (1041L)
2574 lbs |
| <input type="checkbox"/> 30 gallon (113.56L)
280.8 lbs | |

F2315

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

This product is intended for use according to an approved organic system plan. Sea-Crop® is an organic input consisting of natural sea minerals from concentrated sea water. Verify with your organic certifier before using.

APPLICATION RATES AND TIMING

This product is recommended for soil and foliar applications and should be used only in accordance with the manufacturer's instructions. Check with your crop consultant for specific application rates.

Vegetables: Apply up to 1 quart per acre per week in irrigation system or as a foliar.

Broad Acre Crops: Apply up to 1 gallon per acre (must be split application), per season.

Soil Drench or Transplant Solution: Use up to 2 gallons per acre in a 2% solution.

Handling: Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 35°F. Crystallization or sedimentation may occur below 35°F or above 95°F. Shake or stir before use, especially when near empty to dissolve the residue on the bottom of the container. Do not pre-mix or store in diluted form. Always use product within 8 hours of diluting. All foliar sprays perform best when applied early morning or late afternoon.

When tank mixing: A jar test is recommended: Prepare a small amount of the proposed mix and allow it to stand in a glass jar over night observing for signs of incompatibility. Ideally, a sample of the test mix should then be sprayed on a small area of the target crop and observed for 3 to 4 days for any phytotoxic effects. Agitation is the key to usability of this product. AEA products involve tiny particles which are far too small to clog filters; however, if they are inadequately diluted or agitated before and during application, there is a possibility of problems with spray equipment. Consider using an eductor nozzle to agitate prior to use. Compatibility and/or performance cannot be guaranteed when combined with other products.

Disclaimer: AEA products are sold "as is" and all warranties, expressed or implied, are disclaimed. In no event shall AEA or its affiliates be liable for any damages of any kind for the storage or use of this product. The buyer assumes all responsibility for the use of any AEA products.



Advancing Eco Agriculture, LLC
4551 Parks West Road, Middlefield, OH 44062
www.advancingecoag.com • 800-495-6603