

---

# Long-Term Variable Milfoil Management Plan



*Chance Pond Brook  
Franklin, New Hampshire*

---

## Contents

<b>Purpose</b> .....	<b>4</b>
<b>Invasive Aquatic Plant Overview</b> .....	<b>4</b>
<b>Variable Milfoil Infestation in Chance Pond Brook</b> .....	<b>5</b>
<b>Milfoil Management Goals and Objectives</b> .....	<b>6</b>
<b>Local Support</b> .....	<b>6</b>
Town or Municipality Support .....	6
Local Associations .....	6
<b>Waterbody Characteristics</b> .....	<b>6</b>
<b>Beneficial (Designated) Uses of Waterbody</b> .....	<b>7</b>
Aquatic Life .....	8
Recreational Uses and Access Points.....	8
<b>Macrophyte Community Evaluation</b> .....	<b>8</b>
<b>Wells and Water Supplies</b> .....	<b>8</b>
<b>Historical Control Activities</b> .....	<b>9</b>
<b>Aquatic Invasive Plant Management Options</b> .....	<b>9</b>
<b>Feasibility Evaluation of Control Options in this Waterbody</b> .....	<b>10</b>
<b>Recommended Actions, Timeframes and Responsible Parties</b> .....	<b>11</b>
<b>Notes</b> .....	<b>13</b>
Target Specificity .....	13
Adaptive Management.....	13
<b>Figure 1: Map of Variable Milfoil Infestation</b> .....	<b>14</b>
<b>Figure 2: Critical Habitats or Conservation Areas</b> .....	<b>16</b>
<b>Figure 3: Wells and Water Supplies</b> .....	<b>17</b>
<b>Figure 4: Control Actions</b> .....	<b>18</b>
2017 Proposed .....	18

<b>Appendix A</b>	<b>Aquatic Plant Control Techniques .....</b>	<b>19</b>
<b>Appendix B</b>	<b>Control Practices Used New Hampshire.....</b>	<b>23</b>
	Restricted Use Areas and Fragment Barrier: .....	23
	Hand-pulling:.....	23
	Diver Assisted Suction Harvesting .....	24
	Mechanical Harvesting .....	24
	Benthic Barriers: .....	24
	Targeted Application of Herbicides: .....	25
	Extended Drawdown .....	26
	Dredging .....	26
	Biological Control.....	26
	<b>References .....</b>	<b>27</b>

## Purpose

The purposes of this exotic aquatic plant management and control plan are:

1. To identify and describe the historic and current exotic aquatic infestation(s) in the waterbody;
2. To identify short-term and long-term exotic aquatic plant control goals;
3. To minimize any adverse effects of exotic aquatic plant management strategies on non-target species;
4. To recommend exotic plant control actions that meet the goals outlined in this plan; and
5. To evaluate control practices used in this waterbody over time to determine if they are meeting the goals outlined in this plan.

This plan also summarizes the current physical, biological, ecological, and chemical components of the subject waterbody as they may relate to both the exotic plant infestation and recommended control actions, and the potential social, recreational and ecological impacts of the exotic plant infestation.

The intent of this plan is to establish an adaptive management strategy for the long-term control of the target species (in this case variable milfoil) in the subject waterbody, using an integrated plant management approach.

Appendix A and Appendix B detail the general best management practices and strategies available for waterbodies with exotic species, and provide more information on each of the activities that are recommended within this plan.

## Invasive Aquatic Plant Overview

Exotic aquatic plants pose a threat to the ecological, aesthetic, recreational, and economic values of lakes and ponds (Luken & Thieret, 1997, Halstead, 2000), primarily by forming dense growths or monocultures in critical areas of waterbodies that are important for aquatic habitat and/or recreational use. Under some circumstances, dense growths and near monotypic stands of invasive aquatic plants can result, having the potential to reduce overall species diversity in both plant and animal species, and can alter water chemistry and aquatic habitat structure that is native to the system.

Since January 1, 1998, the sale, distribution, importation, propagation, transportation, and introduction of key exotic aquatic plants have been prohibited (RSA 487:16-a) in New Hampshire. This law was designed as a tool for lake managers to help prevent the spread of nuisance aquatic plants.

---

New Hampshire lists 28 exotic aquatic plant species as prohibited in the state (per Env-Wq 1303.02) due to their documented and potential threat to surface waters of the state.

According to the federal Section 305(b) and 303(d) Consolidated Assessment and Listing Methodology (CALM), “exotic macrophytes are non-native, fast growing aquatic plants, which can quickly dominate and choke out native aquatic plant growth in the surface water. Such infestations are in violation of New Hampshire regulation Env-Wq 1703.19, which states that surface waters shall support and maintain a balanced, integrated and adaptive community of organisms having a species composition, diversity, and functional organization comparable to that of similar natural habitats of a region” (DES, 2006). In fact, waterbodies that contain even a single exotic aquatic plant do not attain water quality standards and are listed as impaired.

### Variable Milfoil Infestation in Chance Pond Brook

Variable milfoil (*Myriophyllum heterophyllum*) was documented in Chance Pond Brook in Franklin, New Hampshire in the summer of 2017. Based on the report of a concerned local resident, DES performed an initial site inspection on August 16, 2017, and documented extensive growth along the impounded section of the brook downstream of Webster Lake. A full survey was conducted of Webster Lake upstream, and fortunately no milfoil was found. The boat launch to the lake is located right above the outlet stream, so it is feasible that milfoil was introduced into Webster Lake, and then immediately flowed downstream into Chance Pond Brook. Alternatively, Chance Pond Brook is used by local fisherman who cast in from the shore, so contaminated tackle could be the vector. In any event, variable milfoil was likely present in Chance Pond Brook for 2-3 years prior to it being reported, as growth was expansive when the initial survey was performed.

Figure 1 illustrates the distribution of variable milfoil in this waterbody, and the table below provides a summary of growth over time.

Area	Location/Area Description	Year	Description of Growth	Percent Cover
North of Carr Street	Northern reach of Chance Pond Brook between Webster Lake Road and Carr St.	2017	Patchy areas of growth at the northern end, below the bridge, increasing in density towards Carr St. Some flowering stalks observed.	40%

<b>Area</b>	<b>Location/Area Description</b>	<b>Year</b>	<b>Description of Growth</b>	<b>Percent Cover</b>
South of Carr Street	Southern basin of Chance Pond Brook, south Carr St. to the dam.	2017	High density growth throughout most of this basin, with plants topped out late season in five feet of water. Several flowering stalks observed.	85%

In terms of the impacts of the variable milfoil in the system, there are about a dozen houses around the immediate shoreline and short setback distance of Chance Pond Brook. Lake residents have expressed concern about the exotic plant growth, and concerns about continued expansion of the plant into Webster Lake.

### **Milfoil Management Goals and Objectives**

The goal for Chance Pond Brook is the reduction of overall biomass and distribution of variable milfoil in the system, with the eventual eradication (if feasible) using an Integrated Pest Management Approach.

### **Local Support**

#### **Town or Municipality Support**

The town of Franklin appreciates the importance of keeping the Chance Pond Brook system usable and controlling the variable milfoil. This is the only infested waterbody in the town at this point, and local officials recognize the need to protect other nearby waterbodies.

#### **Local Associations**

There is no local lake association on Chance Pond Brook, though the lake association members from Webster Lake upstream have expressed an interest in this project, and concerns about upstream migration of the milfoil. They will be stepping up their Weed Watcher efforts to help detect new infestations early.

### **Waterbody Characteristics**

The following table summarizes basic physical and biological characteristics of Chance Pond Brook, including the variable milfoil infestation. Note that a current review of the Natural Heritage Bureau (NHB) database was requested and the results from that search are included in the table below, as well as in other key sections of this report as they may pertain to the type of species (fish, wildlife, habitat, or macrophyte).

Parameter/Measure	Value/Description
Waterbody (acres)	Approximately 17
Shoreline Uses	Residential, forested, transportation
Max Depth (ft)	Estimated at 8.0
Mean Depth (ft)	Estimated at 3.0
Trophic Status	Unknown
Color (CPU) in Epilimnion	No data
Clarity (ft)	No data
Flushing Rate (yr-1)	No data
Natural waterbody/Raised by Damming/Other	Impoundment
Invasive Plants	Variable milfoil ( <i>Myriophyllum heterophyllum</i> )
Infested Area (acres)	See Figures
Distribution	See Figures
Sediment type	
Rare, Threatened, or Endangered Species	<u>2018 NHB Review:</u> American Eel ( <i>Anguilla rostrata</i> )

### Beneficial (Designated) Uses of Waterbody

In New Hampshire, beneficial (designated) uses of our waterbodies are categorized into five general categories: Aquatic Life, Fish Consumption, Recreation, Drinking Water Supply, and Wildlife (CALM).

Of these, Aquatic Life, Wildlife and Recreation are the ones most often affected by the presence of invasive plants, though drinking water supplies can also be affected as well in a number of ways.

Following is a general discussion of the most potentially impacted designated uses, including water supplies and near shore wells, as they relate to this system and the actions proposed in this long-term plan.

The goal for aquatic life support is to provide suitable chemical and physical conditions for supporting a balanced, integrated and adaptive community of aquatic organisms having a species composition, diversity, and functional organization comparable to that of similar natural habitats of the region.

## Aquatic Life

### Fisheries Information

The New Hampshire Natural Heritage Bureau (NHB) lists one species of concern from their survey relative to this treatment program: American Eel (*Anguilla rostrata*).

The American eel was documented in Chance Pond Brook in 2009. There are not expected to be any direct impacts to this species as a result of proposed control measures in Chance Pond Brook. Figure 2 shows a map of the historic range of this eel.

Other fish species reported by the Fish and Game Department include margined madtom, common shiner, golden shiner, and pumpkinseed (information from a 6/11/09 survey).

### Recreational Uses and Access Points

Chance Pond Brook is an impounded stream system, that serves as the outflow of Webster Lake in Franklin. It flows through primarily forested areas, with a few private residences along the shoreline. There are no beaches, public access points, or other recreational facilities on the waterbody. Fishermen have been observed along the shoreline, particularly along Carr St., and along the southwestern shoreline of the southern basin on the pond, on a rock outcrop on shore. Due to the shallow depth of the upper basin, swimming is not feasible. The southern basin is deeper, and could be used for undesignated swimming.

## Macrophyte Community Evaluation

The littoral zone is defined as the nearshore areas of a waterbody where sunlight penetrates to the bottom sediments. The littoral zone is typically the zone of rooted macrophyte growth in a waterbody. Due to the shallow nature of this system, the entire waterbody would fall within the littoral zone.

The macrophyte community of Chance Pond Brook is characterized by a mix of native and non-native (variable milfoil) plant growth. Native species include a mix of floating plants (white and yellow water lilies, water shield), emergent plants (bur-reed, pickerelweed), and submergent plants (various pondweeds, tape-grass, spike-rush, water naiad, whorled bladderwort, small bladderwort, filamentous green algae). Native plant communities are mixed, and can be characterized as common.

## Wells and Water Supplies

Figure 3 shows the location of wells, water supplies, well-head protection areas, and drinking water protection areas around the subject waterbody,

---

based on information in the DES geographic information system records. Note that it is likely that Figure 3 does not show the location of all private wells.

Note that the map in Figure 3 cannot be provided on a finer scale than 1:48,000. Due to public water system security concerns, a large-scale map may be made available upon agreement with DES' data security policy. Visit DES' OneStop Web GIS, <http://www2.des.state.nh.us/gis/onestop/> and register to Access Public Water Supply Data Layers. Registration includes agreement with general security provisions associated with public water supply data. Paper maps that include public water supply data may be provided at a larger-scale by DES' Exotic Species Program after completing the registration process.

In the event that an herbicide treatment is needed for this waterbody, the applicator/contractor will provide more detailed information on the wells and water supplies within proximity to the treatment areas as required in the permit application process with the Division of Pesticide Control at the Department of Agriculture. It is beyond the scope of this plan to maintain updated well and water supply information other than that provided in Figure 3.

### **Historical Control Activities**

This is a new infestation, and no historic control actions have been performed.

### **Aquatic Invasive Plant Management Options**

The control practices used should be as specific to the target species as feasible. No control of native aquatic plants is intended.

Exotic aquatic plant management relies on a combination of proven methods that control exotic plant infestations, including physical control, chemical control, biological controls (where they exist), and habitat manipulation. Integrated Pest Management Strategies (IPM) are typically implemented using Best Management Practices (BMPs) based on site-specific conditions so as to maximize the long-term effectiveness of control strategies. Descriptions for the control activities are closely modeled after those prescribed by the Aquatic Ecosystem Restoration Foundation (AERF) (2004). This publication can be found online at <http://www.aquatics.org/bmp.htm>. Additional information can be obtained from a document prepared for the State of Massachusetts called the Generic Environmental Impact Report for Lakes and Ponds, available at <http://www.mass.gov/dcr/watersupply/lakepond/geir.htm>.

---

Criteria for the selection of control techniques are presented in Appendix A. Appendix B includes a summary of the exotic aquatic plant control practices currently used by the State of New Hampshire.

### Feasibility Evaluation of Control Options in this Waterbody

DES has evaluated the feasibility of potential control practices on the subject waterbody. The following table summarizes DES' control strategy recommendations for the subject waterbody:

<b>Control Method</b>	<b>Use on Chance Pond Brook</b>
Restricted Use Areas (RUAs) and/or Fragment Barriers	The purpose of RUAs and fragment barriers is to contain small areas of exotic aquatic plant growth to prevent them from spreading further in a system.  If variable milfoil is reduced by other integrated approaches outlined in this plan, then RUAs and fragment barriers may be a future consideration based on the size, configuration and location of remaining areas of growth.
Hand harvesting/Diver-Assisted Suction Harvesting (DASH)	DES recommends that hand harvesting and/or DASH be employed as appropriate based on milfoil density.
Mechanical Harvesting/Removal	Not recommended due to the risk of fragmentation and drift, and subsequent further spread of the invasive plant.
Benthic Barriers	Recommended for small patches that are 20' x 20' in size or less, and where practical.
Herbicides	Herbicide treatment is recommended as a primary means of control only where infestations of the exotic plant are too widespread and/or dense for non-chemical means of control to be effective, or where water depth or other factor limits the effectiveness of divers.
Extended Drawdown	Not feasible or practical for this waterbody due to lack of an impoundment structure.
Dredge	Cost prohibitive and not often effective for controlling invasive aquatic plants.
Biological Control	No biological controls are yet approved for use on variable milfoil.
No Control	Milfoil is widespread in this basin, and as such, it poses a spread risk to downstream areas such as the Pemigewasset River. Human vectors

Control Method	Use on Chance Pond Brook
	could also spread this plant upstream into Webster Lake through contaminated tackle or other recreational gear (canoes/kayaks/etc).

### Recommended Actions, Timeframes and Responsible Parties

An evaluation of the size, location, and type of variable milfoil infestation, as well as the waterbody uses was conducted at the end of the last growing season (see attached figures for findings). Based on this survey the following recommendations are made for variable milfoil control in the system. Figure 4 includes maps showing control actions.

Year	Action	Responsible Party	Schedule
2018	Perform a spring Weed Watcher training for interested local residents and encourage them to Weed Watch monthly during the growing season	DES/Local Weed Watchers	Spring training, monthly surveys in growing season
	Herbicide treatment	SOLitude Lake Management, LLC	Early June or early September
	Diver work after treatment as needed	DES	As needed in growing season
	Survey waterbody and planning for next season's control actions	DES	September
2019	Weed Watching and marking/reporting of milfoil growth	Local Weed Watchers	Once a month from May through September
	Survey and planning for summer/fall milfoil control actions	DES	May/June
	Diver/DASH work as needed and recommended (areas to be determined based on need)	Contract Diver	May through September as needed

<b>Year</b>	<b>Action</b>	<b>Responsible Party</b>	<b>Schedule</b>
	Herbicide treatment, if needed	SOLitude Lake Management, LLC	Early June or early September
	Survey and planning for next season's control actions	DES	September
2020	Weed Watching and marking/reporting of milfoil growth	Local Weed Watchers	Once a month from May through September
	Survey and planning for summer/fall milfoil control actions	DES	May/June
	Diver/DASH work as needed and recommended (areas to be determined based on need)	Contract Diver	May through September as needed
	Herbicide treatment, if needed	SOLitude Lake Management, LLC	Early June or early September
	Survey and planning for next season's control actions	DES	September
2021	Weed Watching and marking/reporting of milfoil growth	Local Weed Watchers	Once a month from May through September
	Survey and planning for summer/fall milfoil control actions	DES	May/June
	Diver/DASH work as needed and recommended (areas to be determined based on need)	Contract Diver	May through September as needed
	Herbicide treatment, if needed	SOLitude Lake Management, LLC	Early June or early September
	Survey and planning for next season's control actions	DES	September

<b>Year</b>	<b>Action</b>	<b>Responsible Party</b>	<b>Schedule</b>
2022	Weed Watching and marking/reporting of milfoil growth	Local Weed Watchers	Once a month from May through September
	Survey and planning for summer/fall milfoil control actions	DES	May/June
	Herbicide treatment, if needed	SOLitude Lake Management, LLC	Early June or early September
	Survey and planning for next season's control actions	DES	September
2023	Update and revise Long-Term Variable Milfoil Control Plan	DES and Interested Parties	Fall/Winter

## Notes

### Target Specificity

It is important to realize that aquatic herbicide applications are conducted in a specific and scientific manner. To the extent feasible, the permitting authority favors the use of selective herbicides that, where used appropriately, will control the target plant with little or no impact to non-target species, such that the ecological functions of native plants for habitat, lake ecology, and chemistry/biology will be maintained. *Not all aquatic plants will be impacted as a result of an herbicide treatment.*

### Adaptive Management

Because this is a natural system that is being evaluated for management, it is impossible to accurately predict a management course over five years that could be heavily dependent on uncontrolled natural circumstances (weather patterns, temperature, adaptability of invasive species, etc).

This long-term plan is therefore based on the concept of adaptive management, where current field data (from field survey work using DES established field survey standard operating procedures) drive decision making, which may result in modifications to the recommended control actions and timeframes for control. As such, this management plan should be considered a dynamic document that is geared to the actual field conditions that present themselves in this waterbody.

If circumstances arise that require the modification of part or all of the recommendations herein, interested parties will be consulted for their input on revisions that may be needed to further the goal of variable milfoil in the subject waterbody.

**Figure 1: Map of Variable Milfoil Infestation**






Figure 2: Critical Habitats or Conservation Areas

NHB18-0135



Figure 3: Wells and Water Supplies





Figure 4: Control Actions

2017 Proposed



## **Appendix A Aquatic Plant Control Techniques**

### Preliminary Investigations

#### **I. Field Site Inspection**

- Verify genus and species of the plant.
- Determine if the plant is a native or exotic species per RSA 487:16, II.
- Map extent of the exotic aquatic plant infestation (area, water depth, height of the plant, density of the population).
- Document any native plant abundances and community structure around and dispersed within the exotic/nuisance plant population (provide updated native plant map after review of milfoil in the Fall or after treatment)

#### **II. Office/Laboratory Research of Waterbody Characteristics**

- Contact the appropriate agencies to determine the presence of rare or endangered species in the waterbody or its prime wetlands.
- Determine the basic relevant limnological characteristics of the waterbody (size, bathymetry, flushing rate, nutrient levels, trophic status, and type and extent of adjacent wetlands).
- Determine the potential threat to downstream waterbodies from the exotic aquatic plant based on limnological characteristics (water chemistry, quantity, quality as they relate to movement or support of exotic plant growth).

### Overall Control Options

For any given waterbody that has an infestation of exotic plants, one of four options will be selected, based on the status of the infestation, the available management options, and the technical knowledge of the DES Limnologists and other key resource managers who have conducted the field work and who are preparing or contributing to this plan. The options are as follows:

- 1) **Eradication:** The goal is to completely remove the exotic plant infestation over time. In some situations this may be a rapid response that results in an eradication event in a single season (such as for a new infestation), in other situations a longer-term approach may be warranted given the age and distribution of the infestation. Eradication is more feasible in smaller systems without extensive expanded growth (for example, Lake Winnepesaukee is unlikely to achieve eradication of its variable milfoil), or without upstream sources of infestation in other connected systems that continually feed the lake.
  - 2) **Maintenance:** Waterbodies where maintenance is specified as a goal are generally those with expansive infestations, that are larger systems, that have complications of extensive wetland complexes on their periphery, or that have upstream sources of the invasive plant
-

precluding the possibility for eradication. For waterbodies where maintenance is the goal, control activities will be performed on the waterbody to keep an infestation below a desirable threshold. For maintenance projects, thresholds of percent cover or other measurable classification will be indicated, and action will occur when exotic plant growth exceeds the threshold.

- 3) **Containment:** The aim of this approach is to limit the size and extent of the existing infestation within an infested waterbody if it is localized in one portion of that waterbody (such as in a cove or embayment), or if a whole lake is infested action may be taken to prevent the downstream migration of fragments or propagules. This could be achieved through the use of fragment barriers and/or Restricted Use Areas or other such physical means of containment. Other control activities may also be used to reduce the infestation within the containment area.
- 4) **No action.** If the infestation is too large, spreading too quickly, and past management strategies have proven ineffective at controlling the target exotic aquatic plant, DES, in consultation with others, may elect to recommend 'no action' at a particular site. Feasibility of control or control options may be revisited if new information, technologies, etc., develop.

If eradication, maintenance or containment is the recommended option to pursue, the following series of control techniques may be employed. The most appropriate technique(s) based on the determinations of the preliminary investigation will be selected.

Guidelines and requirements of each control practice are suggested and detailed below each alternative, but note that site specific conditions will be factored into the evaluation and recommendation of use on each individual waterbody with an infestation.

#### **A. Hand-Pulling and Diver-Assisted Suction Harvesting**

- Hand-pulling can be used if infestation is in a small localized area (sparsely populated patch of up to 5' X 5', single stems, or dense small patch up to 2' X 2'). For larger areas Diver-Assisted Suction Harvesting (DASH) may be more appropriate.
- Can be used if plant density is low, or if target plant is scattered and not dense.
- Can be used if the plant could effectively be managed or eradicated by hand-pulling or DASH
- Use must be in compliance with the Wetlands Bureau rules.

#### **B. Mechanically Harvest or Hydro-Rake**

- Can not be used on plants which reproduce vegetatively by fragmentation (e.g., milfoil, fanwort, etc.) unless containment can be ensured.
  - Can be used only if the waterbody is accessible to machinery.
-

- Can be used if there is a disposal location available for harvested plant materials.
- Can be used if plant depth is conducive to harvesting capabilities (~ <7 ft. for mower, ~ <12 ft. for hydro-rake).
- If a waterbody is fully infested and no other control options are effective, mechanical harvesting can be used to open navigation channel(s) through dense plant growth.

### **C. Herbicide Treatment**

- Can be used if application of herbicide is conducted in areas where alternative control techniques are not optimum due to depth, current, use, or density and type of plant.
- Can be used for treatment of exotic plants where fragmentation is a high concern.
- Can be used where species specific treatment is necessary due to the need to manage other plants
- Can be used if other methods used as first choices in the past have not been effective.
- A licensed applicator should be contacted to inspect the site and make recommendations about the effectiveness of herbicide treatment as compared with other treatments.

### **D. Restricted Use Areas (per RSA 487:17, II (d))**

- Can be established in an area that effectively restricts use to a small cove, bay, or other such area where navigation, fishing, and other transient activities may cause fragmentation to occur.
- Can not be used when there are several “patches” of an infestation of exotic aquatic plants throughout a waterbody.
- Can be used as a temporary means of control.

### **E. Bottom Barrier**

- Can be used in small areas, preferably less than 10,000 sq. ft.
- Can be used in an area where the current is not likely to cause the displacement of the barrier.
- Can be used early in the season before the plant reaches the surface of the water.
- Can be used in an area to compress plants to allow for clear passage of boat traffic.
- Can be used in an area to compress plants to allow for a clear swimming area.
- Use must be in compliance with the Wetlands Bureau rules.

### **F. Drawdown**

- Can be used if the target plant(s) are susceptible to drawdown control.
-

- Can be used in an area where bathymetry of the waterbody would be conducive to an adequate level of drawdown to control plant growth, but where extensive deep habits exist for the maintenance of aquatic life such as fish and amphibians.
- Can be used where plants are growing exclusively in shallow waters where a drawdown would leave this area “in the dry” for a suitable period of time (over winter months) to control plant growth.
- Can be used in winter months to avoid encroachment of terrestrial plants into the aquatic system.
- Can be used if it will not significantly impact adjacent or downstream wetland habitats.
- Can be used if spring recharge is sufficient to refill the lake in the spring.
- Can be used in an area where shallow wells would not be significantly impacted.
- Reference RSA 211:11 with regards to drawdown statutes.

#### **G. Dredge**

- Can be used in conjunction with a scheduled drawdown.
- Can be used if a drawdown is not scheduled, though a hydraulic pumping dredge should be used.
- Can only be used as a last alternative due to the detrimental impacts to environmental and aesthetic values of the waterbody.

#### **H. Biological Control**

- Grass carp cannot be used as they are illegal in New Hampshire.
  - Exotic controls, such as insects, cannot be introduced to control a nuisance plant unless approved by Department of Agriculture.
  - Research should be conducted on a potential biological control prior to use to determine the extent of target specificity.
-

## Appendix B Control Practices Used New Hampshire

### **Restricted Use Areas and Fragment Barrier:**

Restricted Use Areas (RUAs) are a tool that can be used to quarantine a portion of a waterbody if an infestation of exotic aquatic plants is isolated to a small cove, embayment, or section of a waterbody. RUAs generally consist of a series of buoys and ropes or nets connecting the buoys to establish an enclosure (or exclosure) to protect an infested area from disturbance. RUAs can be used to prevent access to these infested areas while control practices are being done, and provide the benefit of restricting boating, fishing, and other recreational activities within these areas, so as to prevent fragmentation and spread of the plants outside of the RUA.

### **Hand-pulling:**

Hand-pulling exotic aquatic plants is a technique used on both new and existing infestations, as circumstances allow. For this technique divers carefully hand-remove the shoots and roots of plants from infested areas and place the plant material in mesh dive bags for collection and disposal. This technique is suited to small patches or areas of low density exotic plant coverage.

For a new infestation, hand-pulling activities are typically conducted several times during the first season, with follow-up inspections for the next 1-2 years or until no re-growth is observed. For existing infestations, hand-pulling may be done to slow the expansion of plant establishment in a new area or where new stems are removed in a section that may have previously been uninfested. It is often a follow-up technique that is included in most management plans.

In 2007 a new program was created through a cooperative between a volunteer monitor that is a certified dive instructor, and the DES Exotic Species Program. A Weed Control Diver Course (WCD) was developed and approved through the Professional Association of Dive Instructors (PADI) to expand the number of certified divers available to assist with hand-pulling activities. DES has only four certified divers in the Limnology Center to handle problems with aquatic plants, and more help was needed. There is a unique skill involved with hand-removing plants from the lake bottom. If the process is not conducted correctly, fragments could spread to other waterbody locations. For this reason, training and certification are needed to help ensure success. Roughly 100 divers were certified through this program through the 2010 season. DES maintains a list of WCD divers and shares them with waterbody groups and municipalities that seek diver assistance for controlling exotic aquatic plants. Classes are offered two to three times per summer.

---

### **Diver Assisted Suction Harvesting**

Diver Assisted Suction Harvesting (DASH) is an emerging and evolving control technique in New Hampshire. The technique employs divers that perform hand removal actions as described above, however, instead of using a dive bag a mechanical suction device is used to entrain the plants and bring them topside where a tender accumulates and bags the material for disposal. Because of this variation divers are able to work in moderately dense stands of plants that cover more bottom area, with increased efficiency and accuracy.

### **Mechanical Harvesting**

The process of mechanical harvesting is conducted by using machines which cut and collect aquatic plants. These machines can cut the plants up to twelve feet below the water surface. The weeds are cut and then collected by the harvester or other separate conveyer-belt driven device where they are stored in the harvester or barge, and then transferred to an upland site.

The advantages of this type of weed control are that cutting and harvesting immediately opens an area such as boat lanes, and it removes the upper portion of the plants. Due to the size of the equipment, mechanical harvesting is limited to water areas of sufficient size and depth. It is important to remember that mechanical harvesting can leave plant fragments in the water, which if not collected, may spread the plant to new areas. Additionally harvesters may impact fish and insect populations in the area by removing them in harvested material. Cutting plant stems too close to the bottom can result in re-suspension of bottom sediments and nutrients. This management option is only recommended when nearly the entire waterbody is infested, and harvesting is needed to open navigation channels through the infested areas.

### **Benthic Barriers:**

Benthic barriers are fiberglass coated screening material that can be applied directly to the lake bottom to cover and compress aquatic plant growth. Screening is staked or weighted to the bottom to prevent it from becoming buoyant or drifting with current. The barriers also serve to block sunlight and prevent photosynthesis by the plants, thereby killing the plants with time. While a reliable method for small areas of plants (roughly 100 sq. ft. or less), larger areas are not reasonably controlled with this method due to a variety of factors (labor intensive installation, cost, and gas accumulation and bubbling beneath the barrier).

---

### **Targeted Application of Herbicides:**

Application of aquatic herbicides is another tool employed for controlling exotic aquatic plants. Generally, herbicides are used when infestations are too large to be controlled using other alternative non-chemical controls, or if other techniques have been tried and have proven unsuccessful. Each aquatic plant responds differently to different herbicides and concentrations of herbicides, but research performed by the Army Corps of Engineers has isolated target specificity of a variety of aquatic herbicides for different species.

Generally, 2,4-D (Navigate formulation) is the herbicide that is recommended for control of variable milfoil. Based on laboratory data this is the most effective herbicide in selectively controlling variable milfoil in New Hampshire's waterbodies.

A field trial was performed during the 2008 summer using the herbicide Renovate to control variable milfoil. Renovate is a systemic aquatic herbicide that targets both the shoots and the roots of the target plant for complete control. In this application it was dispersed as a granular formulation that sank quickly to the bottom to areas of active uptake of the milfoil plants. A small (<5 acre) area of Captains Pond in Salem was treated with this systemic herbicide. The herbicide was applied in pellet form to the infested area in May 2008, and showed good control by the end of the growing season. Renovate works a little more slowly to control aquatic plants than 2,4-D and it is a little more expensive, but presents DES with another alternative that could be used in future treatments.

During the summer of 2010, DES worked with other researchers to perform field trials of three different formulations of 2,4-D in Lake Winnisquam, to determine which product was most target-specific to the variable milfoil. Navigate formulation was used, as were a 2,4-D amine formulation, and a 2,4-D amine and triclopyr formulation (MaxG). Although the final report has not been completed for this study, preliminary results suggest that all three products worked well, but that Navigate formation may be the most target specific of all three.

Another herbicide, Fluridone, is sometimes also used in New Hampshire, mainly to control growths of fanwort (*Cabomba caroliniana*). Fluridone is a systemic aquatic herbicide that inhibits the formation of carotenoids in plants. Reduced carotenoids pigment ultimately results in the breakdown of chlorophyll and subsequent loss of photosynthetic function of the plants.

Other aquatic herbicides are also used in New Hampshire when appropriate (glyphosate, copper compounds, etc). The product of choice will

---

be recommended based on what the target species is, and other waterbody-specific characteristics that are important to consider when selecting a product.

### **Extended Drawdown**

Extended drawdown serves to expose submersed aquatic plants to dessication and scouring from ice (if in winter), physically breaking down plant tissue. Some species can respond well to drawdown and plant density can be reduced, but for invasive species drawdown tends to yield more disturbance to bottom sediments, something to which exotic plants are most adapted. In waterbodies where drawdown is conducted exotic plants can often outcompete native plants for habitat and come to dominate the system.

Some waterbodies that are heavily infested with exotic plants do conduct drawdowns to reduce some of the invasive aquatic plant density. During this reporting period both Northwood Lake (Northwood) and Jones Pond (New Durham) coordinated deep winter drawdowns to reduce growths of variable milfoil (the drawdown on Northwood Lake is primarily for flood control purposes, but they do see some ancillary benefits from the technique for variable milfoil control).

### **Dredging**

Dredging is a means of physical removal of aquatic plants from the bottom sediments using a floating or land-based dredge. Dredging can create a variety of depth gradients creating multiple plant environments allowing for greater diversity in lakes plant, fish, and wildlife communities. However due to the cost, potential environmental effects, and the problem of sediment disposal, dredging is rarely used for control of aquatic vegetation alone.

Dredging can take place in to fashion, including drawdown followed by mechanical dredging using an excavator, or using a diver-operated suction dredge while the water level remains up.

### **Biological Control**

There are no approved biological controls for submersed exotic aquatic plant at this time in New Hampshire.

---

## References

Department of Environmental Services. 2006: 2006 Section 305(b) and 303(d) Consolidated Assessment and Listing Methodology. November 2005. New Hampshire Department of Environmental Services. NHDES-R-WD-05-29. Available at <http://des.nh.gov/WMB/swqa/calm.html>

Halstead, J.M., J. Michaud, S. Hallas-Burt, and J.P. Gibbs. 2003. "An Hedonic Analysis of Effects of a Nonative Invader (*Myriophyllum heterophyllum*) on New Hampshire (USA) Lakefront Properties." *Environmental Management*. 32 (3): 391 – 398

Luken, J.O. and J.W. Thieret. 1997. *Assessment and Management of Plant Invasions*. Springer-Verlag, New York. 324 pages.

---