



Sopheap Tang, DMD & Associates
General Dentistry For Kids

Post Operative Instructions

Please follow the instruction indicated below:

Your child has had a local anesthetic for the required dental treatment. Children do not always understand the effects of a local anesthetic, and may chew or suck the numb lip, tongue, or cheek. Although this is not a serious thing it is an uncomfortable feeling for your child. Please watch your child closely following the appointment for the duration of the anesthetic, usually one to two hours.

Dentistry has been done in the lower jaw. The tongue, lip, as well as the teeth and surrounding tissue will be numb and asleep for approximately 2 hours.

Dentistry has been done in the upper jaw. The lip, teeth, and surrounding tissue will be numb and asleep for approximately 2 hours. Keep your child on a liquid diet until the anesthetic has worn off.

STAINLESS STEEL CROWNS

GENERAL INFORMATION

Sometimes cavities in children's teeth may become so large that simple fillings cannot restore the tooth. To restore the proper shape to the tooth a stainless steel crown is used. This crown is a pre-made cap, which has been trimmed and fit to your child's tooth.

These crowns cover the entire portion of the tooth at the gum lines. Stainless steel is a long-lasting material in the mouth and it will serve quite well for the life of the baby (deciduous) teeth. This will keep the tooth and root intact until normal tooth loss occurs.

Stainless Steel Crowns are used in Several Circumstances:

1. Broken teeth which cannot be repaired by normal fillings.
2. Badly decayed teeth where little tooth structure remains.
3. To prevent breakage of a tooth, which has had a pulpotomy, (partial removal of the nerve).

We usually can place the stainless steel crown in one appointment. We will shape your child's tooth to the necessary form and then trim a stainless steel crown to fit tightly on the tooth. This crown will then be cemented into place.

INSTRUCTIONS FOR HOME CARE

1. There are no restrictions on your daily activities.
2. Encourage brushing right away. A clean mouth heals faster.
3. The diet should avoid between meal sweets to help prevent cavities.
4. Hard, sticky foods should be avoided.

NOTE:

1. Your child's teeth and gums around the crown may be tender for several days following the appointment.
2. Your child has had a local anesthetic, watch him/her closely for several hours to prevent him/her from chewing his/her cheek, tongue or lip, which are very numb.

NOTIFY US IF ANY OF THE FOLLOWING OCCURS:

1. Gum around the crown becomes very swollen or tender.
2. The tooth, which has the crown, becomes painful or loose.
3. A gum boil (pimple) forming around the crown.

19040 Van Buren Blvd. ste 111, Riverside, CA 92508

Tel: 951-789-0200 Fax: 951-789-0245

31577 Canyon Estates Dr. ste 205, Lake Elsinore CA, 92532

Tel: 951-674-6900 Fax: 951-674-5934