

Geology and Weathering Indices of the Bedrocks and Sediments in Okemesi -Ijero Area, Southwestern Nigeria

Ayodele Olusiji Samuel¹

¹Department of Applied Geology, Federal University of Technology P.M.B. 704, Akure

Abstract

Geological field mapping of the rocks in Okemesi-Ijero area were carried out at a scale of 1:50,000 to determine the underlying lithologic units and petrological characteristics. Also the suitability of the rocks and sediments for geochemical exploration purposes were determined using weathering indices parameters such as chemical index of weathering (CIW), weathering index of Parker (WIP), Plagioclase index of Alteration (PIA) and chemical index of Alteration (CIA). Thirty-five bedrocks and stream sediment samples were collected from various locations within the studied area using grid-controlled sampling techniques. Field studies revealed that the sediments were derived from weathering of the bedrocks such as quartz-biotite-schists, banded gneisses, granite-gneisses, biotite gneisses, calc gneisses, porphyritic granites, charnockites, massive/schistose quartzites and mica schists which are the dominant lithologic units in the studied area. The weathering profiles based on the above parameters showed that sediments from River Elewu and Issa 7c are weakly weathered whereas, sediments from Agbagbara, Ede-1, Ede-2, Efon-Okemesi-2, Esa-Oke-2, Odo-OWa Abidogun, Ofale road, Okemesi-2, 3 and 9 were intensely weathered which makes them unsuitable for geochemical exploration work due to complete weathering of the mineral grains.

Keywords: Okemesi; Ijero; Lithologic units; Sediments; Weathering Indices

1. Introduction

A range of media including stream sediments, soils and waters have been used in many countries and across different scales for geochemical mapping (e.g. Darnley, 1990; BGS, 1990; Riemann and Caritat, 1998; Rice, 1999; Key; De Waele; Linyungu, 2004; Johnson, Breward and Ander, 2005). Stream sediment is a composite of the products of weathering and erosion derived from the catchment basin, and funneled into and along the stream channels. Very fine grained sediments tend to have assemblages rich in zircon which is a result of weathering and sorting from source rock and not of the pressure of zircon-rich source rock. This implies that a regional distribution pattern of heavy minerals is due to various weathering process in source rock and not due to differences in the source.

Geological mapping is the main and one of the basic techniques of gathering geological information on the field where rocks can be observed in-situ. This leads to understanding the geological events of an area or terrain. As disintegration sets in into rocks be it physical or chemical, the aggregates of minerals in the rock are broken down into different fractions which end up in streams and river channels. Stream sediments originate from near the surface of exposed rocks of igneous, sedimentary and volcanic origin. Some of these are easily eroded whereas; others such

as crystalline and metamorphic rocks are affected by stream only when exposed. However, the rocks in the studied area are highly weathered making it difficult to identify minerals and possible geologic structures that could be used to determine the geologic history as well as collection of fresh rock samples. Lots of literatures abound on the various investigations carried out in the study area and related works outside the study area such works include geochemical dispersion of gold in stream sediments in Paleoproterozoic Nyong series, southern Cameroon was also undertaken by Mubfu et al., (2014) in an attempt to explore for gold using stream sediments collected in the Ngo Vayang area of southern Cameroon, the study revealed that the Au-Hf element association from the R-mode factor analysis indicated gold mineralization while U-Th-Pb-W, Nb-Ta-Co-V, Au-Hf-Cu associations reflected lithologic controls. Akintola et al., (2013) carried out the petrography and stream sediment geochemistry of Ede and its environs in order to identify the rock units with their mineralogical appraisal and to determine the concentration and distribution of major and trace elements in the stream sediments with a view to elucidate the mineral potentials of the study area. Emmanuel et al., (2011) carried out geochemical investigation of the southern part of Ilesha with the aim of clarifying the potential source of mineralization in the area. Geologic mapping of

the area revealed that the area is made up of different lithologies such as undifferentiated schists, gneisses and migmatites with pegmatites, schists and epidiorite complex, quartzite and quartz schist. (Olaolun and Oyinloye, 2010; Oyinloye, 1997; Rahaman et al., (1988) and according to Oyinloye and Adebayo (2005), the migmatite-gneiss complex in Ijero area is composed of a mafic portion made up of biotite, hornblende, quartz and opaque minerals, while the felsic portion is quartz-feldsparitic. Other rocks identified by these authors are charnockites, metasediments composed of amphibolite schist, mica-schist and quartzites. Fadipe (1988) also showed that most of the pegmatites in Nigeria especially at Ijero and their environs are associated with cassiterite-tantalite mineralizations which are found associated with migmatite-gneiss and amphibolites. However, this study attempts to unravel the petrological character of the different rock units in the studied area, and carry out geochemical assessment of the bedrocks and sediments for mineral exploration using weathering indices and parameters.

2. Location and Accessibility

The study area lies within latitudes $7^{\circ} 45' N$ and $8^{\circ} 00' N$ and longitudes $4^{\circ} 52' E$ and $5^{\circ} 08' E$. It covers part of the topographic map sheet No. 243 (Ilesha N.E. 1:50,000) and sheet No. 244 (Ado N.W. 1:50,000). The study areas cover parts of Ekiti and Osun, southwestern Nigeria with a total surface area of 821.4 km^2 (Fig.1). Major towns in the area include Okemesi and Ijero Ekiti. Other towns include Epe, Ikoru, Effon, Ipoti, Odo-owa, Ayegunle, while those in Osun State are Oke-ila, Ilupeju, Edemode, Orangun and Oba-sinkin. The areas which fall within Osun, Southwestern Nigeria can be rated moderately motorable due to interconnectivity of roads, while areas within Ekiti can be rated poor because there are only minor roads and footpaths which are not motorable. Localities within Ekiti are mainly small villages with linear settlement along the road, while nucleated settlement predominates in Osun. The study area is generally accessible through network of all seasonal roads and motorable tracks which links it with other part of the country.




Fig.1: Map of Osun and Ekiti, southwestern Nigeria showing the study area (inset: Map of Nigeria showing Osun and Ekiti States).

3. Geology of the study area

The various lithologic descriptions of the different rocks mapped in the study area are discussed below.

Banded gneisses

This lithology covers the southwestern part of the study area around Okokoro, Aba Francis and Aba-Ori-Apata near Ikoru Ekiti. Texturally, they are medium to coarse grained with alternating bands of light and dark coloured portions of about 3cm thickness with complete gradation between them. They occur as low lying outcrops which have been intruded by pegmatite in the south eastern corner of the area. The rock consists of alternating felsic and mafic minerals and structures like banding, microfolds and fractures were noticed on the outcrop. They have whitish grey tones. The samples mapped around Okemesi and Ipoti are medium to coarse grained and are foliated and display compositional banding of the mafic and felsic minerals. The rock strikes at 333° and dips at $60^{\circ} W$. The mineralogy consist of quartz, biotite and orthoclase and microcline feldspars.

Quartzites

These tend to form good topographical features which rise up to about 400metres above the surrounding terrains forming ridges. Quartzites cover the northern part around Oke-Ila, Ilupeju areas to central, western (Ayegunle), extending to the southwestern and southern parts of Okemesi (Ajindo) of the study area. Around the study area, the varieties of quartzite encountered are massive, milky, smoky, sugary, ferruginous and schistose quartzites. However, schistose, ferruginous and smoky varieties are the commonest in Okemesi area. The massive quartzites are not foliated but hard and are compact, the schistose quartzites are foliated and exhibit alternations of felsic mineral such as quartz and mafic mineral such as biotite with planar fabric.

The milky ones have milky appearance by inspection and developed slickenside surfaces with specks of muscovite. The smoky ones are formed by different oxidation states of iron in the crystal lattices of the rock or due to impregnations of some transition elements during the rock formation. The sugary ones have granular textures and are friable when struck with hammer. The ferruginous types are rich in iron. They are very common in Okemesi and Itawure. The quartzites have varying textures from equigranular, medium grained to coarse-grained. The varieties of quartzites are closely related that, often, it is impossible to indicate them as separate units on the map. The quartzites consist of mainly quartz which is usually more than 70% with minor amounts of interlocking grains of biotite and orthoclase. Structurally, the quartzites are jointed, with some having joint sets around Ayikunnugba (Oke-Ila) area, while others are foliated. Dips ranging between 400W-660 W were measured around Soso and Oko Ajindo areas. In some areas for instance, they have very high dips ranging from 660 E to almost horizontal at Okemesi, most especially along the limbs and fold closure. The quartzites in the area have orange-yellow colour due to mineral impurities. Quartz vein is the dominant structural feature on the rocks in this locality.

Biotite-Gneisses

This is a foliated, medium to coarse-grained, dark to almost black coloured rock composed chiefly of biotite and little quartz. It occupies the north central area and extends towards the north western part of the study area. A band of biotite gneiss concordantly lies within the quartzites around Ajindo area. The rock has been severely weathered and covered with sand thereby making field observation difficult to carry out. The varieties seen at Ikoru and Ijero are the highly foliated type with bands of black tints imposed by biotite impregnations alongside felsic minerals such as quartz and plagioclase feldspar. Most of the mineral alignments are conformable with the foliation planes of the adjacent schistose rock.

Granite-gneiss.

The granite gneisses are common in Okemesi and Epe. They exhibit mineralogical banding of felsic and mafic minerals. The felsic minerals are the quartz and feldspar (plagioclase) and mafic minerals are biotite. The texture is medium to coarse grained.

Biotite-schists

The biotite schists encountered in the study area occurs around Arapate Erigbe and Soso area of Ekiti State as a lenticular body within the quartzites and migmatite gneiss, and it is exposed due to stream activity as low lying outcrop. It has undergone

various levels of deformation. Structurally, foliation is present thereby making the name "biotite schist" appended to the rock as confirmed by petrographic studies. Other structural features on the rock include microfolds and joints which control the stream flow. Field observations showed that this rock dips at 480 W to the surrounding rocks. The rock is medium grained and contains quartz and biotite which is the dominant mineral, and it is dark grey in colour. The rock also occurs in Ikoru and Ijero area as a schistose rock with grayish colour, and with black patches of biotite. It exhibits fine grained texture. It covers nearly two-thirds of Ijero area with pegmatite intrusions along Ijero-Ikoru road. It is also found in Arapate/Soso area with structures such as foliations and microfolds. It contains quartz and biotite.

Quartz-Biotite Schists

This group of rock occurs in lowland areas between quartzite and banded gneiss around Oko-Esinkin2 area (eastern part of Okemesi) where it has been exposed by stream channel and road cut. The rock has been highly deformed with the adjacent migmatite-gneiss-quartzite complex, the foliations on the outcrop is defined by biotite streaks. The rock has medium to coarse grained texture and consist mainly of biotite, quartz, plagioclase feldspar and orthoclase feldspar. Structures found on the outcrop are fractures, and the rock is grey to dark in colour.

Pegmatites

This lithologic unit ranges from a few meters in length and is located in the southeastern part of Okemesi where it intrudes the banded gneiss around Aba Francis and Ikoru area as an isolated hill. Texturally, it is extremely coarse-grained with quartz, feldspar and muscovite as distinguished mineral component. Quartz vein, joints and veinlets are the structures observed on the outcrop. Based on field observation, the pegmatite is the complex type with distinct textural and mineralogical variations with an impure white colouration. It is the youngest rock in Ikoru area, unlike the Okemesi pegmatite which is the simple type due to uniform variation of its constituent minerals. Pegmatite intrusions also occur towards the south-western part very close to the centre of Ijero town, which is also complex. It extended to the quarry site in Ijero along Ijero-Ipoti road. The pegmatite in Ijero area is zoned, consisting of massive quartz at the core, and followed by mica schists with smoky quartz impregnated with tourmaline and purplish quartzite. In Ara and Epe area of Ijero, the pegmatites here intruded into the biotite schist and migmatite gneiss that occupies the central part of the area, covering about three-quarter of the total land mass. The pegmatites in this area are very coarse grained

igneous type with phenocrysts over 250mm in length, usually of granitic composition. The pegmatite in Ipoti and Odo Owa areas are associated with cassiterite-tantalite mineralization (Fadipe, 1988). They are found associated with migmatite gneiss and amphibolites with very coarse grained texture and consists of feldspar, tourmaline and garnet while pegmatites in Ijero is enriched with unusual trace elements which thus result in the crystallization of unusual rare minerals such as beryl, tourmaline, columbite and tantalite. The pegmatites in Ijero occur as a low but large elongate hill of average height of about 50m above sea level. Some of the pegmatites in Ijero have been and are still being worked on most especially at Ijero and other villages such as Ikoro, Odo-Owa, Saloro and Oke Asa. The dominant minerals are orthoclase, albite, quartz, microcline and biotite.

Calc-gneiss

This is a typical gneissic rock with abundance of calcium. It exhibits a characteristic black colour with white fragments of quartz. The texture of the rock is mostly fine grained. Joints are the major structures discovered on the outcrop, and are common in most parts of Ikoro. Part of the lithological unit is exposed at Odo agba (Ikoro) where the stream takes its course. They are medium to coarse grained. It is also common in Ijero where it is composed chiefly of calcite and quartz. It occurs mainly as nodules and discontinuous streaks up to four inches in thickness. It is also made up of a mosaic of twinned grains of calcite enclosing isolated rounded crystals or composite spots of silicate minerals.

Granites

Varieties of granites based on mineral composition, texture and grain sizes are very prominent in the south eastern part of the study area most especially in Ofale, Osun/ Epe, Iroko, and Idao parts of the study area and form well defined boundaries with the quartzites. Their textures range from fine, medium to coarse grained. They occur as hilly, low lying, flat and extensive outcrops in most area with sparse vegetations. Structures common the outcrops include quartz vein, veinlets, pegmatite dykes which trend north-south, exfoliation, folds of different styles, xenoliths etc. A typical granitic rock must have > 60% of quartz to be termed an acid igneous rock. However, the granites occur as pockets of rocks within the biotite gneiss in Ikoro and Ijero area. They have colour variations ranging from specks of whitish, grayish to ash colour and patches of dark colour indicating ferromagnesian minerals.

Mica schists

The mica schists extend across most part of Ipoti, Odo Owa and Ijero, but occur prominently around Ipoti and Ijero. They are highly susceptible to weathering and erosion thereby reducing the quantity of fresh samples. In these areas, quartz-muscovite and quartz-muscovite-biotite schist are exposed in many places and have been highly pegmatized.

Migmatite-gneiss complex

This is presumably the oldest group of rocks and the most wide spread of all the lithologies, occupying 30% of the total surface area of Ijero and its surroundings. Its concordant lithostratigraphic relationship with the juxtaposed quartzite at Ayegunle gives credence to its probable metasedimentary origin. It is a mixed rock with a characteristic nature of a typical metamorphic rock which has taken an igneous character through partial melting. The migmatite has a mineralogical composition of quartz, orthoclase feldspar, hornblende and mica. It is not widely distributed in Ara Epe area unlike the pegmatites, but occupies the northern part. It is characterized mostly by alternating light and dark bands of minerals. In Ipoti, Odo Owa area, the rock has been weathered in-situ to give rise to high quality clays of economic importance, whereas, it occurred as a fresh low-lying outcrop in Ijero. It also possesses schlieren texture which is an example of granite formation in migmatites. This is typical of the samples picked from Epe. Migmatites in Epe are characterized by pygmatic folds and are believed to exhibit ductile deformation of the gneissic banding. Structures like folds and microfaults were also seen on the outcrop. The strike value of the rock is 3400 and dip is 320W.

Charnockites

The charnockites occur as an intrusive body north of Odo Owa within the massive quartzites and the gneisses as a coarse grained variety of charnockites which are foliated. They vary in texture and petrography and range from banded types to fine and medium grained. Though, the coarse grained varieties are common around Ipoti and Odo owa area. They also occur as discrete and individual bodies within the migmatite-gneiss-quartzite complex. Mineralogically, it consists of hornblende, biotite with little or no quartz. Structures seen on the rock are solution holes, quartz veins. The various rock units mapped in the course of field work were compiled to produce a geologic and cross-section map of the study area (Fig.2)

Sample ID	TiO ₂	Al ₂ O ₃	MgO	CaO	Na ₂ O	K ₂ O	WIP	PIA	CIW	CIA
Alpa Francis	0.002	0.003	0.033	0.028	0.003	0.024	0.285	-215.37	9.315	5.439
Baba Orskok	0.007	0.011	0.033	0.284	0.212	0.218	58.274	248.19	58.274	12.552
Baba Orskok Sb	0.042	0.075	1.025	1.875	0.010	3.257	50.218	72.29	10.804	12.552
Ida 2	0.354	0.668	1.225	2.043	3.760	4.736	63.981	234.52	10.324	5.061
Ida 2a	0.452	0.824	2.050	3.188	3.188	0.927	53.577	759.89	13.041	8.011
Ida 2b	0.048	0.091	0.033	0.028	0.127	0.422	4.275	312.46	28.915	12.385
Ida Lepidom	0.048	0.091	0.033	0.028	0.127	0.422	4.275	312.46	28.915	12.385
Ida neptun	0.017	0.032	0.005	0.112	0.170	1.028	15.023	117.45	10.058	14.052
Ida 2A	0.003	0.008	0.033	0.490	7.007	2.012	56.875	-36.54	0.081	0.066
Ida 2B	0.003	0.006	0.000	0.000	0.006	0.024	0.000	-62.40	12.088	8.266
Ida 2C	0.018	0.035	0.000	0.000	0.061	0.133	0.000	262.53	36.361	15.210
Ida 2D	0.003	0.006	0.000	0.000	0.028	0.098	0.000	145.81	18.208	4.810
Ida 2E	0.003	0.007	0.000	0.000	0.028	0.301	0.000	144.47	37.626	11.026
Ida 2F	0.522	0.986	1.227	2.895	3.288	3.675	53.690	76.95	13.752	9.024
Ida 2G	0.024	0.047	0.000	0.000	0.028	0.157	0.000	216.88	30.238	15.023
Ida 2H	0.082	0.154	0.192	1.329	2.625	4.230	50.961	3390.19	3.757	1.831
Ida 2I	0.450	0.851	0.763	2.700	4.161	3.049	52.873	47.13	11.031	7.806
Ida 2J	0.024	0.047	0.000	0.000	0.028	0.157	0.000	216.88	30.238	15.023
Ida 2K	0.377	0.712	0.713	1.399	2.600	4.061	50.158	-514.77	15.114	8.118
Ida 2L	0.095	0.180	0.105	0.546	2.058	4.422	53.874	674.42	4.227	2.189
Ida 2M	0.003	0.007	0.000	0.000	0.028	0.301	0.000	144.47	37.626	11.026
Ida 2N	0.552	1.043	0.912	2.965	3.205	4.121	57.106	-96.40	14.259	9.120
Ida 2O	0.191	0.338	0.033	0.042	0.031	0.111	1.220	204.20	24.120	12.960
Ida 2P	0.077	0.145	0.000	0.000	0.064	0.458	0.000	126.75	0.000	0.000
Ida 2Q	0.332	0.627	0.716	0.000	0.030	1.964	0.000	102.27	0.000	0.000
Ida 2R	0.003	0.006	0.000	0.000	0.028	0.301	0.000	144.47	37.626	11.026
Ida 2S	0.277	0.523	0.278	0.588	1.291	4.868	48.251	176.19	21.775	7.194
Ida 2T	0.191	0.338	0.033	0.042	0.031	0.111	1.220	204.20	24.120	12.960
Ida 2U	0.203	0.384	0.829	0.204	0.915	4.579	43.203	140.50	24.123	6.228
Ida 2V	0.150	0.284	1.011	0.797	1.717	3.940	43.599	320.21	10.134	4.208
Ida 2W	0.003	0.007	0.000	0.000	0.028	0.301	0.000	144.47	37.626	11.026
Ida 2X	0.364	0.682	0.514	2.266	2.084	4.157	66.176	-89.21	8.336	5.628
Ida 2Y	0.024	0.047	0.000	0.000	0.028	0.157	0.000	216.88	30.238	15.023
Ida 2Z	0.077	0.145	0.000	0.000	0.067	0.169	0.000	-54.45	0.000	0.000
Ida 2AA	0.053	0.101	0.050	0.028	0.059	0.504	4.283	163.40	30.070	13.194
Ida 2AB	0.012	0.022	0.000	0.000	0.026	0.024	0.000	-5.67	0.000	0.000
Ida 2AC	0.000	0.000	0.000	0.000	0.020	0.000	0.000	0.000	0.000	0.000

Table 2: Weathering indices of stream sediments taken from the study area (%)

Sample ID	Al ₂ O ₃	MgO	CaO	Na ₂ O	K ₂ O	WIP	PIA	CIW	CIA
Alpa Francis	4.042	0.066	0.252	0.062	0.289	3.101	92.284	92.796	87.019
Baba Orskok	7.025	0.215	0.938	1.010	1.119	7.285	86.463	86.660	72.661
Baba Orskok Sb	10.578	0.199	0.330	1.004	0.085	47.140	80.226	88.653	62.162
Baba Orskok Sc	12.411	0.398	0.700	2.049	4.622	50.167	75.804	81.869	64.724
Ida 1	16.661	0.812	0.266	0.670	4.157	38.569	93.017	94.692	76.588
Ida 2	13.242	0.298	0.308	0.522	3.037	28.045	92.483	94.105	77.402
Ida 2a	4.647	0.050	0.070	0.034	0.181	1.794	97.732	97.818	94.233
Ida 2b	4.607	0.050	0.000	0.013	0.186	1.759	97.495	97.691	93.919
Ida 2c	1.341	0.000	0.042	0.105	0.458	4.324	85.723	90.115	68.913
Ida 2d	0.726	0.000	0.000	0.036	0.192	1.750	93.926	95.405	78.727
Ida 2e	1.967	0.050	0.000	0.013	0.186	1.759	97.495	97.691	93.919
Ida 2f	1.379	0.000	0.042	0.038	0.096	1.047	94.149	94.525	88.675
Ida 2g	8.406	0.116	0.148	0.216	0.615	14.693	94.526	95.510	80.718
Ida 2h	2.512	0.216	0.168	0.103	0.759	7.169	86.463	86.660	72.661
Ida 2i	3.721	0.149	0.238	0.286	1.398	15.321	81.609	87.665	65.949
Ida 2j	2.58	0.054	0.062	0.042	0.193	1.074	95.691	95.357	82.357
Ida 2k	6.819	0.298	0.462	0.249	3.145	29.156	74.819	87.552	62.827
Ida 2l	17.681	0.116	0.224	2.596	1.012	40.521	74.123	72.237	72.130
Ida 2m	7.556	0.249	0.518	0.890	3.699	24.554	76.151	86.218	60.627
Ida 2n	5.856	0.415	0.546	0.786	1.747	19.709	75.525	81.475	65.242
Ida 2o	1.473	0.182	0.168	0.069	0.627	5.848	78.160	86.162	65.057
Ida 2p	13.289	0.862	0.126	0.124	2.049	18.234	97.827	93.135	85.264
Ida 2q	0.793	0.000	0.000	0.022	0.217	1.858	96.394	97.533	76.889
Ida 2r	0.112	0.000	0.000	0.011	0.117	0.924	96.661	97.116	82.454
Ida 2s	1.096	0.000	0.000	0.026	0.349	2.942	96.661	97.716	74.497
Ida 2t	2.267	0.066	0.084	0.117	0.778	5.491	89.351	91.847	74.409
Ida 2u	1.096	0.000	0.000	0.013	0.131	2.454	96.661	97.116	82.454
Ida 2v	2.342	0.000	0.036	0.151	0.856	7.787	87.782	91.883	68.705
Ida 2w	1.322	0.000	0.000	0.035	0.494	4.153	95.940	97.418	71.422
Ida 2x	3.227	0.000	0.000	0.017	0.219	29.122	65.919	65.919	65.919
Ida 2y	6.083	0.249	0.196	0.443	2.109	19.960	86.141	90.489	68.880
Ida 2z	7.065	0.033	0.126	0.481	4.651	40.177	79.801	92.086	57.539
Ida 2aa	3.242	0.108	0.126	0.124	1.171	11.118	82.819	82.819	82.819
Ida 2ab	5.459	0.133	0.222	0.429	2.085	19.634	83.219	88.917	66.379

Geological field mapping

Ten distinct lithological units were recognized in the studied area which were compiled to produce a geological map (Fig.2) which are the schistose and the massive quartzites occurring as quartzite ridge (Effon Psammite Formation) which runs north-south, granites occurring as plutons and intrusions, calc-gneiss as intrusions within the migmatite and the quartzite while the schists and migmatites occurred as distinct rock groups. The schists, quartzites and migmatites are interbanded. Other lithologies include the gneisses, pegmatites and charnockites occupying different portions of the study area with well-defined boundaries. The major structure in the area is an antiform which is visible on the schistose quartzites in the studied area.

Weathering Profile of the crustal rocks

Table.1 represents the weathering indices of the crustal rocks from the study area. The weathering index assessment of the rocks was carried out to ascertain whether the rock samples collected from the various locations in the studied area are fresh or weathered. This is to justify the reliability of the geochemical analytical results of the rocks. The weathering indices employed is WIP, PIA, CIW and CIA respectively. The CIA values range from

0.006% -15.10% which indicates that the crustal rocks collected from the study area are largely unweathered (fresh rocks).The CIW and PIA values show a positive correlation with CIA. The WIP shows a negative trend, indicating that the higher value of WIP is a reflection of rocks that are largely fresh samples. Therefore, the rock samples used for this study are fresh samples which give a good result for the interpretation of the nature and the geological processes in the study area.

Weathering Profile of sediments in the source area

The weathering profile of the sediments is presented in Table 2. Weathering indices are useful tools to illustrate weathering profiles and ascertain the degree of weathering of the sediments and minerals incorporated in them. Some minerals are accumulated as deposits due to the process weathering and transportation. Most heavy minerals forming placer or paleo-placer deposits are products of weathering from the parent rock. Therefore, it is necessary to ascertain the level and extent of weathering of the sediments to ascertain their usefulness in mineral genesis and exploration. The weathering indices used in this study to examine the decomposition of unstable minerals are (i). Chemical index of Alteration (CIA) Nebbit & Young (1982) (ii).Weathering index of Parker (WIP) (iii) Plagioclase index of Alteration (PIA)

(iv) Chemical index of weathering (CIW) Harnois (1988).The chemical index of alteration (CIA) is defined as $CIA=100*Al_2O_3/(Al_2O_3+CaO +Na_2O+K_2O)$,Chemical index of weathering (CIW) = $Al_2O_3/CaO + Na_2O$, Weathering Index of Parker (WIP) = $2Na_2O/0.35 + MgO/0.90 + 2K_2O/0.25 +CaO/0.70$ and Plagioclase index of Alteration (PIA) = $100*(Al_2O_3 -K_2O)/Al_2O_3 + Na_2O + CaO -K_2O$. In the formular given above CaO* is the amount of calcium oxide in incorporated in the silicate fraction of the studied stream sediment smpls. Corrections for CaO from carbonate contribution were not done for the studied samples since there was no CO2 data. Accordingly, to compute for CaO* from the silicate fraction, the assumption proposed by Burke et al, (1988) was adopted. In this regard, CaO values were accepted only if CaO is less than equal to Na2O. For the CIA, the low CIA of approximately 50% implies an unweathered upper crust or weak weathering, but high CIA value such as 76% - 100% indicates intense weathering with a complete removal of alkali and alkaline earth elements and an increase in alumina according to McLennan (1993), Fido et al, (1995) and Dupuis et al, (2006). The weathering profile of the stream sediment showed that the least of CIA in the stream sediments collected from the study area ranges from

56.94% -94.23% (Table.2). The weakly weathered stream sediments were collected at these two locations namely River Elewu and Issa 7C area.

From Table 2, stream sediment samples from Effon 1 and Okemesi 3C were intensely weathered with CIA values of 91.84% and 94.23% respectively, including the samples from Agbagbara, Ede 1, Ede 2, Effon 1, Effon-Okemesi 3C, Effon-Okemesi 2, Esa-Oke 2, Esa-Oke 3, Esa-Oke 6, Ido-Ile 2, Odo-owa Abidogun, Ofale Rd, Okemesi2&3 and lastly Okemesi9. These are intensely weathered stream sediments. They do not contain stable minerals grains according to their CIA values. The extent of weathering at the source area can also be determined by Plagioclase index of alteration (PIA). The plagioclase index of alteration shows corresponding values with CIA (Table.2). The high PIA value indicates intense weathering of the plagioclase feldspars from the parent materials. The chemical index of weathering (CIW) shows a positive correlation with the CIA and PIA. This showed that all the listed locations above that have intense weathering are also revealed by chemical index of weathering in Table 2. The weathering index of Parker (WIP) showed a negative correlation with CIA, PIA and CIW, indicating that the low values of WIP is a reflection of intense weathering of the sediments in the source area.c

6. Conclusion

Detailed geological mapping has provided an insight into the nature and disposition of the different rock units in the area. The geology of the study area from field examination confirmed the earlier assertion of Hubbard (1975), who described the geology of the area as being characterized by a thrust interface, known as Ifewara fault which separated the Ilesha schist belt into two groups of contrasting lithologies. The geological mapping revealed the dispositions of the various rock units in the area such as the migmatites which are the oldest rocks in the study area upon which other rocks such as granites, calc gneisses and mica-schists intruded. The massive and schistose quartzites, although of the same rock units do not exist together in the area, they are interbanded with the gneisses and migmatites or occur as distinct rock units with massive topography.. The strike values of the quartzites (schistose and massive) range from 0240-0460 in some places. Also, the rocks dip in the western direction, with values such as 400W - 800W in some areas of study, while other areas it dips in the eastern direction with dip values ranging between 720E - 800E respectively. The high dip values could be attributed to several episodes of deformation that characterize the rocks in the area which is manifested in the brittle nature of the quartzites that display

several joints and fracture sets which also control the drainage pattern in the area. The weathering index analysis of the rocks also confirmed that the rock samples used for this research are fresh using the weathering indices parameters while some sediment samples collected for geochemical analysis also ranges from weakly weathered to intensely weathered which may not be totally suitable for mineral exploration purposes.

7. Recommendation

Detailed geochemical exploration for gold and other precious metals is desirable in the study area because of its geologic and thermo-tectonic setting.

8. References

- [1] Akintola, A.I., Ikhane, P.R., Bankole, S.I., Adeokurolere, D.N. (2013). Petrography and Stream sediment geochemistry of Ede and its environs, southwestern Nigeria. *International Research Journal of Geology and Mining*, 3(6):2276-6618.
- [2] Burke, K.C., Dewey, J.F. and Kidd, W.S.F. (1988). World distribution of Sutures; the sites of former oceans. *Tectonophysics*, 40: 69-99.
- [3] British Geological Survey (BGS). (1990). *Regional Geochemical Atlas: Argyll*. British Geological Survey.
- [4] Darnley, A.G. (1990). International Geochemical Mapping: A new global project. *Journal of Geochemical Exploration* 39 : 1-13.
- [5] Dupuis, C, Hebert, R, Cote, VD, et al., (2006). "Geochemistry of sedimentary rocks melange and flysch units south of the YarlungZangbo suture zone, southern Tibet". *Journal of Asian Earth Sciences* 26: 489- 508.
- [6] Emmanuel, A. Ariyibi, Samuel., L. Folami, Bankole., D. Ako, Taye., R. Ajayi, Debowale.,O. Adelus (2011). Applications of the principal component analysis on geochemical data; A case study in the basement complex of southern Ilesha area, Nigeria. *Arab Journal of Geoscience* 4: 239-247.
- [7] Fadipe, A.A (1988). Chemical composition of Niobium-Tantalite from granitic pegmatite and alkali granites (Egbe and Ijero mineralized region). *Journal of Mining and Geology*, 7: 42-47.
- [8] Fedo, C.M., Nesbitt, H.W., Young, G.M. (1995). "Unraveling the effects of potassium metasomatism sedimentary rock sand paleosols, with implications for paleoweathering conditions and provenance". *Geology*, 23: 921- 924.
- [9] Harnois, L. (1988). The CIW index: A new chemical index of weathering. *Sedimentary Geology*, 55, 319-322.
- [10] Johnson, C.C., Breward, N., Ander, E.L., Ault, L., (2005). G-BASE: baseline geochemical mapping

of Great Britain and Northern Ireland. *Geochemistry: Exploration, Environment Analysis*, 5 : 347-357.

[11] Key, R.M., De Waele, B., Liyungu, A.K., 2004. A multi-element baseline geochemical database from the western extension of the Central African Copperbelt in north western Zambia. *Applied Earth Sciences (Transn. Inst. Min. Metall. B)*, 113, 205–226.

[12] McLennan, S.M. (1993). "Weathering and global denudation". *Journal of Geology*, 101, 295-303.

[13] Mumbfu, E.M., Nforba, M.T., Cheo, E.S. (2014). Geochemical dispersion of gold in stream sediment in Paleoproterozoic Nyong Series, Southern Cameroon. *Science Research*. 2(6):155-165.

[14] Nesbitt, HW, and Young, G.M. (1982). Early Proterozoic climates and plate motions inferred from major element chemistry of lutites. *Nature*, 299, 715–717.

[15] Oyinloye, A.O. and Adebayo O.F.(2005). Petrophysical and chemical properties of Calc-gneiss and clay deposit in Ijero-Ekiti area as industrial raw materials. *Global Journal of Mechanical Engineering*. 6 (1): 30-34

[16] Olaolorun, O.O., Oyinloye, A.O (2010). Geology and geotechnical appraisal of some clay deposits around Ijero Ekiti, Southwestern Nigeria: Implication for Industrial uses. *Pakistan Journal of Science and Industrial Research*. 53: (3), 127-135.

[17] Oyinloye, A.O. (1997). Physical and chemical characteristics of residual clay deposits in Ijero Ekiti. *Journal Technoscience* 1(1): 40-45.

[18] Rahaman, M.A. (1988). Recent advances in the Study of the basement complex of Nigeria. In: *Precambrian Geology of Nigeria*. Geological Survey of Nigeria. 11-43.

[19] Reimann, C. and Caritat, P. D., (1998). *Chemical elements in the environment*. Springer-Verlag, New York.31: 75-85.

[20] Rice, K.C., (1999). Trace-element concentrations in streambed sediment across the contemporaneous United States. *Environment Science and Technology*., 33(15):2499-2504.

[21] Watts, D.W. and Johnson, M.H., (2010). Geochemical mapping using stream sediments in west-central Nigeria. Implications for environmental studies and mineral exploration in west Africa. *Applied geochemistry*, 27 (6): 1035-1062.