

# Temporal controls on dissolved organic matter and lignin biogeochemistry in a pristine tropical river, Democratic Republic of Congo

Robert G. M. Spencer,<sup>1</sup> Peter J. Hernes,<sup>2</sup> Rosmarie Ruf,<sup>3</sup> Andy Baker,<sup>4</sup> Rachael Y. Dyda,<sup>1,2</sup> Aron Stubbins,<sup>5</sup> and Johan Six<sup>1</sup>

Received 18 October 2009; revised 11 February 2010; accepted 10 March 2010; published 6 August 2010.

[1] Dissolved organic carbon (DOC), lignin biomarkers, and the optical properties of dissolved organic matter (DOM) were measured in the Epulu River (northeast Democratic Republic of Congo) with the aim of investigating temporal controls on the quantity and chemical composition of DOM in a tropical rainforest river. Three different periods defined by stages of the hydrologic regime of the region, (1) post dry flushing period, (2) intermediary period, and (3) start of the dry period/post flush, were sampled. Temporal variability in DOM quantity and quality was observed with highest DOC, lignin concentration ( $\Sigma_8$ ) and carbon-normalized ( $\Lambda_8$ ) values during the flushing period attributed to greater surface runoff and leaching of organic-rich horizons, with lowest values in the dry period/post flush once source materials were well leached. Chromophoric DOM (CDOM) was strongly correlated to DOC and  $\Sigma_8$  ( $r^2 = 0.85$  and  $0.83$ , respectively;  $p < 0.001$ ), and CDOM quality measurements ( $SUVA_{254}$ , spectral slope ratio and fluorescence index) were strongly correlated to  $\Lambda_8$  values ( $r^2 = 0.77$ ,  $0.69$ , and  $0.75$ , respectively;  $p < 0.001$ ), demonstrating the ability to derive DOC and lignin export and to track DOM quality in tropical riverine systems from simple optical measurements. This study demonstrates similar effects in the variability of DOM quantity and quality due to changing hydrologic inputs for a tropical river as has been previously reported for temperate and northern high-latitude rivers. Therefore, flushing periods in tropical rivers warrant further study, as they are critical toward understanding ecosystem biogeochemistry as maximal export of freshly leached plant material occurs during this time period.

**Citation:** Spencer, R. G. M., P. J. Hernes, R. Ruf, A. Baker, R. Y. Dyda, A. Stubbins, and J. Six (2010), Temporal controls on dissolved organic matter and lignin biogeochemistry in a pristine tropical river, Democratic Republic of Congo, *J. Geophys. Res.*, 115, G03013, doi:10.1029/2009JG001180.

## 1. Introduction

[2] Dissolved organic matter (DOM) represents a fundamental link between terrestrial and aquatic carbon cycles and plays an essential role in aquatic ecosystem biogeochemistry. The majority of DOM studies in catchments have focused on measuring bulk concentrations such as dissolved

organic carbon (DOC) and dissolved organic nitrogen (DON) in order to calculate export fluxes and observe temporal trends. Such bulk measurements, however, provide little information about the chemical composition or quality of DOM and, thus, its biogeochemical role [Hernes *et al.*, 2008; Baker *et al.*, 2008a; Fellman *et al.*, 2009]. To examine biogeochemical processes in relation to DOM quality previous studies have focused on measurements of bulk DOM properties such as C/N ratios, isotopic composition, and chromophoric (colored) DOM (CDOM) [Hood *et al.*, 2005, 2007; Neff *et al.*, 2006] or measurements of the molecular composition of DOM such as amino acids, carbohydrates, and lignin phenols [Hedges *et al.*, 1994, 2000; Lobbes *et al.*, 2000]. CDOM optical properties, such as absorption and fluorescence, have been the focus of numerous current studies in aquatic sciences due to their potential as proxies for understanding biogeochemical processes such as lability and photochemical degradation [Cory *et al.*, 2007; Stubbins *et al.*, 2008; Fellman *et al.*, 2009].

<sup>1</sup>Department of Plant Sciences, University of California-Davis, USA.


<sup>2</sup>Department of Land, Air, and Water Resources, University of California-Davis, USA.

<sup>3</sup>The Okapi Breeding and Research Station, Institute in Congo for the Conservation of Nature, Gilman International Conservation, Epulu, Orientale Province, Democratic Republic of Congo.

<sup>4</sup>Connected Waters Initiative, Water Research Laboratory, University of New South Wales, Manly Vale, New South Wales, Australia.

<sup>5</sup>Department of Chemistry and Biochemistry, Old Dominion University, Norfolk, Virginia, USA.





**Figure 2.** Annual variation in mean monthly rainfall data [adapted from *Katuala et al.*, 2005] in the Okapi Wildlife Reserve (Ituri Forest).

2000] and have received little attention with respect to temporal variability in DOM quality. The second largest area of tropical rainforest in the world is found in central Africa, and within this rainforest, the Congo River Basin includes vast and unbroken regions from the Atlantic Ocean to the Albertine Rift [*Duveiller et al.*, 2008]. Political instability and limited infrastructure have severely inhibited research efforts in this region [*Baccini et al.*, 2008], thereby limiting our fundamental knowledge of biogeochemical cycling in the Congo River Basin. This study examined the Epulu River, a typical river draining pristine tropical rainforest in central Africa. The aim of this study was to investigate temporal variation in the quantity and chemical composition of DOM, which in turn was used to identify source materials and to address how important seasonality is in the biogeochemical cycling of DOM derived from tropical riverine ecosystems.

## 2. Materials and Methods

### 2.1. Study Site

[5] The Epulu River was sampled near the Okapi Breeding and Research Station (1°24'N, 28°34'E) near Epulu in the Okapi Wildlife Reserve in the northeast of the Democratic Republic of Congo (Figure 1). The Okapi Wildlife Reserve is a World Heritage Site that covers an area of ~13,700 km<sup>2</sup> in the Ituri Forest. The Epulu River is one of two main rivers that drain the Okapi Wildlife Reserve in the Ituri Forest. It joins the other main river (the Ituri) and takes its name as it flows westward and later becomes the Aruwimi River, which continues westward and joins the Congo River at Basoko. The Ituri Forest (~70,000 km<sup>2</sup>) exists in a pristine state and due to the density of its vegetation, has among the highest carbon content per hectare of any rainforest [*Koenig*, 2008]. Vegetation within the Ituri Forest is dominated by

two types of tropical moist forest (mixed forest and monodominant forest) with small areas of swamp forest in areas of poor drainage [*Makana et al.*, 2004; *Katuala et al.*, 2005]. Human impacts within the region are minimal (e.g., traditional hunting-gathering), and the population density is low (~1.5 persons/km<sup>2</sup>). A lack of transportation and recent conflict in the region has restricted logging to the forest edge and the banks of navigable rivers [*Makana et al.*, 2004]. Within the central African countries, the Democratic Republic of Congo has the lowest amount of forest disturbance within its borders, with just 1% currently disturbed [*Laporte et al.*, 2007].

[6] Climatic records for the region are not detailed, and no discharge data exist for the rivers in the Ituri Forest; however, hydrologic regimes can be examined by utilizing existing rainfall data. The annual rainfall is estimated to be between ~1300 and 2100 mm (typically 1650–1800 mm), with April, May, and August through October as the wettest months while December, January, and February (the dry season) typically receive less than 100 mm of precipitation per month [*Makana et al.*, 2004; *Katuala et al.*, 2005, and references therein]. Mean monthly rainfall data collected in Epulu from 2003 to 2008 follows the long-term trends observed previously [*Katuala et al.*, 2005, and references therein] (Figure 2) and the mean annual rainfall for 2003–2008 was ~1670 mm. Temperature shows little variation in the region with a mean annual temperature of ~24°C (mean monthly temperature ~22.8–24.6°C) [*Katuala et al.*, 2005, and references therein].

### 2.2. Water Sample Collection and Processing

[7] Water samples were collected during three different periods of the year (9 or 10 days either consecutive to one another or within a 12 day period). The three sampling periods corresponded to different stages of the hydrologic regime of the area (flushing period, intermediary, and the start of the dry period/post flush). The high-discharge flushing period was sampled at the start of the April/May wet period (16–27 April 2008) for 10 days out of the 12 day period. The intermediary period was sampled daily toward the end of the dry season (February) when individual small storms occurred (5–13 February 2008). The post flush/onset of the dry period was sampled for 10 days after some of the wettest months (August through October) at the onset of the dry season, typically during the middle to end of November/beginning of December (28 November to 6 December 2007). Water samples were collected from the surface of the river in midchannel via a precleaned and prerinced 1 L Nalgene bottle and filtered immediately through pre-combusted (450°C) 0.7 μm glass fiber filters (Whatman, GF/F). Samples were stored frozen, kept in the dark, and shipped on ice to the University of California-Davis (USA) for further analyses.

### 2.3. Dissolved Organic Carbon and Chromophoric Dissolved Organic Matter Analyses

[8] Dissolved organic carbon was measured on filtered water samples using a high-temperature catalytic oxidation (HTCO) Shimadzu TOC 5000A instrument and methodology as described in *Spencer et al.* [2007a]. All reported DOC data are the mean of three to five replicate injections

**Table 1.** Epulu River Dissolved Organic Carbon, Chromophoric Dissolved Organic Matter, Specific UV Absorbance at 254 nm, and Fluorescence Index Data<sup>a</sup>

Sampling Period	Date	DOC (mg L <sup>-1</sup> )	$S_{275-295}$	$S_{350-400}$	$S_R$	SUVA <sub>254</sub> (L mg C <sup>-1</sup> m <sup>-1</sup> )	FI
Wet/flushing	16 Apr 2008	9.0	0.01453	0.01824	0.796	3.57	1.22
Wet/flushing	17 Apr 2008	8.7	0.01464	0.01836	0.797	3.45	1.28
Wet/flushing	18 Apr 2008	8.5	0.01476	0.01828	0.807	3.34	1.24
Wet/flushing	19 Apr 2008	7.6	0.01481	0.01794	0.826	3.38	1.29
Wet/flushing	20 Apr 2008	6.8	0.01520	0.01771	0.858	3.34	1.30
Wet/flushing	21 Apr 2008	6.9	0.01512	0.01775	0.852	3.35	1.30
Wet/flushing	23 Apr 2008	8.0	0.01487	0.01806	0.823	3.43	1.27
Wet/flushing	25 Apr 2008	7.2	0.01492	0.01711	0.872	3.36	1.28
Wet/flushing	26 Apr 2008	6.9	0.01479	0.01707	0.867	3.34	1.29
Wet/flushing	27 Apr 2008	6.6	0.01463	0.01735	0.843	3.29	1.33
Intermediary	05 Feb 2008	7.4	0.01484	0.01716	0.865	3.33	1.30
Intermediary	06 Feb 2008	6.8	0.01459	0.01600	0.912	3.30	1.31
Intermediary	07 Feb 2008	6.0	0.01475	0.01604	0.920	3.27	1.38
Intermediary	08 Feb 2008	5.2	0.01491	0.01562	0.954	3.21	1.41
Intermediary	09 Feb 2008	5.5	0.01533	0.01571	0.976	3.17	1.44
Intermediary	10 Feb 2008	5.4	0.01497	0.01614	0.928	3.19	1.43
Intermediary	11 Feb 2008	7.8	0.01485	0.01662	0.893	3.29	1.27
Intermediary	12 Feb 2008	8.4	0.01443	0.01569	0.920	3.31	1.25
Intermediary	13 Feb 2008	8.0	0.01454	0.01595	0.912	3.29	1.25
Dry period/post flush	28 Nov 2007	5.3	0.01470	0.01518	0.968	3.09	1.45
Dry period/post flush	29 Nov 2007	6.0	0.01431	0.01464	0.978	3.12	1.42
Dry period/post flush	30 Nov 2007	5.5	0.01463	0.01484	0.986	3.08	1.45
Dry period/post flush	01 Dec 2007	5.3	0.01417	0.01354	1.046	3.09	1.45
Dry period/post flush	02 Dec 2007	5.4	0.01440	0.01352	1.066	3.13	1.43
Dry period/post flush	03 Dec 2007	5.4	0.01479	0.01409	1.050	3.11	1.42
Dry period/post flush	04 Dec 2007	6.6	0.01473	0.01498	0.983	3.15	1.39
Dry period/post flush	05 Dec 2007	5.6	0.01469	0.01459	1.007	3.16	1.41
Dry period/post flush	06 Dec 2007	5.7	0.01412	0.01497	0.944	3.20	1.41


<sup>a</sup>DOC, dissolved organic carbon; CDOM, chromophoric dissolved organic matter (spectral slope  $S_{275-295}$  and  $S_{350-400}$  and their ratio  $S_R$ ); SUVA<sub>254</sub>, specific UV absorbance at 254 nm; FI, fluorescence index.

with a coefficient of variance of less than 2%. For both CDOM absorbance and fluorescence measurements, all samples were measured in a 10 mm quartz cuvette at a constant laboratory temperature of 20°C, in duplicate and with ultrapure laboratory water (Milli-Q, Millipore) used as a blank. CDOM absorbance was measured with a Shimadzu UV-2501PC UV/Vis dual-beam spectrophotometer and Napierian absorbance coefficients ( $a$  m<sup>-1</sup>) were determined following *Hu et al.* [2002]. SUVA<sub>254</sub> values were calculated by dividing the UV absorbance ( $A$ ) at  $\lambda = 254$  nm by the DOC concentration (mg L<sup>-1</sup>) [*Weishaar et al.*, 2003]. Spectral slope ( $S$ ) was calculated using a nonlinear fit of an exponential function to the absorption spectrum over the ranges 275–295 and 350–400 nm. These ranges were chosen because *Helms et al.* [2008] observed in an extensive study of  $S$  across a range of aquatic ecosystems and DOM sources that the first derivative of the natural log spectra indicated the largest variations in these ranges. The spectral slope ratio ( $S_R$ ) was calculated as the ratio of  $S_{275-295}$  to  $S_{350-400}$  [*Helms et al.*, 2008]. DOM fluorescence spectra were measured on a Varian Cary Eclipse spectrofluorometer with a xenon excitation source and slits were set to 5 nm for both excitation and emission. A single emission scan was obtained from 400 to 550 nm at an excitation wavelength of 370 nm, and fluorescence intensity was blank corrected and also corrected for instrument specific bias and any inner filter effects [*Cory and McKnight*, 2005; *Spencer et al.*, 2007a]. Fluorescence index (FI) was then calculated as the ratio of fluorescence emission wavelength intensities at 470

and 520 nm at an excitation wavelength of 370 nm [*McKnight et al.*, 2001; *Cory and McKnight*, 2005]. Samples ranged in pH from 5.68 to 5.97 and, thus, were within the natural range typically observed in freshwaters in which the response of fluorescence to pH has been previously shown to be limited [*Ahmad and Reynolds*, 1999; *Spencer et al.*, 2007b]. As samples were stored frozen, the impact of freeze/thaw (thawing was conducted in the dark in a refrigerator at 4°C) was examined on DOC and CDOM parameters. After freezing and subsequent thawing DOC,  $a_{350}$ ,  $S_{275-295}$ ,  $S_{350-400}$ ,  $S_R$ , SUVA<sub>254</sub>, and FI exhibited changes typically within analytical error and always less than  $\pm 2\%$ .

#### 2.4. Lignin Phenol Analysis

[9] Lignin phenols were measured on whole water samples (~150 mL) acidified to pH 2 with 12 N HCl to minimize precipitation during rotary evaporation, and then rotary evaporated to ~3 mL. The concentrate was transferred to monel reaction vessels (Prime Focus, Inc.) and dried under vacuum centrifugation in preparation for lignin analysis. Lignin phenols were determined by alkaline CuO oxidation, followed by acidification and ethyl acetate extraction [*Hedges and Ertel*, 1982; *Spencer et al.*, 2008]. Quantification was carried out on a GC-MS (Agilent 6890 gas chromatograph equipped with an Agilent 5973 mass selective detector and a DB5-MS capillary column; 30 m, 0.25 mm inner diameter, Agilent) using cinnamic acid as an internal standard and a five-point calibration scheme



**Figure 3.** Box plots of (a) DOC and (b)  $a_{350}$  during the flushing, intermediary, and post flush sampling periods. The black dash-dotted line and the solid black line in the box represent the mean and the median, respectively. The horizontal edges of the box denote the 25th and 75th percentiles, and the error bars denote the 10th and 90th percentiles.

[Hernes and Benner, 2002]. Eight lignin phenols were quantified for all samples, including three vanillyl phenols (vanillin, acetovanillone, and vanillic acid), three syringyl phenols (syringaldehyde, acetosyringone, and syringic acid), and two cinnamyl phenols (*p*-coumaric acid and ferulic

acid). One blank was run for every 10 sample oxidations, and all samples were blank corrected. Blank concentrations of lignin phenols were low (30–50 ng) and, thus, never exceeded 2% of the total lignin phenols in a sample.

### 3. Results

#### 3.1. Dissolved Organic Carbon


[10] Epulu River DOC concentrations ranged from 5.2 to 9.0 mg L<sup>-1</sup> (mean = 6.7 mg L<sup>-1</sup>;  $n = 28$ ; Table 1 and Figure 3a) dependent on the time of sampling in the annual hydrological cycle. Concentrations of DOC from the dry period/post flush were consistently lower, ranging from 5.3 to 6.6 mg L<sup>-1</sup> (mean = 5.6 mg L<sup>-1</sup>;  $n = 9$ ), increasing in the intermediary period to 5.2–8.4 mg L<sup>-1</sup> (mean = 6.7 mg L<sup>-1</sup>;  $n = 9$ ), with highest DOC values in the flushing period ranging from 6.6 to 9.0 mg L<sup>-1</sup> (mean = 7.6 mg L<sup>-1</sup>;  $n = 9$ ) (Table 1 and Figure 3a).

#### 3.2. Chromophoric Dissolved Organic Matter

[11] Chromophoric DOM absorption coefficients at 350 nm ranged from 10.41 to 19.20 m<sup>-1</sup> (mean = 13.75 m<sup>-1</sup>;  $n = 28$ ; Figure 3b) in the Epulu River. Similar to DOC, CDOM varied seasonally, with  $a_{350}$  exhibiting highest values during the flushing period ranging from 13.12 to 19.20 m<sup>-1</sup> (mean = 15.38 m<sup>-1</sup>;  $n = 10$ ), lowest values in the dry period/post flush ranging from 11.47 to 13.58 m<sup>-1</sup> (mean = 12.48 m<sup>-1</sup>;  $n = 9$ ), and with the intermediary period in between, with values ranging from 10.41 to 16.39 m<sup>-1</sup> (mean = 13.20 m<sup>-1</sup>;  $n = 9$ ) (Figure 3b).

[12] Spectral slope calculated over the range 275–295 nm ( $S_{275-295}$ ) showed little variation in the Epulu River samples ranging from 0.01533 to 0.01412 nm<sup>-1</sup> (mean = 0.01472 nm<sup>-1</sup>;  $n = 28$ ; Table 1) and no significant difference between samples from different times of year. However,  $S_{350-400}$  ranged from 0.01836 to 0.01352 nm<sup>-1</sup> (mean = 0.01618 nm<sup>-1</sup>;  $n = 28$ ; Table 1) over the year. In contrast to  $S_{275-295}$ ,  $S_{350-400}$  showed seasonal variability with steeper  $S$  values associated with the flushing period ranging from 0.01836 to 0.01707 nm<sup>-1</sup> (mean = 0.01779 nm<sup>-1</sup>;  $n = 10$ ) but shallower in the intermediary period ranging from 0.01716 to 0.01562 nm<sup>-1</sup> (mean = 0.01610 nm<sup>-1</sup>;  $n = 9$ ). The shallowest  $S_{350-400}$  values were observed in the dry period/post flush ranging from 0.01518 to 0.01352 nm<sup>-1</sup> (mean = 0.01448 nm<sup>-1</sup>;  $n = 9$ ). The  $S_R$  in the Epulu River ranged from 0.796 to 1.066 (mean = 0.916;  $n = 28$ ; Table 1 and Figure 4a), with seasonal variations driven by changes in  $S_{350-400}$ . The lowest  $S_R$  values were observed in the flushing period ranging from 0.796 to 0.872 (mean = 0.834;  $n = 10$ ) and the highest values in the dry period/post flush ranging from 0.944 to 1.066 (mean = 1.003;  $n = 9$ ), whereas  $S_R$  ranged from 0.865 to 0.976 (mean = 0.920;  $n = 9$ ) in the intermediary period.

[13]  $SUVA_{254}$  values ranged from 3.08 to 3.57 L mg C<sup>-1</sup> m<sup>-1</sup> (mean = 3.26 L mg C<sup>-1</sup> m<sup>-1</sup>;  $n = 28$ ; Table 1 and Figure 4b). Seasonal variability was observed in  $SUVA_{254}$  with highest values during the flushing period, ranging from 3.29 to 3.57 L mg C<sup>-1</sup> m<sup>-1</sup> (mean = 3.39 L mg C<sup>-1</sup> m<sup>-1</sup>;  $n = 10$ ), whereas the lowest values were observed in the dry period/post flush ranging from 3.08 to 3.20 L mg C<sup>-1</sup> m<sup>-1</sup> (mean = 3.13 L mg C<sup>-1</sup> m<sup>-1</sup>;  $n = 9$ ). In the intermediary period  $SUVA_{254}$  values fell in between the range



**Figure 4.** Box plots of (a) spectral slope ratio  $S_R$ , (b)  $SUVA_{254}$ , and (c) fluorescence index during the flushing, intermediary, and post flush sampling periods. The black dash-dotted line and the solid black line in the box represent the mean and the median, respectively. The horizontal edges of the box denote the 25th and 75th percentiles, and the error bars denote the 10th and 90th percentiles.

shown in the flushing and dry/post flush periods ranging from 3.17 to 3.33  $L\ mg\ C^{-1}\ m^{-1}$  (mean = 3.26  $L\ mg\ C^{-1}\ m^{-1}$ ;  $n = 9$ ).

[14] The fluorescence index (FI) ranged from 1.22 to 1.45 (mean = 1.35;  $n = 28$ ; Table 1 and Figure 4c). As with the parameters derived from CDOM absorbance, the FI also showed seasonal variability in the Epulu River samples. Samples taken during the flushing period had the lowest values ranging from 1.22 to 1.33 (mean = 1.28;  $n = 10$ ) and samples from the dry period/post flush had the highest values ranging from 1.39 to 1.45 (mean = 1.43;  $n = 9$ ). As with the CDOM absorbance-derived parameters, the FI of samples from the intermediary period was within the ranges shown during the flushing and dry/post flush periods ranging from 1.25 to 1.44 (mean = 1.34;  $n = 9$ ).

### 3.3. Lignin Concentrations and Carbon-Normalized Yields

[15] Lignin concentrations ( $\Sigma_8$ ) and carbon-normalized yields ( $\Lambda_8$ ) are presented as the sum of three vanillyl phenols, three syringyl phenols, and two cinnamyl phenols.  $\Sigma_8$  for the Epulu River samples ranged from 41.9 to 131.9  $\mu g\ L^{-1}$  (mean = 77.8  $\mu g\ L^{-1}$ ;  $n = 26$ ; Table 2 and Figure 5a).  $\Lambda_8$  and the carbon-normalized vanillyl yields ( $V$ ) for Epulu River samples ranged from 0.78 to 1.47 ( $mg\ (100\ mg\ OC)^{-1}$ ) (mean = 1.14 ( $mg\ (100\ mg\ OC)^{-1}$ );  $n = 26$ ) and 0.39 to 0.81 ( $mg\ (100\ mg\ OC)^{-1}$ ) (mean = 0.59 ( $mg\ (100\ mg\ OC)^{-1}$ );  $n = 26$ ), respectively (Table 2 and Figures 5b and 5c).  $\Sigma_8$ ,  $\Lambda_8$ , and  $V$  all exhibited temporal variation with highest concentrations or yields in the flushing period (mean = 97.6  $\mu g\ L^{-1}$ , 1.27 ( $mg\ (100\ mg\ OC)^{-1}$ ) and 0.68 ( $mg\ (100\ mg\ OC)^{-1}$ ), respectively), declining in the intermediary period (mean = 76.1  $\mu g\ L^{-1}$ , 1.15 ( $mg\ (100\ mg\ OC)^{-1}$ ) and 0.60 ( $mg\ (100\ mg\ OC)^{-1}$ ), respectively) and with lowest values in the dry period/post flush (mean = 54.6  $\mu g\ L^{-1}$ , 0.96 ( $mg\ (100\ mg\ OC)^{-1}$ ) and 0.46 ( $mg\ (100\ mg\ OC)^{-1}$ ), respectively) (Table 2 and Figure 5).

### 3.4. Lignin Composition

[16] Cinnamyl/vanillyl ( $C/V$ ) and syringyl/vanillyl phenol ( $S/V$ ) ratios in the Epulu River samples ranged from 0.10 to 0.16 (mean = 0.12;  $n = 26$ ) and 0.71 to 0.84 (mean = 0.79;  $n = 26$ ), respectively (Table 2). The ratios of vanillic acid to vanillin ( $Ad/Al$ )<sub>v</sub> and syringic acid to syringaldehyde ( $Ad/Al$ )<sub>s</sub> ranged from 1.08 to 1.17 (mean = 1.13;  $n = 26$ ) and 0.80 to 0.91 (mean = 0.86;  $n = 26$ ), respectively (Table 2). The  $C/V$ ,  $S/V$ , ( $Ad/Al$ )<sub>v</sub>, and ( $Ad/Al$ )<sub>s</sub> ratios did not show any significant temporal trends.

## 4. Discussion

### 4.1. Epulu River Dissolved Organic Matter Temporal Trends

[17] The DOC and dissolved lignin phenol concentrations ( $\Sigma_8$ ) in the Epulu River are high in comparison to other studied rivers (Tables 1 and 2 and Figures 3a and 5a) and highlight the organic-rich nature of the Ituri tropical rainforest ecosystem [Koenig, 2008]. Epulu River DOC concentrations are comparable to previously reported DOC concentrations in the Congo River and its tributaries [Coynel *et al.*, 2005; Spencer *et al.*, 2009a] and in other rivers draining tropical rainforest [Hedges *et al.*, 1994; Battin,


**Table 2.** Epulu River Dissolved Organic Carbon, Lignin Phenol Concentration, Carbon-Normalized Yield, Vanillyl Carbon-Normalized Yield, Cinnamyl/Vanillyl Phenol Ratios, Syringyl/Vanillyl Phenol Ratios, Vanillic Acid/Vanillin Ratios, and Syringic Acid/Syringaldehyde Ratios<sup>a</sup>

Sampling Period	Date	DOC (mg L <sup>-1</sup> )	$\Sigma_8$ ( $\mu\text{g L}^{-1}$ )	$\Lambda_8$ (mg (100 mg OC) <sup>-1</sup> )	V (mg (100 mg OC) <sup>-1</sup> )	C/V	S/V	(Ad/Al) <sub>v</sub>	(Ad/Al) <sub>s</sub>
Wet/flushing	16 Apr 2008	9.0	131.9	1.47	0.81	0.10	0.80	1.11	0.86
Wet/flushing	17 Apr 2008	8.7	125.2	1.44	0.76	0.10	0.79	1.15	0.88
Wet/flushing	18 Apr 2008	8.5	107.8	1.27	0.71	0.10	0.71	1.15	0.87
Wet/flushing	19 Apr 2008	7.6	94.4	1.24	0.66	0.10	0.79	1.14	0.91
Wet/flushing	20 Apr 2008	6.8	83.0	1.21	0.66	0.10	0.76	1.14	0.83
Wet/flushing	21 Apr 2008	6.9	84.5	1.23	0.66	0.10	0.77	1.12	0.84
Wet/flushing	23 Apr 2008	8.0	111.9	1.39	0.72	0.12	0.81	1.12	0.85
Wet/flushing	25 Apr 2008	7.2	91.4	1.27	0.66	0.12	0.81	1.11	0.85
Wet/flushing	26 Apr 2008	6.9	77.3	1.12	0.59	0.12	0.80	1.13	0.86
Wet/flushing	27 Apr 2008	6.6	68.8	1.05	0.60	0.10	0.78	1.09	0.82
Intermediary	05 Feb 2008	7.4	78.6	1.07	0.57	0.10	0.77	1.11	0.82
Intermediary	06 Feb 2008	6.8	90.0	1.32	0.68	0.11	0.84	1.14	0.84
Intermediary	07 Feb 2008	6.0	64.8	1.09	0.57	0.12	0.80	1.08	0.84
Intermediary	08 Feb 2008	5.2	54.1	1.05	0.54	0.14	0.82	1.08	0.83
Intermediary	09 Feb 2008	5.5	51.8	0.94	0.49	0.12	0.79	1.12	0.81
Intermediary	10 Feb 2008	5.4	60.4	1.11	0.57	0.14	0.80	1.10	0.85
Intermediary	11 Feb 2008	7.8	101.6	1.31	0.69	0.11	0.78	1.12	0.88
Intermediary	12 Feb 2008	8.4	107.2	1.28	0.68	0.11	0.76	1.17	0.89
Intermediary	13 Feb 2008	8.0	-	-	-	-	-	-	-
Dry period/post flush	28 Nov 2007	5.3	46.8	0.88	0.45	0.12	0.79	1.16	0.89
Dry period/post flush	29 Nov 2007	6.0	60.5	1.02	0.49	0.15	0.81	1.11	0.88
Dry period/post flush	30 Nov 2007	5.5	56.0	1.02	0.48	0.11	0.78	1.17	0.86
Dry period/post flush	01 Dec 2007	5.3	-	-	-	-	-	-	-
Dry period/post flush	02 Dec 2007	5.4	41.9	0.78	0.39	0.16	0.78	1.12	0.86
Dry period/post flush	03 Dec 2007	5.4	52.0	0.96	0.49	0.11	0.79	1.16	0.80
Dry period/post flush	04 Dec 2007	6.6	66.2	1.00	0.45	0.11	0.80	1.15	0.89
Dry period/post flush	05 Dec 2007	5.6	54.2	0.96	0.44	0.13	0.80	1.11	0.90
Dry period/post flush	06 Dec 2007	5.7	59.3	1.04	0.50	0.13	0.79	1.10	0.85

<sup>a</sup>DOC, dissolved organic carbon;  $\Sigma_8$ , lignin phenol concentration;  $\Lambda_8$ , carbon-normalized yield; V, vanillyl carbon-normalized yield; C/V, cinnamyl/vanillyl phenol ratios; S/V, syringyl/vanillyl phenol ratios; (Ad/Al)<sub>v</sub>, vanillic acid/vanillin ratios; (Ad/Al)<sub>s</sub>, syringic acid/syringaldehyde ratios.

1998]. The mean  $\Sigma_8$  value for the Epulu River of 77.8  $\mu\text{g L}^{-1}$  is comparable to concentrations for the main stem of the Congo River (76.5  $\mu\text{g L}^{-1}$ ) [Spencer et al., 2009a], blackwater tributaries of the Amazon (up to 72  $\mu\text{g L}^{-1}$ ) [Ertel et al., 1986], and the Yukon River at the height of the spring flush (up to 73.5  $\mu\text{g L}^{-1}$ ) [Spencer et al., 2008]. Maximum DOC concentrations during flushing events as observed in the Epulu River have been reported previously in a diverse range of watersheds including other tropical rivers [Newbold et al., 1995; Coynel et al., 2005], temperate rivers [Inamdar et al., 2006; Fellman et al., 2009], and northern high-latitude rivers [Raymond et al., 2007; Striegl et al., 2007]. As with DOC, the highest  $\Sigma_8$  values for the Epulu River occurred during the April flushing period (up to 131.9  $\mu\text{g L}^{-1}$ ; Table 2 and Figure 5a) and are comparable to values from blackwater tributaries of the Yukon River (e.g., Hess Creek, 124.4  $\mu\text{g L}^{-1}$  and Black River, 114.9  $\mu\text{g L}^{-1}$ ) during the spring flush period [Spencer et al., 2008] and storm event samples from a Californian agricultural catchment (107  $\mu\text{g L}^{-1}$ ) [Hernes et al., 2008]. To further assess relative contributions of vascular plant-derived material to organic matter pools, carbon-normalized lignin yields ( $\Lambda_8$ ) are frequently used. The mean  $\Lambda_8$  value of 1.14 (mg (100 mg OC)<sup>-1</sup>) in the Epulu River is high when compared to previously studied rivers, including the main stem of the Congo River (0.72 (mg (100 mg OC)<sup>-1</sup>)) [Spencer et al., 2009a]. This elevated  $\Lambda_8$  is indicative of high vascular plant inputs to the Epulu River.

[18] Carbon-normalized yields based on solely the vanillyl phenols (V) has recently been suggested by Hernes et al. [2007] as a representative end-member for estimation of vascular plant inputs to DOM in riverine systems, as it removes the variation in sources and reactivity associated with syringyl and cinnamyl phenols. In the Epulu River samples, V was typically greater than the combined sum of the carbon-normalized syringyl and cinnamyl phenol yields (Table 2) as has been observed in other rivers [Lobbet et al., 2000; Spencer et al., 2008, 2009a]. The Epulu River mean V yield of 0.59 (mg (100 mg OC)<sup>-1</sup>) is comparable to that of 0.62 (mg (100 mg OC)<sup>-1</sup>) observed in storm events in a Californian agricultural catchment by Hernes et al. [2007] and 0.67 (mg (100 mg OC)<sup>-1</sup>) in high molecular weight (HMW) DOC in the Amazon River [Hedges et al., 2000]. Employing the mean leachate/sorption end-member V yield of 1.53 (mg (100 mg OC)<sup>-1</sup>) from an assorted range of angiosperm and gymnosperm sources [Hernes et al., 2007] and the Epulu River mean V yield, it is estimated that the vascular plant component of DOC is 39% in the Epulu River. Utilizing this end-member V yield for samples from the flushing period (mean V yield = 0.68 (mg (100 mg OC)<sup>-1</sup>)) and dry period/post flush (mean V yield = 0.46 (mg (100 mg OC)<sup>-1</sup>)) results in an estimated vascular plant component of DOC as 44% and 30%, respectively, demonstrating temporal variation in the Epulu River with highest inputs during the flushing period in April. However, the actual vascular plant component of the Epulu River



**Figure 5.** Box plots of (a) lignin phenol concentration ( $\Sigma_8$ ), (b) carbon-normalized yield ( $\Lambda_8$ ), and (c) vanillyl carbon-normalized yield ( $V$ ) during the flushing, intermediary, and post flush sampling periods. The black dash-dotted line and the solid black line in the box represent the mean and the median, respectively. The horizontal edges of the box denote the 25th and 75th percentiles, and the error bars denote the 10th and 90th percentiles.

could be even higher than that estimated here if DOM sources within the catchment generate lower carbon-normalized yields [Spencer *et al.*, 2008], and this seems likely in an organic-rich tropical river such as the Epulu.

[19] The ratio of spectral slopes ( $S_R$ ) was developed as a rapid and easily reproducible method for characterizing CDOM and has been correlated to molecular weight and source, i.e., samples of greater allochthonous character with higher molecular weight DOM have lower  $S_R$  values [Helms *et al.*, 2008]. The range of  $S_R$  in the Epulu River samples (0.796–1.066; Table 1 and Figure 4a) is comparable to that from other allochthonous dominated freshwater systems [Helms *et al.*, 2008; Spencer *et al.*, 2009a, 2009b].  $SUVA_{254}$  has been positively correlated to the percentage of aromaticity of DOM [Weishaar *et al.*, 2003] and the range of values observed in the Epulu River are comparable to the temporal variation observed in other rivers (Table 1 and Figure 4b) [Hood *et al.*, 2005; Spencer *et al.*, 2008; Fellman *et al.*, 2009]. The fluorescence index (FI) changes with aromaticity and indicates the relative contribution of lower molecular weight nonaromatic DOM versus HMW aromatic DOM [McKnight *et al.*, 2001; Cory and McKnight, 2005; Asrat *et al.*, 2007]. For example, fulvic acids with aromaticities of 12%–17% yielded FI values of 1.7–2.0, while fulvic acid samples including Suwannee River with aromaticities of 25%–30% yielded FI values of 1.3–1.4 [McKnight *et al.*, 2001]. FI is conceptually similar to utilization of the emission wavelength of the fulvic-like fluorophore (excitation 300–350 nm, emission 400–460 nm), which has also been linked to DOM hydrophobicity and its functional properties [Baker *et al.*, 2008a] and DOM source [Spencer *et al.*, 2007a]. The range of FI values in the Epulu River (1.22–1.45) are typical of those observed in allochthonous dominated rivers (Table 1 and Figure 4c) [McKnight *et al.*, 2001; Cory *et al.*, 2007]. Fluorescence index was observed to be linearly correlated to  $SUVA_{254}$  and  $S_R$  ( $r^2 = 0.81$ ;  $p < 0.001$  and 0.70;  $p < 0.001$ , respectively;  $n = 28$ ) and  $SUVA_{254}$  and  $S_R$  were linearly correlated to one another ( $r^2 = 0.83$ ;  $p < 0.001$ ;  $n = 28$ ), emphasizing the common drivers of aromaticity and molecular weight underlying these measurements.

[20] The observed temporal variation in Epulu River DOC,  $\Sigma_8$ ,  $\Lambda_8$ , and  $V$  with elevated values in the flushing period, declining in the intermediary period and exhibiting lowest values in the dry period/post flush (Tables 1 and 2 and Figures 3a and 5) is indicative of different DOM source pools, hydrologic flow paths, and residence times between the sampling periods [McGlynn and McDonnell, 2003; Striegl *et al.*, 2005]. The highest riverine DOC,  $\Sigma_8$ ,  $\Lambda_8$ , and  $V$  values are observed during the period of high rainfall after the dry period due to increased surface runoff and leaching of organic rich horizons [Neff *et al.*, 2006; Spencer *et al.*, 2008] during the April flushing period. For example, during this flushing period, the highest  $\Lambda_8$  values (Table 2 and Figure 5b) are comparable to those observed by Hernes *et al.* [2008] (up to 1.6 (mg (100 mg OC) $^{-1}$ )), during storm events in a Californian agricultural catchment concurrent with strong leaching of surface vegetation and organic rich soil layers. Conversely, lower DOC,  $\Sigma_8$ ,  $\Lambda_8$ , and  $V$  values are observed in the dry period/post flush likely due to increased flow path, residence time, and, thus greater microbial mineralization of DOM [Striegl *et al.*, 2005], as well as due to previous extensive flushing of the organic rich



horizons during the preceding wet period (August through October; Figure 2). The extensive flushing of the organic-rich surface layers in the August through October wet period subsequently contributes to lower  $\Lambda_8$  and V yields during the dry period/post flush sampling period, as lignin phenols have been shown to leach at a faster rate than the bulk DOC pool [Spencer *et al.*, 2009b]. The intermediary period shows a large range of DOC,  $\Sigma_8$ ,  $\Lambda_8$ , and V values driven by storm pulses as this period undergoes flushing and drying cycles during the transition out of the dry season. Such variation in the DOM source pools and processing has previously been shown in temperate and northern high-latitude rivers to impact DOC concentration and to cause shifts in the biochemical composition, age, and reactivity of DOM [Holmes *et al.*, 2008; Fellman *et al.*, 2009; Sanderman *et al.*, 2009].

[21] The larger relative terrestrial signature as evidenced by the higher  $\Lambda_8$  and V values in the Epulu River during the April flushing period is supported by the trends in the CDOM data. The lowest  $S_R$ , highest SUVA<sub>254</sub>, and lowest FI values (Table 1 and Figure 4) observed during the post dry flushing period indicate that DOM in this period is enriched in HMW compounds and aromatic moieties compared to other times of year due to the leaching of fresh organic rich layers [Neff *et al.*, 2006; Spencer *et al.*, 2008]. The highest FI and  $S_R$  values and lowest SUVA<sub>254</sub> values were found in the dry period/post flush, indicating that DOM is relatively less aromatic in nature at this time due to a shift in source pools (i.e., this material is already heavily leached from the earlier wet period of August through October), hydrologic flow paths, and residence times between the sampling periods [McGlynn and McDonnell, 2003; Striegl *et al.*, 2005]. Therefore DOM in the Epulu River exhibits temporal variations in its composition, which presumably have a significant impact upon its reactivity in aquatic ecosystems.

#### 4.2. Epulu River Lignin Composition Temporal Trends


[22] The ratios of syringyl (S) and cinnamyl (C) phenols to the ubiquitous vanillyl phenols (V) are used to discriminate between angiosperm and gymnosperm sources (S/V) and between nonwoody and woody tissues (C/V) as syringyl phenols and cinnamyl phenols are specific to angiosperms and nonwoody tissues, respectively [Hedges and Mann, 1979]. In the Epulu River samples, these ratios showed no temporal trends and mean ratios of C/V (0.12) and S/V (0.79) are comparable to those observed in the Congo River [0.15 and 0.68, respectively; Spencer *et al.*, 2009a]. The C/V and S/V ratios in the Epulu River point toward the dominance of angiosperm sources in the region [Hart, 1995; Makana *et al.*, 2004], if no phase change fractionation or degradation has taken place [Hedges and Mann, 1979; Hernes *et al.*, 2007]. The ratios of vanillic acid to vanillin (Ad/Al)<sub>v</sub> and syringic acid to syringaldehyde (Ad/Al)<sub>s</sub> have been shown to have increased relative yields in degradation experiments [Opsahl and Benner, 1995; Hernes and Benner, 2003; Spencer *et al.*, 2009a]. Hernes *et al.* [2007] also showed that the processes of leaching and sorption can result in the fractionation of lignin phenols including the relative increase of (Ad/Al)<sub>v</sub> and (Ad/Al)<sub>s</sub>, thus, these ratios may not be indicative solely of degradation in riverine studies. Epulu River (Ad/Al)<sub>v</sub> and (Ad/Al)<sub>s</sub> ratios showed no

temporal trends and mean ratios of 1.13 and 0.86, respectively. These (Ad/Al)<sub>v</sub> and (Ad/Al)<sub>s</sub> ratios are lower than those observed in the Congo River main stem (1.38 and 1.11, respectively) [Spencer *et al.*, 2009a] and may reflect fresher DOM in the small tributaries with lower residence times and degradation or other sources throughout the Congo Basin with elevated (Ad/Al)<sub>v</sub> and (Ad/Al)<sub>s</sub> ratios. Previous riverine studies employing lignin have shown (Ad/Al)<sub>v</sub> and (Ad/Al)<sub>s</sub> ratios that are comparable to each other from a diverse range of systems [Lobbess *et al.*, 2000; Hedges *et al.*, 2000; Hernes *et al.*, 2008], thus suggesting that (Ad/Al)<sub>v</sub> and (Ad/Al)<sub>s</sub> are primarily governed by common sources and physicochemical processing as opposed to degradation.

#### 4.3. Relationships Between DOC, Lignin, and CDOM

[23] The absorption coefficient of CDOM has previously been utilized as a proxy for DOC in rivers as it is rapid and inexpensive to measure [Worrall *et al.*, 2003; Baker *et al.*, 2008b] and can also be measured in situ to allow for increased spatial and temporal resolution [Spencer *et al.*, 2007c; Downing *et al.*, 2009]. Epulu River  $a_{350}$  values exhibited the same temporal trend as DOC (Figure 3) and a strong linear relationship was found between  $a_{350}$  and DOC ( $r^2 = 0.85$ ;  $p < 0.001$ ;  $n = 28$ ). This further demonstrates the potential of CDOM absorption measurements to derive fluxes of DOC from rivers that have been “ground truthed” [Spencer *et al.*, 2009b].

[24] The utility of molecular biomarkers such as lignin phenols for investigating organic biogeochemical sources and processing has been extensively demonstrated. However, analysis of lignin phenols is expensive and analytically intensive, thus making it poorly suited for the detailed temporal and spatial measurements that are often required to understand complex highly dynamic natural systems. Temporal and spatial coverage of DOM dynamics can be greatly improved by combining CDOM measurements that have been “ground truthed” to molecular biomarkers such as lignin, as has been shown in estuarine [Hernes and Benner, 2003], agricultural river [Hernes *et al.*, 2008], and high-latitude northern river [Spencer *et al.*, 2008, 2009b] systems. In this tropical riverine system,  $a_{350}$  was strongly linearly correlated with  $\Sigma_8$  ( $r^2 = 0.83$ ;  $p < 0.001$ ;  $n = 26$ ), indicating that  $\Sigma_8$  and DOC fluxes (see above) from tropical rivers can be reliably derived from CDOM absorption coefficients. Such measurements will allow for the quantification of fluxes of DOC and  $\Sigma_8$  from CDOM but do not assist in delineating its role in the environment because the spectral parameters of CDOM are largely independent of concentration [Del Vecchio and Blough, 2004; Helms *et al.*, 2008]. In order to “ground truth” CDOM measurements associated with compositional properties, we investigated relationships between lignin carbon-normalized yields ( $\Lambda_8$  and V) and CDOM measurements related to aromaticity and molecular weight (i.e.,  $S_R$ , SUVA<sub>254</sub>, and FI). Linear correlations were observed between  $\Lambda_8$  and  $S_R$ , SUVA<sub>254</sub>, and FI ( $r^2 = 0.69$ , 0.77, and 0.75, respectively;  $p < 0.001$ ; Figure 6). Linear correlations were also observed between V and  $S_R$ , SUVA<sub>254</sub> and FI ( $r^2 = 0.78$ , 0.84, and 0.82, respectively;  $p < 0.001$ ). These relationships indicate that lignin has a fundamental role in CDOM measurements related to aroma-



**Figure 6.** Relationships between lignin carbon-normalized yields ( $\Lambda_8$ ) and CDOM measurements related to aromaticity and molecular weight: (a) spectral slope ratio ( $S_R$ ), (b) SUVA $_{254}$ , and (c) fluorescence index.

ticity, which are used to assess DOM quality. As such these simple and inexpensive optical properties show great potential for future studies aiming to assess the quality and composition of DOM in tropical riverine systems. The

prospective of “ground truthing” CDOM measurements to an array of biomarkers and indicators of DOM reactivity (e.g., lability) serves to increase our fundamental understanding of DOM catchment dynamics [Cory *et al.*, 2007; Fellman *et al.*, 2009].

## 5. Conclusions

[25] To date there has been little work on DOM dynamics in tropical ecosystems, and there is especially a lack of data on temporal variation in DOM quality. Although tropical rainforest systems do not undergo the temporal variability with respect to seasonal climatic extremes observed in temperate and northern high-latitude rivers, this study clearly shows both temporal variation in DOM quantity and more significantly DOM quality. Previous studies in temperate and northern high-latitude rivers have shown the potential to use CDOM-derived optical measurements to examine temporal changes in DOM composition [Hernes *et al.*, 2008; Spencer *et al.*, 2008, 2009b] and to examine lability [Fellman *et al.*, 2009] in riverine systems. Here we show the possibility of deriving DOC and lignin phenol fluxes from CDOM measurements in a tropical river system. This study also shows the ability to track DOM quality by relating lignin carbon-normalized yields (which indicate the contribution of vascular plant-derived material in the DOM pool) to simple optical measurements (e.g.,  $S_R$ , SUVA $_{254}$ , and FI). CDOM-derived measurements, such as  $S_R$  and FI, can be measured in situ and, thus, allow for increased temporal and spatial coverage to elucidate DOM catchment dynamics.

[26] The temporal variability in DOM quality in the Epulu River is attributed to greater surface runoff and leaching of the organic-rich horizons during the April wet period and increased flow path, residence time, and, therefore, greater microbial mineralization in the intermediary period in conjunction with an already well-leached source material in the dry period/post flush [Strigel *et al.*, 2005; Spencer *et al.*, 2008]. This shift in DOM source and processing results in DOM exported during the flushing period with greater HMW and more aromatic character (as indicated by the CDOM quality measurements and the higher  $\Lambda_8$  and V yields) versus DOM exported during the dry period/post flush. In high-latitude northern rivers, DOM exported during the spring flush has a greater proportion derived from vascular plant sources than at other times of the year (i.e., higher  $\Lambda_8$  and V yields as well as relatively lower  $S_R$  and higher SUVA $_{254}$  values [Neff *et al.*, 2006; Spencer *et al.*, 2008, 2009b]) and has also been shown to be younger, more labile, and more photoreactive [Raymond *et al.*, 2007; Holmes *et al.*, 2008; Osburn *et al.*, 2009]. Thus, it seems likely that the Epulu River and other tropical rivers during wet periods are exporting a greater proportion of freshly leached plant material that tends to be highly labile and more susceptible to photochemical degradation. Therefore, temporal variations in DOM quality in tropical rivers will impact the fundamental role the material plays in the receiving ecosystem’s biogeochemistry. Future work is urgently required to better constrain and understand the dynamics and ultimate fate of tropical riverine DOM, which represents >28% of all terrestrial DOC inputs to the oceans [Coynel *et al.*, 2005].

[27] **Acknowledgments.** We would like to thank Cort Anastasio for use of the Shimadzu UV-2501PC UV/Vis spectrophotometer and Tad Doane, Sarah Flores, and Tim Ingrum for their assistance. We thank Kevin Burkhill at the University of Birmingham for producing Figure 1. Finally, R.G.M.S. is very grateful to Major Paul Naish (RET. South African Army) of African Byways for sharing information garnered from his antipoaching intelligence/information gathering work in the Democratic Republic of Congo and his assistance in getting this project off the ground.

## References

- Ahmad, S. R., and D. M. Reynolds (1999), Monitoring of water quality using fluorescence technique: prospect of online process control, *Water Res.*, **33**, 2069–2074.
- Aitkenhead, J. A., and W. H. McDowell (2000), Soil C:N ratio as a predictor of annual riverine DOC flux at local and global scales, *Global Biogeochem. Cycles*, **14**(1), 127–138.
- Asrat, A., A. Baker, M. U. Mohammed, M. J. Leng, P. V. Calsteren, and C. Smith (2007), A high-resolution multi-proxy stalagmite record from Mechara, Southeastern Ethiopia: palaeohydrological implications for speleothem palaeoclimate reconstruction, *J. Quaternary Sci.*, **22**, 53–63.
- Baccini, A., N. Laporte, S. J. Goetz, M. Sun, and H. Dong (2008), A first map of tropical Africa's above-ground biomass derived from satellite imagery, *Environ. Res. Lett.*, **3**, 045011.
- Baker, A., E. Tipping, S. A. Thacker, and D. Gondar. (2008a), Relating dissolved organic matter fluorescence and functional properties, *Chemosphere*, **73**, 1765–1772.
- Baker, A., L. Bolton, M. Newson, and R. G. M. Spencer (2008b), Spectrophotometric properties of surface water dissolved organic matter in an afforested upland peat catchment, *Hydrocarbon Process.*, **22**, 2325–2336.
- Battin, T. J. (1998), Dissolved organic matter and its optical properties in a blackwater tributary of the upper Orinoco River, Venezuela, *Org. Geochem.*, **28**, 561–569.
- Cory, R. M., and D. M. McKnight (2005), Fluorescence spectroscopy reveals ubiquitous presence of oxidized and reduced quinones in dissolved organic matter, *Environ. Sci. Technol.*, **39**, 8142–8149.
- Cory, R. M., D. M. McKnight, Y. P. Chin, P. Miller, and C. L. Jaros (2007), Chemical characteristics of fulvic acids from Arctic surface waters: Microbial contributions and photochemical transformations, *J. Geophys. Res.*, **112**, G04S51, doi:10.1029/2006JG000343.
- Coyne, A., P. Seyler, H. Etcheber, M. Meybeck, and D. Orange (2005), Spatial and seasonal dynamics of total suspended sediment and organic carbon species in the Congo River, *Global Biogeochem. Cycles*, **19**(4), GB4019, doi:10.1029/2004GB002335.
- Dalzell, B. J., T. R. Filley, and J. M. Harbor (2005), Flood pulse influences on terrestrial organic matter export from an agricultural watershed, *J. Geophys. Res.*, **110**(G2), G02011, doi:10.1029/2005JG000043.
- Del Vecchio, R., and N. V. Blough (2004), On the origin of the optical properties of humic substances, *Environ. Sci. Technol.*, **38**, 3885–3891.
- Downing, B. D., E. Boss, B. A. Bergamaschi, J. A. Fleck, M. A. Lionberger, N. K. Ganju, D. H. Schoellhamer, and R. Fujii (2009), Quantifying fluxes and characterizing compositional changes of dissolved organic matter in aquatic systems in situ using combined acoustic and optical measurements, *Limnol. Oceanogr. Methods*, **7**, 119–131.
- Duveiller, G., P. Defourny, B. Desclee, and P. Mayaux (2008), Deforestation in central Africa: Estimates at regional, national and landscape levels by advanced processing of systematically distributed Landsat extracts, *Remote Sens. Environ.*, **112**, 1969–1981.
- Ertel, J. R., J. I. Hedges, A. H. Devol, J. E. Richey, and M. D. G. Ribeiro (1986), Dissolved humic substances of the Amazon River system, *Limnol. Oceanogr.*, **31**, 739–754.
- Evans, C. D., C. Freeman, L. G. Cork, D. N. Thomas, B. Reynolds, M. F. Billett, M. H. Garnett, and D. Norris (2007), Evidence against recent climate-induced destabilisation of soil carbon from  $^{14}\text{C}$  analysis of riverine dissolved organic matter, *Geophys. Res. Lett.*, **34**(7), L07407, doi:10.1029/2007GL029431.
- Fellman, J. B., E. Hood, R. T. Edwards, and D. V. D'Amore (2009), Changes in the concentration, biodegradability, and fluorescent properties of dissolved organic matter during stormflows in coastal temperate watersheds, *J. Geophys. Res.*, **114**, G01021, doi:10.1029/2008JG000790.
- Hart, T. B. (1995), Seed, seedling and subcanopy survival in monodominant and mixed forests of the Ituri Forest, Africa, *J. Trop. Ecol.*, **11**, 443–459.
- Hedges, J. I., and J. R. Ertel (1982), Characterization of lignin by gas capillary chromatography of cupric oxide oxidation-products, *Anal. Chem.*, **54**, 174–178.
- Hedges, J. I., and D. C. Mann (1979), Characterization of plant tissues by their lignin oxidation products, *Geochim. Cosmochim. Acta*, **43**, 1803–1807.
- Hedges, J. I., G. L. Cowie, J. E. Richey, P. D. Quay, R. Benner, M. Strom, and B. R. Forsberg (1994), Origins and processing of organic matter in the Amazon River as indicated by carbohydrates and amino acids, *Limnol. Oceanogr.*, **39**, 743–761.
- Hedges, J. I., et al. (2000), Organic matter in Bolivian tributaries of the Amazon River: A comparison to the lower main stream, *Limnol. Oceanogr.*, **45**, 1449–1466.
- Helms, J. R., A. Stubbins, J. D. Ritchie, E. C. Minor, D. J. Kieber, and K. Mopper (2008), Absorption spectral slopes, and slope ratios as indicators of molecular weight, source, and photobleaching of chromophoric dissolved organic matter, *Limnol. Oceanogr.*, **53**, 955–969.
- Hernes, P. J., and R. Benner (2002), Transport and diagenesis of dissolved and particulate terrigenous organic matter in the North Pacific Ocean, *Deep Sea Res. Part 1*, **49**(12), 2119–2132.
- Hernes, P. J., and R. Benner (2003), Photochemical and microbial degradation of dissolved lignin phenols: Implications for the fate of terrigenous dissolved organic matter in marine environments, *J. Geophys. Res.*, **108**(C9), 3291, doi:10.1029/2002JC001421.
- Hernes, P. J., A. C. Robinson, and A. K. Aufdenkampe (2007), Fractionation of lignin during leaching and sorption and implications for organic matter “freshness,” *Geophys. Res. Lett.*, **34**(17), L17401, doi:10.1029/2007GL031017.
- Hernes, P. J., R. G. M. Spencer, R. Y. Dyda, B. A. Pellerin, P. A. M. Bachand, and B. A. Bergamaschi (2008), The role of hydrologic regimes on dissolved organic carbon composition in an agricultural watershed, *Geochim. Cosmochim. Acta*, **72**, 5266–5277.
- Holmes, R. M., J. W. McClelland, P. A. Raymond, B. B. Frazer, B. J. Peterson, and M. Stieglitz (2008), Lability of DOC transported by Alaskan rivers to the Arctic Ocean, *Geophys. Res. Lett.*, **35**(3), L03402, doi:10.1029/2007GL032837.
- Hood, E., M. W. Williams, and D. M. McKnight (2005), Sources of dissolved organic matter (DOM) in a Rocky Mountain stream using chemical fractionation and stable isotopes, *Biogeochem.*, **74**, 231–255.
- Hood, E., J. Fellman, and R. T. Edwards (2007), Salmon influences on dissolved organic matter in a coastal brown-water stream: An application of fluorescence spectroscopy, *Limnol. Oceanogr.*, **52**, 1580–1587.
- Hu, C. M., F. E. Muller-Karger, and R. G. Zepp (2002), Absorbance, absorption coefficient, and apparent quantum yield: A comment on common ambiguity in the use of these optical concepts, *Limnol. Oceanogr.*, **47**(4), 1261–1267.
- Inamdar, S. P., N. O'Leary, M. J. Mitchell, and J. T. Riley (2006), The impact of storm events on solute exports from a glaciated forested watershed in western New York, USA, *Hydrol. Processes*, **20**, 3423–3439.
- Katuala, P. G. B., J. A. Hart, R. Hutterer, H. Leirs, and A. Dudu (2005), Biodiversity and ecology of small rodents (Rodents and Shrews) of the “Reserve de Faune a Okapis,” Democratic Republic of the Congo, *Belg. J. Zool.*, **135**, 191–196.
- Koenig, R. (2008), Critical time for African rainforests, *Science*, **320**, 1439–1441.
- Laporte, N. T., J. A. Stabach, R. Grosch, T. S. Lin, and S. J. Goetz (2007), Expansion of industrial logging in central Africa, *Science*, **316**, 1451–1451.
- Lobbis, J. M., H. P. Fitznar, and G. Kattner (2000), Biogeochemical characteristics of dissolved and particulate organic matter in Russian rivers entering the Arctic Ocean, *Geochim. Cosmochim. Acta*, **64**, 2973–2983.
- Makana, J. R., T. B. Hart, I. Liengola, C. Ewango, J. A. Hart, and R. Condit (2004), Ituri Forest Dynamics Plots, Democratic Republic of Congo, in *Tropical Forest Diversity and Dynamism: Findings From a Large-Scale Plot Network*, edited by E. C. Losos and E. G. Leigh, pp. 492–505, Univ. of Chicago Press.
- McGlynn, B. L., and J. J. McDonnell (2003), Role of discrete landscape units in controlling catchment dissolved organic carbon dynamics, *Water Resour. Res.*, **39**(4), 1090, doi:10.1029/2002WR001525.
- McKnight, D. M., E. W. Boyer, P. K. Westerhoff, P. T. Doran, T. Kulbe, and D. T. Andersen (2001), Spectrofluorometric characterization of dissolved organic matter for indication of precursor organic material and aromaticity, *Limnol. Oceanogr.*, **46**, 38–48.
- Neff, J. C., J. C. Finlay, S. A. Zimov, S. P. Davydov, J. J. Carrasco, E. A. G. Schuur, and A. I. Davydova (2006), Seasonal changes in the age and structure of dissolved organic carbon in Siberian rivers and streams, *Geophys. Res. Lett.*, **33**(23), L23401, doi:10.1029/2006GL028222.
- Newbold, J. D., B. W. Sweeney, J. K. Jackson, and L. A. Kaplan (1995), Concentrations and export of solutes from 6 mountain streams in northwestern Costa Rica, *J. N. Am. Benthol. Soc.*, **14**, 21–37.
- Opsahl, S., and R. Benner (1995), Early diagenesis of vascular plant tissues, lignin and cutin decomposition and biogeochemical implications, *Geochim. Cosmochim. Acta.*, **59**(23), 4889–4904.

- Osburn, C. L., L. Retamal, and W. F. Vincent (2009), Photoreactivity of chromophoric dissolved organic matter transported by the Mackenzie River to the Beaufort Sea, *Mar. Chem.*, *115*, 10–20.
- Raymond, P. A., J. W. McClelland, R. M. Holmes, A. V. Zhulidov, K. Mull, B. J. Peterson, R. G. Striegl, G. R. Aiken, and T. Y. Gurtovaya (2007), Flux and age of dissolved organic carbon exported to the Arctic Ocean: A carbon isotopic study of the five largest rivers, *Global Biogeochem. Cycles*, *21*(4), GB4011, doi:10.1029/2007GB002934.
- Sanderman, J., K. A. Lohse, J. A. Baldock, and R. Amundson (2009), Linking soils and streams: Sources and chemistry of dissolved organic matter in a small coastal watershed, *Water Resour. Res.*, *45*(3), W03418, doi:10.1029/2008WR006977.
- Spencer, R. G. M., A. Baker, J. M. E. Ahad, G. L. Cowie, R. Ganeshram, R. C. Upstill-Goddard, and G. Uher (2007a), Discriminatory classification of natural and anthropogenic waters in two U.K. estuaries, *Sci. Total Environ.*, *373*, 305–323.
- Spencer, R. G. M., L. Bolton, and A. Baker (2007b), Freeze/thaw and pH effects on freshwater dissolved organic matter fluorescence and absorbance properties from a number of U.K. locations, *Water Res.*, *41*, 2941–2950.
- Spencer, R. G. M., B. A. Pellerin, B. A. Bergamaschi, B. D. Downing, T. E. C. Kraus, D. R. Smart, R. A. Dahlgren, and P. J. Hernes (2007c), Diurnal variability in riverine dissolved organic matter composition determined by in-situ optical measurement in the San Joaquin River (California, U.S.A.), *Hydrol. Processes*, *21*, 3181–3189.
- Spencer, R. G. M., G. R. Aiken, K. P. Wickland, R. G. Striegl, and P. J. Hernes (2008), Seasonal and spatial variability in dissolved organic matter quantity and composition from the Yukon River basin, Alaska, *Global Biogeochem. Cycles*, *22*(4), GB4002, doi:10.1029/2008GB003231.
- Spencer, R. G. M., et al. (2009a), Photochemical degradation of dissolved organic matter and dissolved lignin phenols from the Congo River, *J. Geophys. Res.*, *114*, G03010, doi:10.1029/2009JG000968.
- Spencer, R. G. M., G. R. Aiken, K. D. Butler, M. M. Dornblaser, R. G. Striegl, and P. J. Hernes (2009b), Utilizing chromophoric dissolved organic matter measurements to derive export and reactivity of dissolved organic carbon to the Arctic Ocean: A case study of the Yukon River, Alaska, *Geophys. Res. Lett.*, *36*(6), L06401, doi:10.1029/2008GL036831.
- Striegl, R. G., G. R. Aiken, M. M. Dornblaser, P. A. Raymond, and K. P. Wickland (2005), A decrease in discharge-normalized DOC export by the Yukon River during summer through autumn, *Geophys. Res. Lett.*, *32*(21), L21413, doi:10.1029/2005GL024413.
- Striegl, R. G., M. M. Dornblaser, G. R. Aiken, K. P. Wickland, and P. A. Raymond (2007), Carbon export and cycling by the Yukon, Tanana, and Porcupine rivers, Alaska, 2001–2005, *Water Resour. Res.*, *43*(2), W02411, doi:10.1029/2006WR005201.
- Stubbins, A., V. Hubbard, G. Uher, C. S. Law, R. C. Upstill-Goddard, G. R. Aiken, and K. Mopper (2008), Relating carbon monoxide photo-production to dissolved organic matter functionality, *Environ. Sci. Tech.*, *42*, 3271–3276.
- Weishaar, J. L., G. R. Aiken, B. A. Bergamaschi, M. S. Fram, R. Fuji, and K. Mopper (2003), Evaluation of specific ultraviolet absorbance as an indicator of the chemical composition and reactivity of dissolved organic carbon, *Environ. Sci. Technol.*, *37*, 4702–4708.
- Worrall, F., T. Burt, and R. Shedden (2003), Long term records of riverine dissolved organic matter, *Biogeochem.*, *64*, 165–178.
- 
- A. Baker, Connected Waters Initiative, Water Research Laboratory, University of New South Wales, 110 King Street, Manly Vale, NSW, 2093, Australia.
- R. Y. Dyda, J. Six, and R. G. M. Spencer, Department of Plant Sciences, University of California–Davis, One Shields Avenue, Davis, CA 95616, USA. (rgspencer@ucdavis.edu)
- P. J. Hernes, Department of Land, Air and Water Resources, University of California–Davis, One Shields Avenue, Davis, CA 95616, USA.
- R. Ruf, The Okapi Breeding and Research Station, Institute in Congo for the Conservation of Nature, Gilman International Conservation, Epulu, Orientale Province, Democratic Republic of Congo.
- A. Stubbins, Department of Chemistry and Biochemistry, Old Dominion University, Norfolk, VA 23529, USA.